

# Journal of Statistical Software

MMMMMM YYYY, Volume VV, Issue II.

doi: 10.18637/jss.v000.i00

# The chords R Package- A Principled Approach to Respondent Driven Sampling

Jonathan D. Rosenblatt

Ben Gurion University

Yakir Brechenko Gertner Institute for

Epidemiology and Health Policy Research Simon D. Frost Cambridge University

Abstract

The abstract of the article.

Keywords: RDS, estimation, counting-process, R.

# 1. Introduction

As the name suggests, Respondent Driven Sampling (RDS) is a framework for sampling by chain-referral. RDS is a bundle of a sampling mechanism and analysis methods, most common in the study of marginalized populations which do not lend themselves to simple sampling (Heckathorn 1997, 2002).

In RDS seeds are selected – usually by convenience – from the target population, and given coupons. They use these coupons to recruit others, who themselves become recruiters. Recruits are given an incentive, usually money, for taking part in the survey, and also for recruiting others. This process continues in recruitment waves until the survey is stopped, usually when a target sample size is reached.

With the above sampling mechanism, highly connected individuals will be overrepresented in the sample. If the attribute of interest is correlated with an individual's degree, as is often the case (e.g. HIV), naïveestimates will be biased towards the state of the highly connected subgroups. An unbiased Horowitz-Thompson-type estimator (Horvitz and Thompson 1952) would require the knowledge of frequency of each degree. Clearly, the frequency of each degree will also be biased towards higher degrees, and thus cannot be recovered from the knowledge of individuals' degrees. A common

# 2. Work Flow

# 3. Some Technicalities

# 4. Conclusion

#### 5. Future Work

### References

Heckathorn D (1997). "Respondent-driven sampling: a new approach to the study of hidden populations." *Social Problems*, **44**(2), 174–199.

Heckathorn D (2002). "Respondent-driven sampling II: deriving valid population estimates from chain-referral samples of hidden populations." Social Problems, 49(1), 11–34.

Horvitz DG, Thompson DJ (1952). "A generalization of sampling without replacement from a finite universe." *Journal of the American Statistical Association*, **47**(260), 663–685.

http://www.jstatsoft.org/

http://www.foastat.org/

Submitted: yyyy-mm-dd

Accepted: yyyy-mm-dd

# Affiliation:

Jonathan D. Rosenblatt
Department of Industrial Engineering and Management
Faculty of Engineering
Ben Gurion University of the Negev
P.O. 653, Beer Sheva, 8410501
E-mail: johnros@bgu.ac.il

URL: http://www.john-ros.com/

Journal of Statistical Software published by the Foundation for Open Access Statistics MMMMMM YYYY, Volume VV, Issue II doi:10.18637/jss.v000.i00