

***METHODS SECTION OF THE PAPER ENTITLED: Acute effects of inter-effort recovery hypoxia on physiological responses and training quality during sprint interval training and small-sided games in handball: a randomised crossover trial***

**## Disclaimer**

This file (‘methods.pdf’) is provided **\*\*for consultation purposes only\*\***.

It must **\*\*not\*\*** be cited, reproduced, or used as part of the final manuscript.

Its sole purpose is to assist in understanding and interpreting the analysis scripts in this repository.

The related article is currently under peer review. If accepted and published, the official reference will be available on ResearchGate, at the link below.

**## Author**

***Germano Marcolino Putti***

***GitHub: [germanomputti](https://github.com/germanomputti)***

***ORCID: https://orcid.org/0000-0001-8191-598X***

***Research gate: https://www.researchgate.net/profile/Germano-Putti-2?ev=hdr\_xprf***

***Email: germanomputti@gmail.com***

***Lattes: http://lattes.cnpq.br/9960140114038219***


**1. Methods**

**1.1.1. Experimental Design**

To investigate the effects of IEH during SIT and SSG on physiological indicators and training quality in handball, a randomised, crossover, single-blind study was conducted. Participants completed four training sessions, two SIT and two SSG, under environmental conditions of normoxia (NOR, FiO<sub>2</sub> = 0.21, SIT-NOR and SSG-NOR) or IEH (FiO<sub>2</sub> = 0.13, SIT-IEH and SSG-IEH). The training sessions were filmed and the distances covered (SIT) or each athlete's score in PlayerScore (Wagner et al., 2023a) (SSG) were determined. In addition, lactate concentrations, heart rate and perceptual responses were monitored as indicators of

physiological stress during each session. Figure 1 illustrates experimental design and training protocol. The sessions were held in August 2024, in the morning, on a 40x20m court located at an altitude of 546m above sea level ( $FiO_2=0.21$ ).

Figure 1 – Experimental design and training session



The SIT sessions took place before the SSG sessions, with only the environmental condition (NOR or IEH) being randomised. SIT: sprint interval training; SSG: small-sided games; NOR: normoxia; IEH: inter-effort recovery hypoxia;  $FiO_2$ : fraction of inspired oxygen.

### 1.1.2. Participants

Ten handball players voluntarily took part in this study. The characteristics of the participants are shown in Table 1, with body composition values obtained using the dual-energy X-ray absorptiometry technique (Lunar DPX-NT, General Electric Healthcare, Buckinghamshire, Reino Unido). The participants' fitness level were classified according to criteria previously proposed by Mackay et al (2022) as Tier 2: Trained/Developmental. The study's inclusion criteria included male handball players aged between 18 and 35; while the exclusion criteria consisted of not completing one or more of the scheduled sessions (SIT-NOR, SIT-IEH, SSG-NOR, SSG-IEH). Two participants did not complete all sessions and were removed from final analysis. The participant flow chart is shown in Figure 2. All participants


were informed about the risks and benefits of the study, and each provided their voluntary agreement by signing a document in accordance with the Declaration of Helsinki and approved by the university's ethics committee.

Table 1 – Participants' characteristics

Height (cm)	Weight (kg)	Age (year)	Lean mass (kg)	Fat percentage (%)
177.1 ± 4.1	78.6 ± 9.0	22.4 ± 3.3	61.5 ± 7.0	18.3 ± 5.3

Values are mean ± standard deviation. N=8.

Figure 2 – Participants flow diagram



SIT: sprint interval training; SSG: small-sided games; NOR: normoxia; IEH: inter-effort recovery hypoxia.

### 1.1.3. Hypoxic exposure

A balloon system previously proposed and validated by Norberto et al. (2024) was used to administer hypoxia exposure during the training sessions, which allows the storage of

air with low FiO<sub>2</sub>, obtained by using a commercial hypoxia generator (ACT – 12 air unit, Everest Summit II, Cardiff, EUA). Unidirectional hoses and masks were connected to this system, allowing participants to inhale hypoxic air and exhale air into the environment. To ensure blinding, a damper system was implemented which allowed the air flow to be redirected so that the participants in the normoxic situation inhaled atmospheric air. Only the researchers were aware of the participants' experimental conditions. After a standardized warm-up of approximately 10 minutes including mobility exercises, ballistic stretching, squats, jumps, running drills, and short sprints, the participants remained breathing the air corresponding to the randomised condition for 5 minutes and then began their first effort. Between efforts, the recovery and exposure time was 4 minutes (4x) and after the last effort another 5 minutes of passive exposure, for a total of 26 minutes of exposure (Figure 1).

#### ***1.1.4. Training protocol***

The participants included in the study underwent four training sessions, the first two being SIT and the last two SSG, in a non-randomised way. This format was chosen to avoid a dropout after the SSG sessions had been carried out – if this were to happen, it would be necessary to repeat the session with all the participants. Only the environmental conditions were randomised. Randomisation was carried out by an online drawer. The sessions were carried out at least 48 hours apart. In addition, participants were instructed not to perform intense physical exercise in the 24 hours prior to the study protocol sessions.

##### ***1.1.4.1. Sprint interval training (SIT)***

The SIT sessions consisted of five maximal running efforts lasting 30 seconds, with a 4-minute break between each effort. The efforts were made between the side lines of the court, with a 180° change of direction every 20 metres. The assessors verbally encouraged the participants to make a maximum effort throughout the training session. The sessions were filmed and then the distances covered in each of the 5 efforts were determined, with a precision of 2.5 metres ( $\pm 1.8\%$  of the average value of all the sprints). In addition to the distance values, the average power output (PO) during the effort was also estimated using the formula proposed by Kalva-Filho et al. (2013):

$$\text{Eq 1: Power output (W)} = \frac{\text{Body weight (kg)} * [\text{distance (m)}]^2}{[\text{time (s)}]^3}$$

It's important to note that, due to the time spent changing direction and the consequent decrease in speed and distance travelled during the 30 s, this estimated PO is underestimated. Even so, it was decided to use it as a value that somehow includes the external load that had to

be moved to make the effort (body mass) and the effort itself (the distance travelled). PO was analysed as maximum (PO in the longest sprint,  $PO_{MAX}$ ), mean (average of all sprints,  $PO_{MEAN}$ ) and fatigue index:

$$\text{Eq 2: Fatigue index(\%)} = 100 - \left( \frac{PO_{MEAN}}{PO_{MAX}} \right) * 100$$

#### **1.1.4.2. Small-sided games (SSG)**

The sessions with SSG consisted of five games lasting 2.5 min with a 4 min break between each game, on an official court (40 × 20 m). Each team consisted of four court players plus a goalkeeper (who was not exposed to hypoxia at any time). Two rules were changed in order to increase the intensity of the session: dribbling was not allowed and the ball was replaced directly by the goalkeeper, even after a goal (Buchheit et al., 2009). To ensure that the technical level of the teams was balanced, a coach who knew all the players divided them into pairs of a similar level and then they were randomised within this pair. The teams were maintained for both games.

To assess performance during the SSG, the sessions were filmed and then analysed using the PlayerScore observation tool (Wagner et al., 2023a). This systematic observation tool makes it possible to classify different actions in the offensive and defensive phases of the game, with positive and negative scores based on how much a given action contributes to or impedes the team in scoring a goal or preventing a goal.

#### **1.1.5. Internal load**

##### **1.1.5.1. Heart rate**

Heart rate values were monitored during the SSG and SIT sessions. The values were recorded immediately after the efforts and every 10 s during all the recoveries. The values were analysed as the maximum value (immediately after the effort,  $HR_{MAX}$ ) during the session and the sum of the area under the curve ( $HR_{AUC}$ ) (Malta et al., 2023) during recovery bouts.

##### **1.1.5.2. Blood lactate**

Before warming up and one minute after the end of each effort, 25 µL of blood was collected from the earlobe in a previously heparinised glass capillary and stored in a tube containing 50 µL of 1% NaF. The samples were frozen at -20 °C and the lactate concentration ([La]) was subsequently estimated on a biochemical analyser calibrated according to the manufacturer's instructions (YSI 2700 Biochemistry Analyzer, Yellow Springs Instrument Company, Ohio, EUA). The [La] values were then analysed as the difference between the resting value and the maximum value obtained during the session ( $\Delta$  [La]).

### **1.1.5.3. Perceptual responses**

A modified Borg scale (0-10) was used to monitor perceptual responses. The four responses monitored were: the rate of perceived effort (*How much effort did you give compared to a maximal effort?*), the perception of overall discomfort (*How uncomfortable do you feel overall?*), the perception of difficulty breathing (*How uncomfortable does it feel to breathe?*), and the perception of discomfort in the lower limbs (*How uncomfortable do your legs feel?*) (Christian et al., 2014).

### **1.1.6. Data analysis**

The data was analysed using generalised linear mixed models, with fixed effects for training modality (SSG or SIT) and environmental condition (IEH or NOR) and random effects for individuals. For the variables relating to distance or PO in SIT or PlayerScore in SSG, the only fixed effect was the environmental condition. Before fitting the models, the presence of outliers was checked using box and scatter plots. The Gamma and Gaussian distributions were tested, and the assumptions of normality of the residuals were checked by visually inspecting the histogram and the QQplot, as well as homoscedasticity by analysing the Pearson residual plot as a function of the predicted values. If the assumptions of the models were met, the final model adopted would be the one with the lowest Akaike information criterion (AIC). Based on the model selected, hypothesis tests were carried out on the coefficients, with Bonferroni correction, adopting a significance level of 5%.

The data is presented as estimated mean or mean difference and 95% confidence interval. Cohen's d for paired samples was used to estimate the effect size, calculated as the quotient of the difference between the means and the product of the standard error of the difference and the square root of the sample size. The qualitative interpretation of the effect size was: small (0.0-0.2), medium (0.2-0.5), large (0.5-0.8) and very large (>0.8). The analyses were conducted in the R environment (R Core Team, 2024), with the following packages: readxl (Wickham & Bryan, 2023), dplyr (Wickham et al., 2023), writexl (Ooms, 2024), lme4 (Bates et al., 2015), ggResidpanel (Goode & Rey, 2019), ggplot2 (Wickham, 2016), lmerTest (Kuznetsova et al., 2017), and emmeans (Lenth, 2024).

## REFERENCES

- Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models Using lme4. *Journal of Statistical Software*, 67(1). <https://doi.org/10.18637/jss.v067.i01>
- Buchheit, M., Laursen, P., Kuhnle, J., Ruch, D., Renaud, C., & Ahmaidi, S. (2009). Game-based Training in Young Elite Handball Players. *International Journal of Sports Medicine*, 30(04), 251–258. <https://doi.org/10.1055/s-0028-1105943>
- Christian, R. J., Bishop, D. J., Billaut, F., & Girard, O. (2014). The role of sense of effort on self-selected cycling power output. *Frontiers in Physiology*, 5. <https://doi.org/10.3389/fphys.2014.00115>
- Goode, K., & Rey, K. (2019). *ggResidpanel: Panels and Interactive Versions of Diagnostic Plots using “ggplot2”* [Computer software]. R package (<https://CRAN.R-project.org/package=ggResidpanel>).
- Kalva-Filho, C. A., Loures, J. P., Franco, V. H., Kaminagakura, E. I., Zagatto, A. M., & Papoti, M. (2013). Comparação da potência anaeróbia mensurada pelo teste de RAST em diferentes condições de calçado e superfícies. *Revista Brasileira de Medicina Do Esporte*, 19(2), 139–142. <https://doi.org/10.1590/S1517-86922013000200014>
- Kuznetsova, A., Brockhoff, P. B., & Christensen, R. H. B. (2017). lmerTest Package: Tests in Linear Mixed Effects Models. *Journal of Statistical Software*, 82(13). <https://doi.org/10.18637/jss.v082.i13>
- Lenth, R. (2024). *emmeans: Estimated Marginal Means, aka Least-Squares Means* [Computer software]. R package (<https://CRAN.R-project.org/package=emmeans>).
- Malta, E. S., Lopes, V. H. F., Esco, M. R., & Zagatto, A. M. (2023). Repeated cold-water immersion improves autonomic cardiac modulation following five sessions of high-

- intensity interval exercise. *European Journal of Applied Physiology*, 123(9), 1939–1948. <https://doi.org/10.1007/s00421-023-05205-4>
- McKay, A. K. A., Stellingwerff, T., Smith, E. S., Martin, D. T., Mujika, I., Goosey-Tolfrey, V. L., Sheppard, J., & Burke, L. M. (2022). Defining Training and Performance Caliber: A Participant Classification Framework. *International Journal of Sports Physiology and Performance*, 17(2), 317–331. <https://doi.org/10.1123/ijsp.2021-0451>
- Norberto, M. S., Torini, J. V. G., Firmino, M. S., & Papoti, M. (2024). Validation of Air Storage System for Hypoxia Exposure During Exercise. *High Altitude Medicine & Biology*, 25(2), 122–128. <https://doi.org/10.1089/ham.2023.0122>
- Ooms, J. (2024). *writexl: Export Data Frames to Excel “xlsx” Format* (Version 1.5.0) [Computer software]. <https://CRAN.R-project.org/package=writexl>
- R Core Team. (2024). *R: A Language and Environment for Statistical Computing* [Computer software]. R Foundation for Statistical Computing <<https://www.R-project.org/>>.
- Wagner, H., Hinz, M., Melcher, K., Radic, V., & Uhrmeister, J. (2023a). The PlayerScore: A Systematic Game Observation Tool to Determine Individual Player Performance in Team Handball Competition. *Applied Sciences*, 13(4), 2327. <https://doi.org/10.3390/app13042327>
- Wagner, H., Hinz, M., Melcher, K., Radic, V., & Uhrmeister, J. (2023b). The PlayerScore: A Systematic Game Observation Tool to Determine Individual Player Performance in Team Handball Competition. *Applied Sciences*, 13(4), 2327. <https://doi.org/10.3390/app13042327>
- Wickham, H. (2016). *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York. <https://ggplot2.tidyverse.org>
- Wickham, H., & Bryan, J. (2023). *readxl: Read Excel Files* (Version 1.4.3) [Computer software]. <https://CRAN.R-project.org/package=readxl>



Wickham, H., François, R., Henry, L., Müller, K., & Vaughan, D. (2023). *dplyr: A Grammar of Data Manipulation* (Version 1.1.4) [Computer software]. <https://CRAN.R-project.org/package=dplyr>