PHYS 615 – HW 10

Types of homework questions

- RQ (Reading questions): prompt you to go back to the text and read and think about the text more carefully and explain in your own words. While not directly tested in quizzes, can help you think more deeply about quiz questions.
- BF (Building foundations): gives you an opportunity to build and practice foundational skills that you have, presumably, seen before.
- TQQ (typical quiz questions): Similar questions (though perhaps longer or shorter) will be asked on quizzes. But the difficulty level and skills tested will be similar.
- Design (D): These are questions in which you are given a desired outcome and asked to figure out how to make it happen. These will often also be TQQ's, but always starting with desired motion/behavior as the given.
- COMP (Computing): computing questions often related to TQQ but will never be asked on a quiz (since debugging can take so long). You will need to do at least four computing questions over the semester
- FC (free choice): allows you to decide where to put your time. Any of the following are possible: work through a section of the text or a lecture in detail; redo a problem from before; do an unassigned problem in the text; extend a computing project; try a problem using a different analytical approach (e.g. forces instead of conservation of energy).
- ACT (in-class activity): These questions are repeats of questions (or similar to) that occurred in a previous in-class activity.
- Standard Reading Questions: How does the reading connect with what you already know? What was something new? Ask an "I wonder" question OR give an example applying the idea in the reading.

Please remember to say something about the "Check/Learn" part at the end of solving a problem!

Full credit will be given at 75% of the total points possible, so you can choose a subset of problems (you can do more / all, but the score is capped at 75%)

This homework contains some previous group activities. I'm including them here in order to try to help gradescope, but you can of course hand in the original paper version I handed out in class.

1. TQQ (10 points) Conjugate momentum to relative position

The momentum \vec{p} conjugate to the relative position \vec{r} is defined with components $p_x = \partial \mathcal{L}/\partial \dot{x}$ and so on. Show that $\vec{p} = \mu \dot{\vec{r}}$. Prove also that in the center of mass frame, \vec{p} is the same as the momentum of particle $\vec{p_1}$ (and also $-\vec{p_2}$).

| 2. TQQ/D (20 points) Circular orb | 2. | TQQ/D (20 |) points) |) Circula | r orbits |
|-----------------------------------|----|-----------|-----------|-----------|----------|
|-----------------------------------|----|-----------|-----------|-----------|----------|

(a) Using elementary Newtonian mechanics, find the period of a mass m on a circular orbit of radius r around a **fixed** mass m_2 (under the influence of Newton's Law of Gravity).

(b) Using the separation into CM and relative motions, find the corresponding period for the case that m_2 is not fixed and the masses circle each other a constant distance r apart. Discuss the limit of this result as $m_2 \to \infty$.

(c) What would be the orbital period if Earth was replaced by a star of mass equal to the solar mass, with in a circular orbit, with the distance between the Sun and the star equal to the present earth-sun distance? (The mass of the sun is more than $300,000 \times$ that of Earth.)


3. TQQ (20 points) Two masses and a spring in a plane

Consider two particles of equal masses $m=m_1=m_2$ attached to each other by a light straight spring (spring constant k, natural length L) and free to slide on top of a frictionless horizontal table.

(a) Write down the Lagrangian in terms of the coordinates $\vec{r_1}$ and $\vec{r_2}$ and rewrite in terms of CM and relative positions, \vec{R} and \vec{r} , using polar coordinates (r, ϕ) for \vec{r} .

(b) Write down and solve the Lagrange equations for the CM coordinates X,Y.

(c) Write down the Lagrangian equations for r and ϕ . Solve these for the two special cases that r remains constant and ϕ remains constant. Describe the corresponding motions. In particular, show that the frequency of oscillations in the 2nd case is $\omega = \sqrt{2k/m}$.



5. TQQ / ACT (40 points) Hand in Activity 8.1