

17-5-11

8-9 Month

A90181

Csmocote® Blend

Recommended Rates

Longevity at average media temperature

60°F (15°C)	70°F (21°C)	80°F (26°C)	90°F (32°C)
9-10 months	8-9 months	6-7 months	5-6 months

Surface application (grams)

Common container volumes	Approx. containers per cubic yard ¹	Low	Med	High
1 quart	850	4	6	9
2 quart	400	9	14	18
1 gallon trade	300	11	18	24
1 gallon	210	16	26	35
2 gallon trade	125	27	44	58
2 gallon	102	33	53	71
3 gallon	70	49	78	104
5 gallon	52	65	105	140
7 gallon	35	97	156	208

Large container volumes	Surface area in square feet	Low	Med	High
10 gallon (17" diam.)	1.4	118	188	251
15 gallon (17.5" diam.)	1.5	126	202	269
20 gallon (21" diam.)	2.3	193	309	413
25 gallon (22.5" diam.)	2.8	235	377	502
30 gallon (26.5" diam.)	3.8	319	511	682
45 gallon (30" diam.)	4.8	404	646	861
65 gallon (30" diam.)	4.8	404	646	861
100 gallon (36" diam.)	7.1	597	955	1273
200 gallon (48.5" diam.)	12.8	1076	1722	2296
24" box	4.0	336	538	717
30" box	6.25	525	841	1121
36" box	9.0	757	1211	1614
48" box	16.0	1345	2152	2870
For containers not listed, multiply surface area by:		84	135	179

Incorporation

	Low	Med	High
Pounds per cubic yard	7.5	12.0	16.0
Kilograms per cubic meter	4.4	7.1	9.5
Grams per liter	4.4	7.1	9.5

Landscape²

	Low	Med	High
Pounds per 1000 square feet	12.0	20.5	29.5
Kilograms per 100 square meters	5.9	10.0	14.4
Pounds of N per 1000 square feet	2.0	3.5	5.0

¹May vary depending on container brand, media, and fill method.²ICL recommends low rate on heavy or clay soils, high rate on light or sandy soils.

Approximate Volume Measurements

ICL spoons

	#1	#2	#3	#4	#5	#6	#7
Grams	9	13	17	36	47	69	94

Conventional measures

	1 tsp.	1 tbsp.	1/4 c.	1/3 c.	1/2 c.	1 c.
Grams	5	15	66	88	132	263

Guaranteed Analysis

Total Nitrogen (N) ³	17%
9% Ammoniacal Nitrogen (N-NH ₄)	
8% Nitrate Nitrogen (N-NO ₃)	
Available Phosphate (P ₂ O ₅) ³	5%
Soluble Potash (K ₂ O) ³	11%
Calcium (Ca) ³	2.2%
Sulfur (S) ³	5.7%
5.7% Combined Sulfur (S)	
Copper (Cu).....	0.130%
0.130% Water Soluble Copper (Cu)	
Iron (Fe) ³	2.21%
2.21% Water Soluble Iron (Fe)	
Manganese (Mn) ³	0.32%
0.32% Water Soluble Manganese (Mn)	
Molybdenum (Mo) ³	0.0060%
Zinc (Zn).....	0.130%
0.130% Water Soluble Zinc (Zn)	

Derived from: Polymer-coated: Ammonium Nitrate, Ammonium Phosphate, Calcium Phosphate, Potassium Sulfate; Copper Sulfate, Ferrous Sulfate, Manganese Sulfate, Sodium Molybdate, Zinc Sulfate

³The nitrogen, phosphate, and potash sources have been coated to provide 17% coated slow-release nitrogen (N), 5% coated slow-release available phosphate (P₂O₅), 11% coated slow-release soluble potash (K₂O), 1.33% coated slow-release calcium (Ca), and 4.14% coated slow-release sulfur (S).

Directions

- recommended for use in covered greenhouses (low to medium rates suggested), nurseries, landscape beds, and containers
- verify product analysis, longevity, and rates (for assistance, contact your regional ICL Territory Manager or call ICL Customer Service at 800-492-8255)
- thoroughly blend into growing media to ensure uniform distribution without over-mixing
- growing media should be used two to four weeks after incorporation
- for top-dress applications, spread fertilizer evenly on container surface (avoid piling fertilizer directly against plant stem)
- irrigate after application (irrigation frequency and volume should be monitored and adjusted during the crop production cycle)
- a product trial is recommended before adopting a new fertilizer program or making full-scale changes to standard local practices
- use caution when applying to plants being over-wintered under cover (if you can't monitor soluble salts and/or adjust irrigation, avoid Fall and Winter applications)
- store in a clean, cool, dry place

All information is intended for use as a guideline only and may not be suitable for all regions and conditions.