## A correction to the photoelectric current in the Planck's constant experiment

Evan S. Snyder

Citation: The Physics Teacher 23, 98 (1985); doi: 10.1119/1.2341733

View online: https://doi.org/10.1119/1.2341733

View Table of Contents: http://aapt.scitation.org/toc/pte/23/2 Published by the American Association of Physics Teachers

### Articles you may be interested in

Improved student laboratory on the measurement of Planck's constant using the photoelectric effect American Journal of Physics **56**, 86 (1988); 10.1119/1.15387


## Planck's constant experiment

The Physics Teacher 23, 334 (1985); 10.1119/1.2341835

What was measured in Millikan's study of the photoelectric effect? American Journal of Physics **83**, 765 (2015); 10.1119/1.4928173

Photoelectric Effect and Planck's Constant in th Introductory Laboratory

American Journal of Physics 39, 50 (1971); 10.1119/1.1986055



# A correction to the photoelectric current in the Planck's constant experiment

Evan S. Snyder Ursinus College, Collegeville, Pennsylvania 19426

The experiment which uses the photoelectric effect to determine Planck's constant is important both for its historical value and its conceptual value. One of the problems with this experiment is the photoemission from the collector which causes a negative current through the cell. This makes the determination of the cutoff potential difficult.

To make our correction we shall assume that the relation between the current from the collector and the voltage across the photocell is roughly the same as the relation between the current from the photocathode and the voltage. There may be reason to question the validity of this assumption because of the different geometrical configuration of the two electrodes. We shall see later that there is an empirical check that seems to justify this assumption.

In Fig. 1 the curve labeled I (+V) represents the true current from the photocathode. It is this curve that we wish to extract from the observed curve I' (+V). The curve labeled fI (-V) represents the current from the collector. Note that fI (-V) is negative with respect to I (+V) both in direction and in its dependence on V. This means that, for some negative voltage, -V, we expect the current from the collector to be some negative fraction, f, of the current from the photocathode for the corresponding positive




Fig. 1. The observed photoelectric current is the sum of the electron emission from the cathode and from the collector.

voltage +V. The value of f should lie somewhere between -1 and 0. The measured current, I' (+V), is the sum of the two components:

$$I'(+V) = I(+V) + fI(-V)$$
If  $I'(+V)$  is the massured current for some voltage

If I' (+V) is the measured current for some voltage




Fig. 2. A plot of |I'(+V)/I'(-V)| versus |V|. The value of this ratio of |V|greater than the cutoff potential should give f.




Fig. 3. Experimental data and corrected data using the value of f from Fig. 2.

+V, then I' (-V) should give the measured current for the voltage -V. That is

$$I'(-V) = I(-V) + fI(+V)$$

Multiplying this equation by f and subtracting from the equation for I' (+V) yields

$$I(+V) = \frac{I'(+V) - fI'(-V)}{1 - f^2}$$

To obtain the value of f we note in Fig. 1 that I (+V) goes to zero at the cutoff potential  $-V_c$  and fI (-V) goes to zero at  $+V_c$ . In the range  $-V_c$  to  $+V_c$  the current I' (+V) is the sum of the two components. Below  $-V_c$  the current I' (+V) should be due to fI (-V) only and above  $V_c$  it should be due to I (+V) only. The f may be found by taking the ratio of I' (-V) for a negative voltage less than  $-V_c$  and dividing it by the value of I' (+V) for the corresponding positive voltage above  $V_c$ . That is

$$f = \frac{I'(-V)}{I'(+V)}$$

Figure 2 was obtained by measuring I' (V) for voltages from -2 V to +2 V for  $\lambda$  = 436 nm (Hg arc spectrum). The ratio I' (-V)/I' (+V) was found for each negative voltage and the corresponding positive voltage. The ratio is plotted versus the absolute value of voltage. It may be seen that this ratio is reasonably constant from 2.0 V to 1.5 V. This suggests the assumption made earlier concerning the similarities of the two components of the current is a valid approximation.

The mean value of the ratio over this nearly constant interval yields a value of  $f = -9.08 \times 10^{-3}$ . Figure 3 shows the raw data and the corrected data using this value of f. Other lines in the mercury spectrum may be treated in the same way. The value of f obtained is slightly different for different lines, however.

This treatment of the data does not result in a phenomenal improvement in the value of Planck's constant, but the cutoff potentials may be read from the curves with more confidence and fewer apologies for the reverse current.

#### ASTRONOMY EDUCATION MATERIALS

Do you want to share astronomy instructional materials you have created and at the same time get credit for them? The Astronomy Education Materials Project provides a way for this to happen. A dissemination network has been established with funds from the National Academy of Science (V. M. Slipher Fund) to foster the exchange of useful astronomy instructional units and aids. If you have developed lab and field activities, instructional modules, film,

games, planetarium activities, computer simulations or other materials you wish to share on a nonprofit basis (or would like to receive these types of materials) write for further information and submission survey forms to: Dennis W. Sunal, Astronomy Education Materials Project, Department of Curriculum and Instruction, West Virginia University, Morgantown, West Virginia 26506