

DEFENSIVE PUBLICATION

Technical Disclosure Document

**AI-Enabled Wearable and Software Systems
for
Continuous Mental Health Monitoring,
Behavioral Insight Generation, and
Collaborative Elder Care Coordination**

Publication Date: February 25, 2026

Disclosure ID: GC-DP-2026-001

This document constitutes a public disclosure of inventive concepts, methods, systems, and architectures for the purpose of establishing prior art under applicable patent law.

The authors disclaim any intent to patent the disclosed concepts and dedicate them to the public domain for the benefit of open research and innovation.

Section 1: Purpose and Scope of This Disclosure

1.1 Purpose

This document serves as a defensive publication and public technical disclosure. Its purpose is to establish prior art for a broad range of inventive concepts, systems, methods, and architectures related to: AI-enabled wearable devices for continuous mental health monitoring; software platforms for behavioral insight generation in elder care and memory care settings; collaborative care coordination systems connecting care teams, families, and healthcare providers; and the integration of sensor data, artificial intelligence, and healthcare interoperability standards for the detection, monitoring, and management of cognitive and mental health conditions.

By publicly disclosing these concepts, the authors intend to prevent any party from obtaining patent rights that would restrict open research, innovation, or commercial development in these domains. This disclosure covers both implemented systems and anticipated future developments that are the natural extensions of the described architectures.

1.2 Scope of Conditions Addressed

The systems, methods, and devices disclosed herein address the full spectrum of mental health and cognitive conditions that may affect individuals, with particular emphasis on elder care populations. These include but are not limited to:

- Dementia (all types including Alzheimer's disease, vascular dementia, Lewy body dementia, frontotemporal dementia, and mixed dementia)
- Mild cognitive impairment (MCI) and pre-dementia cognitive changes
- Depression and major depressive disorder
- Anxiety disorders (generalized anxiety, social anxiety, panic disorder, agoraphobia)
- Paranoia and persecutory ideation
- Delirium (acute confusional states)
- Sundowning syndrome and circadian rhythm disruptions
- Agitation, aggression, and behavioral and psychological symptoms of dementia (BPSD)
- Apathy and withdrawal
- Sleep disorders (insomnia, sleep apnea, REM sleep behavior disorder, restless leg syndrome)
- Medication-induced cognitive and behavioral changes

- Post-traumatic stress disorder (PTSD) in elderly populations
- Bipolar disorder and mood cycling
- Psychosis and hallucinations
- Social isolation and loneliness-related decline
- Pain-related behavioral changes and undiagnosed pain conditions
- Substance use disorders and withdrawal in geriatric populations

1.3 Legal Notice

The concepts described herein are disclosed to the public without restriction. The authors do not claim patent rights to any of the described inventions, methods, or systems. Any person or organization is free to implement, modify, extend, or commercialize any concept described in this document. This disclosure is intended to serve as prior art under 35 U.S.C. § 102 (United States), Article 54 of the European Patent Convention, and equivalent provisions under patent law in all jurisdictions worldwide.

Section 2: Wearable Device Systems for Mental Health Monitoring

2.1 Sensor Modalities

The disclosed system encompasses wearable devices that incorporate one or more of the following sensor modalities for the purpose of detecting, monitoring, or predicting mental health conditions. Each sensor modality may be used independently or in combination with others (sensor fusion) to improve detection accuracy.

2.1.1 Motion and Activity Sensors

1. Tri-axial accelerometers for detecting gait patterns, tremors, falls, wandering behavior, repetitive movements, agitation-related motor patterns, and activity levels.
2. Gyroscopes for measuring rotational movement, postural sway, balance deterioration, and movement quality degradation indicative of cognitive decline.
3. Magnetometers for compass-based orientation tracking, spatial navigation assessment, and wandering pattern characterization.
4. Barometric pressure sensors for altitude-based activity tracking (stairs, elevation changes) and detection of positional changes (lying, sitting, standing).
5. Pedometers and step counters for daily activity quantification, circadian activity rhythm analysis, and detection of activity pattern changes associated with depression or cognitive decline.

2.1.2 Physiological Sensors

1. Photoplethysmography (PPG) sensors for continuous heart rate monitoring, heart rate variability (HRV) analysis, stress detection, autonomic nervous system assessment, and anxiety episode identification.
2. Electrodermal activity (EDA) / galvanic skin response (GSR) sensors for measuring sympathetic nervous system arousal, emotional state changes, stress responses, and anxiety indicators.
3. Skin temperature sensors for circadian rhythm assessment, fever detection, stress-related temperature changes, and thermoregulation monitoring.
4. Blood oxygen saturation (SpO2) sensors for detecting hypoxia-related confusion, sleep apnea monitoring, and respiratory status assessment.
5. Electroencephalography (EEG) sensors (dry electrode or flexible electrode arrays) for brainwave monitoring, sleep stage classification, seizure detection, cognitive load assessment, and attention tracking.

6. Electromyography (EMG) sensors for muscle tension monitoring, stress-related tension patterns, involuntary movement detection, and restlessness quantification.
7. Blood pressure estimation sensors using pulse transit time (PTT), pulse wave analysis, or oscillometric methods for cardiovascular monitoring related to cognitive health.
8. Cortisol and biomarker sensors using sweat analysis, interstitial fluid sampling, or transdermal measurement for stress hormone monitoring and neuroendocrine assessment.

2.1.3 Environmental and Contextual Sensors

1. Ambient light sensors for light exposure tracking, circadian rhythm assessment, and environmental context determination (indoor/outdoor, time of day validation).
2. Microphones for voice analysis, speech pattern monitoring, vocal biomarker extraction, conversational engagement assessment, and environmental sound classification.
3. GPS and location services for geofencing, wandering detection, route deviation analysis, spatial behavior pattern tracking, and place-based behavioral correlation.
4. Bluetooth Low Energy (BLE) beacons and proximity sensors for indoor positioning, room-level location tracking, social proximity detection, and interaction frequency measurement.
5. Near-field communication (NFC) sensors for identifying interaction with objects, medication compliance tracking, and activity verification.
6. Ultra-wideband (UWB) sensors for precise indoor positioning, gesture recognition, and fine-grained spatial behavior analysis.

2.1.4 Optical and Visual Sensors

1. Infrared (IR) sensors for sleep monitoring without visible light, nighttime activity detection, and non-invasive vital sign estimation.
2. Time-of-flight (ToF) depth sensors for gesture recognition, fall detection, and movement pattern analysis.
3. Camera modules (privacy-preserving, edge-processed only) for facial expression analysis, pupil dilation tracking, and non-verbal communication assessment.

2.2 Wearable Form Factors

The disclosed sensor systems may be implemented in any of the following form factors, each optimized for different use cases, comfort levels, and monitoring capabilities:

- Wrist-worn devices (smartwatches, fitness bands, medical-grade wrist monitors)

- Ring-form devices for unobtrusive continuous monitoring
- Pendant or necklace devices for fall detection and location tracking
- Clip-on or pin-on devices for clothing-attached monitoring
- Headband or ear-worn devices for EEG, brain activity, and audio monitoring
- Smart insole or shoe-based devices for gait analysis and fall prediction
- Patch-type adhesive sensors for extended physiological monitoring
- Smart textile and fabric-embedded sensors woven into everyday clothing
- Belt-mounted or waistband devices for core body motion tracking
- Eyeglass-mounted devices for visual field analysis and head movement tracking
- Bed-integrated sensor mats for sleep monitoring without wearable compliance
- Chair-integrated pressure and motion sensors for seated behavior analysis

2.3 On-Device AI Processing

The disclosed system includes methods for performing artificial intelligence and machine learning inference directly on the wearable device, reducing latency, preserving privacy, and enabling operation without continuous network connectivity. On-device processing methods include:

1. Quantized neural networks (INT8, INT4) optimized for microcontroller-class processors for real-time classification of sensor data into behavioral states.
2. TinyML models deployed on ARM Cortex-M class processors or RISC-V processors for edge inference of mood states, activity classification, and anomaly detection.
3. Federated learning approaches where model updates are computed locally on the device and only aggregated model parameters (never raw data) are transmitted to improve global models.
4. Adaptive threshold algorithms that learn individual baseline patterns and trigger alerts based on personalized deviation thresholds rather than population-level norms.
5. Hierarchical inference pipelines where low-power always-on classifiers trigger more computationally expensive secondary models only when initial screening detects potential events of interest.
6. Temporal convolutional networks and transformer-based architectures adapted for time-series sensor data classification at the edge.
7. Anomaly detection using autoencoders, isolation forests, or one-class SVMs trained on individual baseline behavioral patterns to flag deviations without requiring labeled training data for every condition.

Section 3: AI Methods for Mental Health Condition Detection

3.1 Cognitive Decline and Dementia Detection

The following methods and signal combinations are disclosed for the detection and monitoring of cognitive decline, dementia progression, and related cognitive impairments:

3.1.1 Gait and Movement Analysis

Machine learning models that analyze accelerometer and gyroscope data to extract gait parameters including stride length, cadence, gait variability, dual-task gait cost, turning behavior, and gait symmetry. Changes in these parameters over time serve as biomarkers for cognitive decline. Specific methods include: convolutional neural networks applied to raw accelerometer waveforms for gait phase segmentation; recurrent neural networks (LSTM, GRU) for modeling temporal gait pattern evolution; statistical process control methods for detecting significant gait parameter drift over weeks and months; and correlation analysis between gait degradation and scores on standardized cognitive assessments.

3.1.2 Speech and Language Analysis

Natural language processing and audio signal processing methods applied to voice recordings from wearable microphones to detect cognitive decline indicators. These include: analysis of speech rate, pause patterns, and hesitation frequency; vocabulary richness metrics (type-token ratio, lexical diversity) tracked over time; semantic coherence analysis using word embedding similarity between consecutive utterances; automated speech recognition with error analysis where increased ASR errors correlate with speech degradation; prosody analysis including pitch variation, volume modulation, and emotional expression range; detection of word-finding difficulties, circumlocution, and semantic paraphasia; and naming accuracy assessment through structured conversational prompts.

3.1.3 Circadian Rhythm and Sleep Analysis

Methods for detecting circadian rhythm disruptions that are early indicators of cognitive decline and dementia. These include: actigraphy-based rest-activity rhythm analysis using accelerometer data; sleep architecture estimation from movement and heart rate data; sundowning detection through late-afternoon and evening agitation pattern recognition; sleep fragmentation quantification and correlation with daytime cognitive function; REM sleep behavior disorder detection as a prodromal marker for Lewy body dementia; and longitudinal tracking of sleep timing drift, sleep efficiency changes, and nocturnal wandering patterns.

3.1.4 Spatial Navigation and Wandering

GPS, indoor positioning, and inertial navigation methods for assessing spatial cognition. Disclosed approaches include: route complexity analysis comparing traveled paths against efficient routes; familiar route deviation detection using personalized geospatial models; wandering classification algorithms that distinguish purposeful walking from confused wandering based on path geometry, speed variation, and temporal context; progressive geographic range monitoring that detects shrinking or expanding movement territories over time; and integration of spatial behavior analysis with time-of-day context to identify sundowning-related wandering.

3.2 Depression Detection and Monitoring

Methods for continuous, passive monitoring of depressive symptoms using wearable sensor data:

1. Activity level analysis: Tracking daily step counts, movement intensity, and active minutes with statistical models that detect progressive activity reduction patterns characteristic of depression onset or deepening.
2. Social interaction quantification: Using proximity sensors, Bluetooth scanning, microphone-based voice activity detection, and location patterns to estimate social engagement frequency, duration, and quality.
3. Sleep disruption patterns: Identifying depression-associated sleep patterns including insomnia (increased sleep onset latency, nocturnal awakenings), hypersomnia (excessive total sleep time), and early morning awakening.
4. Heart rate variability analysis: Reduced HRV as a physiological biomarker for depressive states, with longitudinal tracking to detect trends rather than relying on single measurements.
5. Diurnal variation modeling: Tracking daily mood-related physiological patterns where morning worsening and evening improvement (or vice versa) patterns are characteristic of specific depressive subtypes.
6. Psychomotor retardation detection: Using fine-grained accelerometer analysis to detect slowed movements, reduced gesture frequency, and decreased movement complexity associated with depression.
7. Voice biomarker tracking: Monitoring changes in speech rate, pitch, energy, spectral characteristics, and phonation quality that correlate with depressive severity.

3.3 Anxiety Detection and Monitoring

Disclosed methods for detecting anxiety states, panic episodes, and chronic anxiety patterns:

1. Physiological arousal detection: Multi-sensor fusion of heart rate elevation, HRV reduction, electrodermal activity increase, and skin temperature changes for real-time anxiety state classification.

2. Panic attack detection: Rapid-onset physiological signature detection combining sudden heart rate acceleration, respiratory rate increase, electrodermal spike, and motor agitation for automated panic episode identification and timestamping.
3. Anticipatory anxiety modeling: Pattern recognition of physiological arousal preceding regular events (meals, activities, transitions) to identify anxiety triggers and enable preemptive intervention.
4. Restlessness and fidgeting quantification: Accelerometer-based detection of repetitive non-purposeful movements, postural shifts, and hand/limb movements indicative of anxious states.
5. Hypervigilance detection: Analysis of startle responses (sudden movement amplitudes), environmental scanning behavior (frequent head/body orientation changes), and sleep-onset latency patterns.

3.4 Paranoia and Psychosis Indicators

Methods for detecting behavioral indicators associated with paranoid ideation and psychotic episodes:

1. Social avoidance pattern detection: Tracking progressive reduction in proximity to others, avoidance of common areas, and changes in social interaction patterns over time.
2. Agitation escalation modeling: Sequential pattern recognition across physiological and movement data to identify escalating agitation that may precede paranoid episodes.
3. Sleep disruption correlation: Analyzing the relationship between sleep deprivation patterns and subsequent behavioral disturbance, with predictive models for psychosis-risk periods.
4. Speech pattern anomaly detection: Identification of changes in speech including increased speech rate, pressured speech, tangential communication, and self-referential content increases.
5. Environmental response analysis: Detecting behavioral responses to environmental stimuli that may indicate hallucinations or misperceptions, such as responding to non-present stimuli, sudden startles without apparent cause, or extended attention directed at empty spaces.

3.5 Multi-Condition Differential Analysis

Methods for distinguishing between conditions that share overlapping symptoms, which is a critical challenge in mental health monitoring:

- Delirium vs. dementia differentiation using onset speed analysis, fluctuation patterns, attention metrics, and temporal trajectory modeling
- Depression vs. apathy in dementia using motivation-related behavioral probes, response to social stimuli, and hedonic capacity assessment

- Anxiety vs. pain-related agitation using movement quality analysis, positional preferences, and physiological pattern differentiation
- Medication side effects vs. disease progression using temporal correlation with medication administration, dose changes, and pharmacokinetic modeling
- Sundowning vs. general agitation using time-of-day analysis, light exposure correlation, and preceding sleep quality assessment
- UTI-related behavioral changes vs. psychiatric symptoms using temperature trends, activity pattern acute changes, and historical episode pattern matching

Section 4: Sensor Fusion and Multi-Modal Analytics

4.1 Fusion Architectures

The disclosed system employs multiple approaches to combining data from heterogeneous sensor sources for improved detection accuracy and reduced false positive rates:

1. Early fusion: Raw sensor data from multiple modalities is concatenated or interleaved into a unified feature vector before being processed by a single model. This approach captures cross-modal correlations at the data level.
2. Late fusion: Each sensor modality is processed by a dedicated model that produces an independent prediction or confidence score. These scores are then combined using weighted voting, Bayesian inference, or a meta-learning ensemble to produce a final output.
3. Attention-based fusion: Transformer-style attention mechanisms that learn to dynamically weight the importance of different sensor modalities based on context, time of day, activity state, and the specific condition being assessed.
4. Hierarchical fusion: A multi-level architecture where low-level sensor features are first fused within related groups (e.g., all motion sensors, all physiological sensors), and group-level representations are then combined at a higher level for final inference.
5. Graph neural network fusion: Representing sensor data as a graph where nodes represent sensor readings and edges represent spatial, temporal, and causal relationships, enabling the model to learn complex inter-sensor dependencies.
6. Missing data resilient fusion: Architectures that gracefully degrade when one or more sensor modalities are temporarily unavailable (sensor removal, charging, malfunction) by using imputation, marginal inference, or modality dropout training.

4.2 Temporal Pattern Mining

Methods for extracting clinically meaningful patterns from longitudinal multi-sensor time-series data:

- Change point detection algorithms that identify statistically significant shifts in behavioral baselines over periods of days, weeks, or months
- Seasonal and cyclical pattern extraction accounting for day-of-week effects, monthly patterns, seasonal affective patterns, and holiday/event-related behavioral changes
- Episode pattern recognition that identifies recurring sequences of physiological and behavioral states that precede adverse events

- Progressive trend analysis using regression, spline fitting, and state-space models to characterize gradual cognitive or functional decline trajectories
- Causal discovery methods that attempt to identify directional relationships between environmental factors, interventions, physiological states, and behavioral outcomes
- Transfer learning approaches where patterns learned from one individual's data are adapted to improve detection for another individual with limited historical data

4.3 Personalized Baseline Learning

The system discloses methods for establishing and continuously updating individual behavioral and physiological baselines, recognizing that population-level norms are insufficient for accurate detection in elderly populations with heterogeneous health profiles:

- Initial enrollment period algorithms that establish personal baselines from the first 7-30 days of monitoring, with uncertainty quantification that improves as more data is collected
- Adaptive baseline methods that account for expected progressive changes (gradual cognitive decline) while still detecting acute deviations (delirium, UTI-related changes)
- Context-aware baselines that maintain separate baseline profiles for different contexts (morning routine, mealtime, social activities, nighttime) rather than a single global baseline
- Caregiver-validated baseline calibration where care team observations are used to label periods as “typical” or “atypical” to improve baseline model accuracy

Section 5: Behavioral Insight and Care Coordination Platform

5.1 System Architecture

The disclosed software platform provides an integrated environment for collecting, analyzing, and acting on behavioral and health data from multiple sources. The architecture encompasses:

1. A data ingestion layer that receives and normalizes data from wearable devices, environmental sensors, electronic health records, care team observations, and family reports into a unified data model based on HL7 FHIR (Fast Healthcare Interoperability Resources) standards.
2. An analytics engine that applies machine learning models, statistical analysis, and rule-based systems to detect patterns, generate insights, and produce actionable recommendations.
3. A care coordination interface that presents insights to care teams in context-appropriate formats, enabling structured observations, shift handoffs, intervention tracking, and longitudinal outcome analysis.
4. A family engagement portal that translates clinical observations and trend data into accessible, meaningful summaries for family members, supporting transparency and collaborative care planning.
5. An administrative dashboard that provides facility-level analytics, quality metrics, staff performance insights, regulatory compliance documentation, and resource optimization recommendations.
6. An interoperability layer based on FHIR R4 standards and Model Context Protocol (MCP) server architecture that enables data exchange with external healthcare systems, research institutions, and AI analysis tools.

5.2 Structured Behavioral Observation System

The platform includes a guided observation system for care workers that captures both structured data and narrative context:

- Multi-component assessments where a single observation captures mood, agitation level, confusion episodes, communication quality, social engagement, eating independence, activity participation, and sleep quality as related but independently valued dimensions
- Coded value systems using industry standard terminologies (SNOMED CT, LOINC) augmented by custom code systems for elder-care-specific concepts (agitation scales, engagement scores, intervention response ratings)
- Reference range definitions that enable automated flagging of abnormal values, with facility-customizable and individual-customizable threshold definitions

- Free-text narrative capture with natural language processing to extract structured concepts, sentiment, and intervention mentions from care worker notes
- Voice-to-observation conversion where care workers can speak observations that are automatically transcribed, parsed into structured components, and presented for review and confirmation
- Photo and short video attachment capabilities for documenting observable conditions (skin changes, mobility patterns, environmental factors) with privacy controls and automated redaction options

5.3 AI-Powered Insight Generation

The platform applies artificial intelligence to generate actionable insights from collected data:

1. Pattern recognition that identifies what time of day residents experience most confusion, agitation, or positive engagement, and correlates these patterns with environmental and care factors.
2. Intervention effectiveness analysis that tracks which approaches (music therapy, redirection, physical activity, social interaction, environmental modification) produce the best outcomes for specific residents in specific contexts.
3. Predictive alerting that identifies early warning patterns (changes in sleep, activity, social behavior) that historically precede adverse events for a specific individual, enabling proactive intervention.
4. Care plan optimization that synthesizes longitudinal observation data, intervention outcomes, and resident preference information to generate evidence-based care plan recommendations.
5. Shift handoff intelligence that summarizes the most relevant recent observations, ongoing concerns, and recommended approaches for incoming care staff, personalized to each resident.
6. Family communication generation that produces meaningful, compassionate summaries of a resident's week including positive moments, areas of concern, and care team responses, written in accessible language.
7. Staff pattern analysis that identifies which care workers achieve the best outcomes with specific residents or in specific situations, enabling optimized scheduling and mentorship.
8. Population-level trend analysis that identifies facility-wide patterns such as seasonal behavioral changes, environmental factors affecting multiple residents, or systemic care process improvements.

5.4 Large Language Model Integration

The disclosed system integrates large language models (LLMs) through a Model Context Protocol (MCP) server architecture to provide conversational AI analysis capabilities:

- FHIR-to-LLM translation layer that converts structured healthcare data into contextually rich prompts, and translates LLM outputs back into structured data formats
- Conversational query interface where care workers can ask natural language questions about resident patterns, intervention histories, and care recommendations
- Automated report generation where LLMs synthesize structured observation data, trend analysis, and intervention outcomes into narrative clinical reports and family-friendly summaries
- Clinical decision support where LLMs analyze current resident status against historical patterns, published care guidelines, and facility best practices to suggest appropriate interventions
- Multi-modal context assembly where the LLM receives structured data, narrative notes, sensor trends, and contextual information to provide holistic analysis that no single data source could support independently

Section 6: Healthcare Data Architecture and Interoperability

6.1 FHIR-Based Data Model

The system employs HL7 FHIR R4 as its healthcare data standard, mapping all observations, assessments, and care activities to FHIR resources. The disclosed FHIR implementation includes:

- Observation resources with multi-component structure for behavioral assessments, where each assessment captures multiple related dimensions (mood, cognition, behavior, function) as components of a single observation
- Custom FHIR profiles defining elder-care-specific observation types including daily behavioral assessments, incident reports, functional status evaluations, and family communication summaries
- Custom CodeSystems and ValueSets for elder-care-specific concepts not adequately covered by existing terminologies, including nuanced agitation scales, engagement metrics, and intervention response categories
- Patient, Practitioner, Encounter, and CarePlan resources linked to Observations to maintain full clinical context and enable relationship-aware queries
- Provenance and AuditEvent resources for complete data lineage, supporting regulatory compliance and research data integrity requirements

6.2 MCP Server Architecture

A Model Context Protocol server that exposes care data as FHIR resources to AI systems, enabling intelligent analysis while maintaining data governance:

- Read operations: retrieving individual observations, searching by patient/date/type/performer, generating patient timelines, and fetching recent assessments
- Write operations: creating new observations from care worker input, amending existing observations, appending contextual notes, and recording intervention outcomes
- Analytics operations: behavioral trend analysis, intervention comparison, family report generation, concerning pattern flagging, and population health queries
- Privacy-preserving operations: de-identification pipelines, consent management, role-based access control, and audit trail generation for all data access

6.3 Open Protocol Design

The system is designed around open protocols and standards to prevent vendor lock-in and enable collaborative innovation:

- All data is stored in FHIR-compliant format, enabling export to any FHIR-compatible system
- The MCP server interface is documented publicly, enabling third-party AI tools and analytics platforms to connect
- De-identified data export capabilities support academic research partnerships using standard FHIR queries
- The system supports bidirectional data exchange with electronic health record (EHR) systems through FHIR REST APIs
- Custom extensions are documented and published as FHIR implementation guides, enabling other organizations to implement compatible systems

Section 7: Intervention Recommendation and Response Systems

7.1 Personalized Intervention Engine

The disclosed system includes methods for recommending personalized interventions based on individual history, current state, and evidence from the broader population:

1. Individual intervention history tracking that records what approaches have been tried for each resident, in what context, by which care worker, and with what outcome.
2. Contextual recommendation algorithms that consider current time of day, recent events, physiological state, caregiver identity, environmental factors, and historical response patterns when suggesting interventions.
3. Reinforcement learning models that continuously update intervention recommendations based on observed outcomes, balancing exploitation of known-effective approaches with exploration of potentially beneficial new interventions.
4. Music therapy personalization using documented genre preferences, era associations, emotional responses, and physiological responses to specific music to curate personalized therapeutic playlists.
5. Environmental modification recommendations including lighting adjustments, noise reduction, temperature changes, and spatial reconfiguration based on individual sensory profiles and current behavioral state.
6. Activity recommendation engines that match resident preferences, functional capabilities, social preferences, and current cognitive state to available structured and unstructured activities.
7. De-escalation protocol selection that recommends specific approaches (redirection, validation therapy, physical comfort measures, environmental changes) based on the specific presentation and individual history.

7.2 Alert and Escalation Systems

Multi-tiered alert systems that balance sensitivity with alert fatigue management:

- Tiered alert severity based on statistical deviation magnitude, historical context, and potential risk, ranging from informational notes to urgent notifications
- Smart alert routing that directs notifications to the most appropriate recipient based on the alert type, resident assignment, staff proximity, and care worker expertise
- Alert suppression and grouping logic that prevents alert flooding by combining related signals and suppressing expected patterns (e.g., known medication effect windows)

- Escalation chains that automatically escalate unacknowledged or unresolved alerts to supervisory staff, nursing leadership, or external providers based on configurable time and severity thresholds
- False positive learning where care worker feedback on alert accuracy is used to refine detection models and thresholds over time

7.3 Family Engagement Systems

Methods for keeping families informed and engaged in care without overwhelming them or sharing inappropriate clinical detail:

- AI-generated weekly summaries that highlight positive moments, notable activities, social engagement, and areas where the care team is focusing attention
- Real-time milestone and positive event sharing (participated in a group activity, recognized a family member, enjoyed a favorite meal) with privacy-appropriate context
- Concern communication that frames clinical observations in accessible, empathetic language with clear explanations of care team responses
- Family input channels where relatives can share biographical information, preference updates, historical context, and observations from visits that inform care planning
- Video and photo sharing with consent management, privacy controls, and contextual information about when and why content was captured

Section 8: Privacy, Security, and Ethical Framework

8.1 Privacy-Preserving Architecture

The disclosed system incorporates privacy-by-design principles throughout its architecture:

- On-device processing that performs initial analysis of sensitive sensor data (voice, location, physiological signals) locally before transmitting only derived insights and aggregated metrics
- Differential privacy techniques applied to aggregated analytics to prevent re-identification of individuals from population-level reports
- Federated learning architectures where individual models are improved using local data and only model weight updates are shared, never raw personal data
- Data minimization principles where the system collects, transmits, and stores only the minimum data necessary for each specific function
- Automated data retention policies with configurable retention periods and secure deletion procedures for different data categories
- Homomorphic encryption methods for performing analytics on encrypted data where supported by the computational complexity of the analysis

8.2 Consent and Autonomy

The system addresses the complex consent landscape in elder care and dementia monitoring:

- Tiered consent models that allow granular control over data collection, sharing, and analysis activities
- Surrogate decision-making support for individuals who lack capacity to provide informed consent, with documentation and review workflows
- Dynamic consent interfaces that allow consent decisions to be reviewed and modified as conditions change
- Assent monitoring that detects behavioral indicators of distress or resistance related to monitoring devices and alerts care teams to reconsider monitoring approaches
- Ethical review integration with institutional review board (IRB) workflows for research use of collected data

8.3 Bias and Fairness

Methods for detecting and mitigating algorithmic bias in mental health monitoring systems:

- Demographic parity testing across age, gender, ethnicity, and cognitive impairment level for all detection algorithms
- Training data auditing to identify and address representation gaps in model training datasets
- Explainable AI methods that enable care teams to understand why a particular alert or recommendation was generated, enabling informed clinical judgment
- Regular bias audits comparing system performance and alert rates across different demographic groups within and across facilities
- Human-in-the-loop design where AI provides recommendations and insights but clinical decisions remain with qualified care providers

Section 9: Research Enablement and Open Science

9.1 Research Data Infrastructure

The disclosed system is designed to facilitate research collaboration while maintaining privacy protections:

- Automated de-identification pipelines that remove or generalize personal identifiers from FHIR resources while maintaining analytical utility
- Research cohort definition tools that enable researchers to define study populations based on clinical criteria without accessing individual records
- Standardized data export in FHIR format that enables researchers to use existing FHIR tooling and query languages without custom integration
- Longitudinal data repositories that maintain temporal integrity of observations for time-series research
- Multi-site data federation that enables cross-facility research without centralizing sensitive data

9.2 Open Research Domains

The following research areas are enabled by the disclosed system architecture and are specifically disclosed to ensure they remain available for open investigation:

7. Digital biomarker discovery: Identifying novel sensor-derived markers that correlate with cognitive decline progression, treatment response, or quality of life in elder care populations.
8. Intervention optimization: Using collected outcome data to refine evidence-based intervention strategies for specific behavioral and psychological symptoms of dementia.
9. Predictive modeling: Developing and validating models that predict adverse events (falls, delirium episodes, emergency hospitalizations) from longitudinal sensor and observation data.
10. Caregiver impact research: Analyzing how AI-assisted care tools affect care worker stress, burnout, job satisfaction, and care quality.
11. Family engagement outcomes: Studying the impact of data-driven family communication on family satisfaction, trust, care participation, and resident outcomes.
12. Health economics: Evaluating the cost-effectiveness of sensor-based monitoring, AI-assisted care planning, and proactive intervention strategies in memory care settings.

13. Pharmacovigilance: Using continuous monitoring data to detect and characterize medication side effects, drug interactions, and optimal dosing in elderly populations.
14. Environmental design: Using behavioral data to evaluate and optimize physical environment design in memory care facilities.

Section 10: Additional Disclosed Methods and Systems

The following additional methods, systems, and approaches are disclosed to ensure comprehensive prior art coverage for the field of AI-enabled mental health monitoring and elder care technology:

10.1 Communication and Coordination Methods

- Cross-shift continuity algorithms that automatically identify and highlight the most relevant information for incoming care staff based on current resident status and recent events
- Multi-disciplinary care team coordination systems that route specific observations and insights to physicians, therapists, pharmacists, and social workers based on relevance to their role
- Automated regulatory documentation generation that compiles required assessments, incident reports, and care plan updates from routine observation data
- Emergency department transition summaries that package relevant behavioral history, medication information, and recent changes for acute care providers

10.2 Environmental Integration

- Smart environment control systems that automatically adjust lighting, sound, temperature, and visual displays based on individual resident needs, time of day, and detected behavioral states
- Ambient sensor networks (motion sensors, door sensors, pressure mats, sound level monitors) that provide environmental context for wearable device data interpretation
- Robotic assistance integration where behavioral insights inform the behavior of companion robots, assistive robots, or autonomous delivery systems
- Digital signage and wayfinding systems that adapt to individual cognitive capabilities, providing appropriate visual cues and navigation assistance

10.3 Training and Education Systems

- AI-generated care worker training scenarios based on real (de-identified) behavioral patterns and intervention outcomes
- Real-time coaching systems that provide in-the-moment guidance to care workers based on current resident state and evidence-based best practices
- Competency tracking that identifies specific skill development needs based on a care worker's observed intervention outcomes

- Simulation environments that use historical behavioral data to create realistic training scenarios for new care workers

10.4 Financial and Operational Methods

- Staffing optimization algorithms that use predicted behavioral patterns to recommend staffing levels and skill mix for upcoming shifts
- Cost-of-care analysis that links intervention approaches to outcomes and resource utilization for value-based care reporting
- Insurance and payer reporting integration that generates claims-supporting documentation from routine observation data
- Quality measure computation that automatically calculates CMS-required quality measures from collected observation data

10.5 Device and Infrastructure Methods

- Adaptive battery management systems that adjust sensor sampling rates based on current behavioral state (higher sampling during detected events, lower during stable periods)
- Mesh networking among wearable devices for peer-to-peer data relay, range extension, and proximity-based social interaction detection
- Seamless device replacement and upgrade protocols that maintain continuous monitoring across hardware generations without data loss or baseline reset
- Multi-device correlation where data from a resident's wearable is combined with data from ambient sensors, medication dispensing systems, and care team devices for comprehensive context

Section 11: Summary of Disclosed Inventive Concepts

This section enumerates the specific inventive concepts disclosed in this document for clarity in establishing prior art. Each concept is disclosed individually and in all reasonable combinations with other disclosed concepts.

Concept A: A system combining wearable sensor data with AI analysis to detect onset or progression of mental health conditions including dementia, depression, anxiety, paranoia, and psychosis in elder care populations.

Concept B: A method of using multi-modal sensor fusion (motion, physiological, environmental, audio) processed by machine learning models to classify behavioral and cognitive states in real time.

Concept C: An on-device AI processing architecture for mental health monitoring wearables that performs inference at the edge to preserve privacy and reduce latency.

Concept D: A personalized baseline learning system that establishes individual behavioral and physiological norms and detects clinically meaningful deviations from those norms over time.

Concept E: A software platform that combines structured behavioral observations with AI-generated insights to enable collaborative care coordination among care teams, families, and healthcare providers.

Concept F: A FHIR-based healthcare data architecture that maps behavioral observations, sensor data, and care activities to interoperable healthcare standards with MCP server integration for AI analysis.

Concept G: An intervention recommendation engine that uses reinforcement learning and individual outcome history to suggest personalized care approaches in context.

Concept H: A family engagement system that uses AI to generate meaningful, accessible summaries of care observations and resident wellbeing for non-clinical family members.

Concept I: A privacy-preserving architecture for mental health monitoring that employs on-device processing, federated learning, differential privacy, and data minimization principles.

Concept J: A research enablement platform that provides de-identified, FHIR-formatted behavioral and sensor data for academic research in elder care, digital biomarkers, and intervention optimization.

Concept K: A multi-condition differential analysis system that distinguishes between overlapping mental health conditions using temporal pattern analysis, contextual data, and multi-sensor correlation.

Concept L: An adaptive alert and escalation system for elder care that uses AI to manage alert severity, routing, suppression, and escalation based on individual context and care team response patterns.

Concept M: Voice and speech analysis methods applied to wearable microphone data for detecting cognitive decline, depression, anxiety, and psychosis through changes in speech patterns, vocabulary, prosody, and semantic coherence.

Concept N: Gait and movement analysis methods using wearable motion sensors for detecting cognitive decline progression, fall risk, medication effects, and behavioral state changes.

Concept O: Environmental integration systems that combine wearable device data with ambient sensor networks and smart environment controls to optimize care environments for individual residents.

All disclosed concepts are intended to be available for open research and commercial implementation by any party. The combination of any two or more disclosed concepts into an integrated system is also specifically disclosed as prior art.

Section 12: Publication Attestation

This document is published as a defensive disclosure for the purpose of establishing prior art. The contents of this document describe systems, methods, and architectures that the authors have conceived, developed, or reasonably anticipated as extensions of their work in AI-enabled elder care and mental health monitoring technology.

This publication is made in good faith to support open innovation in healthcare technology and to prevent the encumbrance of these concepts through patent claims by any party, including the authors.

Document Identifier: GC-DP-2026-001

Publication Date: February 25, 2026

Version: 1.0

Author Signature and Date

Author Signature and Date

Witness Signature and Date