

Set 2 Homework, Analysis of Algorithms

Jay R Bolton

May 4, 2012

- P 52: 3.1-1, 3.1-2
- P 60: 3.2-1, 3.2-2 and Problems: 3-1, 3-3, 3-4
- P 107 Problems: 4-1, 4-2, 4-4

Chapter 3

3.1-1 Prove: $\max(f(n), g(n)) = \Theta(f(n) + g(n))$

By theorem 3.1, in order for a func to be big-Theta, it should be both big-O and big-Omega.

\mathcal{O} :

$$\max(f(n), g(n)) \leq f(n) + g(n)$$

$$\max(f(n), g(n)) = O(f(n) + g(n))$$

Ω :

$$2 * \max(f(n), g(n)) \geq f(n) + g(n)$$

$$\max(f(n), g(n)) \geq f(n) + g(n) * 1/2$$

$$\max(f(n), g(n)) = \Omega(f(n) + g(n))$$

3.1-2 Prove: $(n + a)^b = \Theta(n^b)$

Similarly to 3.1-1, we need to prove that the RHS is both big-O and big-Omega of the LHS.

\mathcal{O} :

$$\text{Show : } (n + a)^b \leq n^b * c$$

for some constant c

Where : $b > 0$

Cases :

$a \leq 0$:

$$(n - a)^b < n^b$$

$$(n - a)^b = O(n^b)$$

$a > 0$:

$$n + a \leq n * a$$

$$(n + a)^b \leq (n * a)^b = n^b * a^b$$

$$(n + a)^b \leq n^b * a^b$$

$$(n + a)^b = O(n^b)$$

with constant a^b for $a > 0$

$\Omega :$

$a \geq 0 :$

$$(n + a) \geq n$$

$$(n + a)^b \geq n^b$$

$$(n + a)^b = \Omega(n^b)$$

$a < 0 :$

$$(n - a) \geq n \cdot -a$$

$$(n - a)^b \geq (n \cdot -a)^b$$

$$(n - a)^b \geq n^b \cdot -a^b$$

$$(n - a)^b = \mathcal{O}(n^b)$$

with constant a^b for $a < 0$

3.2-1

Show :

If $f(n)$ and $g(n)$ are monotonically increasing, then so are:

$$f(n) + g(n) :$$

$$f(n) \leq f(m)$$

$$g(n) \leq g(m)$$

$$f(n) + g(n) \leq f(m) + g(m)$$

$$f(g(n)) :$$

$$f(n) \leq f(m)$$

$$g(n) \leq g(m)$$

$$f(g(n)) \leq f(g(m))$$

* Let: $g(n) = p$ and $g(m) = q$

* We know that $p \leq q$ because it was stated that $g(n) \leq g(m)$

* We already said $f(n) \leq f(m)$ for all $n \leq m$, and that $p \leq q$

* Thus $f(p) \leq f(q)$, that is $f(g(n)) \leq f(g(m))$

Show :

If $f(n)$ and $g(n)$ are nonnegative, then:

$f(n) \cdot g(n)$ is monotonically increasing

Definitions :

* $f(n) \leq f(m)$ for all $n \leq m$

* $g(n) \leq g(m)$ for all $n \leq m$

* $f(n) > 0$ for all n

* $g(n) > 0$ for all n

Conclusions :

* Since $f(n)$ and $g(n)$ are monotonically increasing and only positive, then they will only be positively increasing.

* $f(n) \cdot g(n) \leq f(m) \cdot g(m)$ for all $n \leq m$

* This holds true because increasing positive integers multiplied will still be increasing.

3.2-2

Prove :

$$a^{\log(b,c)} = c^{\log(b,a)}$$

I assume we can use the equations above this one.

$$\text{Definition : } q = b^y \Leftrightarrow \log(b, q) = y$$

$$a^{\log(b,c)} = c^{\log(b,a)}$$

$$= \log(c, a^{\log(b,c)}) = \log(b, a)$$

$$= \log(b, c) * \log(c, a) = \log(b, a)$$

$$= \log(c, a) = \log(b, a) / \log(b, c)$$

$$= \log(c, a)$$

This used equations on p56 above the equation we proved.

3-1 The following is a lemma that I'll use for this problem:

a. Prove: $k \geq d \rightarrow p(n) = \mathcal{O}(n^k)$

$$\text{Show : } \sum_{i=0}^d a_i n^i \leq c \cdot n^k \text{ for some constant } c$$

$$\text{Let } a_m = \max(a_i)$$

$$\sum_{i=0}^d a_i n^i \leq (a_m d) \cdot n^d \leq (a_m d) \cdot n^k$$

$$\sum_{i=0}^d a_i n^i = \mathcal{O}(n^k) \quad \text{with constant } (a_m \cdot d)$$

b. Prove: $k \leq d \rightarrow p(n) = \Omega(n^k)$

$$\text{Show : } \sum_{i=0}^d a_i n^i \geq c \cdot n^k \text{ with some constant } c$$

$$\sum_{i=0}^d a_i n^i \geq n^d \geq n^k$$

$$\sum_{i=0}^d a_i n^i = \Omega(n^k) \quad \text{with constant } 1$$

c. Prove: $k = d \rightarrow p(n) = \Theta(n^k)$

See proof in (a) and (b); by Theorem 3.1, n^d is also Θ .

Show : $\sum_{i=0}^d a_i n^i \geq c \cdot n^d$ with some constant c

Also : $\sum_{i=0}^d a_i n^i \leq e \cdot n^d$ with some constant e

$$\sum_{i=0}^d a_i n^i \leq (a_m d) \cdot n^d$$

$$\sum_{i=0}^d a_i n^i \geq n^d$$

d. Prove: $k > d \rightarrow p(n) = o(n^k)$

Show : $\sum_{i=0}^d a_i n^i < c \cdot n^k$ with some constant c

$$\sum_{i=0}^d a_i n^i \leq (a_m d) \cdot n^d < (a_m d) \cdot n^k$$

e. Prove: $k < d \rightarrow p(n) = \omega(n^k)$

Show : $\sum_{i=0}^d a_i n^i > c \cdot n^k$ with some constant c

$$\sum_{i=0}^d a_i n^i \geq n^d > n^k$$

3-3 From largest to smallest:

2^{2^n}
 $(n+1)!$
 $n!$
 e^n
 $n \cdot 2^n$
 2^n
 $(3/2)^n$
 $(\lg n)^{\lg n}$
 $(\lg n)!$
 n^3
 n^2
 $n \lg n, \lg(n!)$
 n
 $2^{\sqrt{2 \lg n}}$
 $(\lg n)^2$
 $\lg n$
 $\sqrt{\lg n}$
 $\lg \lg n$
 $2^{\lg \cdot n}$
 $(\lg n)^*$
 $n^{1/\lg n}$

Some more of them may be in equivalence classes...

- 3-4**
- a. False by counterexample: n and n^2
 - b. False by counterexample: n and n^2
 - c. True: $f(n) \leq g(n)$ and $\lg(f(n)) \leq \lg(g(n))$
 - d. True: $f(n) \leq g(n)$ and $2^{f(n)} \leq 2^{g(n)}$
 - e. True: $f(n) \leq f(n)^2$
 - f. True by transpose symmetry.
 - g. False: $n^2 > c * (n/2)^2$
 - h. True:

$$f(n) + o(n) \leq 2 \cdot f(n)$$

$$f(n) + o(n) \geq 1/2 \cdot f(n)$$

Chapter 4

- 4-1**
- a.
 - b.
 - c.
 - d.

e.

f.

g.

4-2 a.

b.

4-4

a.

b.

c.

d.