

The pair singular series of a polynomial V : the dilated pieces below $H^{1/2}$

Jasper Reichardt*

16 September 2026 — working draft, not for circulation

Abstract

[Draft. The proofs of Theorems 3 and 5 are in [research/explore/PAPER-V-thmM.md](#) and [PAPER-V-smallend-draft.md](#) and are under adversarial reading; nothing here is to be circulated before those readings are recorded.] For a monic irreducible quadratic f without fixed prime divisor, part III decomposed the Cesàro off-diagonal $\text{Off}_f^*(H)$ into pieces indexed by the dilation u and proved that the pieces with $u > H^{1/2+\varepsilon}$ contribute a main term $c_{\text{off}}(f)H$. This paper treats dilations below that threshold. We prove that the pieces with $H^{0.47} < u \leq H^{0.53}$ contribute $O(H^{1-\delta})$, so that $\sum_{u > H^{0.47}} w(u) \mathcal{P}_u(H/u) = c_{\text{off}}(f)H + O(H^{1-\delta})$; the input is a bilinear estimate for Kloosterman fractions of Bourgain and Garaev together with a bound for the additive energy of reciprocals, valid for every modulus, which we deduce from an argument of Cilleruelo and Garaev. For negative discriminants we prove a second result at the other end: the pieces with $u \leq H^{1/3-c}$ contribute $O(H^{1-\delta(c)})$, by running the automorphic kernel method of Grimmelt and Merikoski with the leading coefficient u^2 tracked. What remains is a neighbourhood of $u = H^{1/3}$.

1 Introduction

[Working draft: the proofs are in [research/explore/PAPER-V-thmM.md](#) and [PAPER-V-smallend-draft.md](#) and are under adversarial reading.]

The problem. Let $f = t^2 + bt + c \in \mathbb{Z}[t]$ be monic, irreducible, without fixed prime divisor, of discriminant D . Part III [6] wrote the Cesàro off-diagonal of the pair singular series as a sum of *pieces*,

$$\text{Off}_f^*(H) = P(1) \sum_u \lambda(u) \omega(u) \mathcal{P}_u(H/u),$$

one for each squarefree *dilation* u , the piece $\mathcal{P}_u(Y)$ being the same problem at length $Y = H/u$ with the roots of $x^2 \equiv D$ replaced by those of $u^2 x^2 \equiv D$. Each piece is $O(Y)$, and the weights satisfy $\sum_{u \leq H} |w(u)|/u \asymp \log H$: the trivial bound for the whole sum is $H \log H$, which is exactly the size of the statement to be proved. Every piece must therefore be small, and the difficulty is uniformity in u .

Exponent coordinates. Write $u = H^\alpha$. Each method available for the pieces gives a saving $H^{-s(\alpha)}$ over the trivial bound on an interval of α , and degenerates at the ends of that interval. In these coordinates part III proves $s > 0$ for $\alpha > 2/3$ (averaging the dilations along u), and the present author's extension of that theorem, recorded in the revised part III, gives $s > 0$ for $\alpha > 1/2$ by a dispersion argument with Weil's bound. Below $1/2$ both mechanisms fail: the

*Independent researcher. Correspondence: jasperreichardt@gmail.com.

incomplete Kloosterman sums that dispersion produces have modulus $[d_1, d_2] \asymp Y^2$, and Weil's bound is trivial once the range of u is shorter than the square root of that modulus. This paper supplies two further mechanisms, one on each side of $\alpha = 1/3$.

Results. Theorem 3 treats α near $1/2$: the weight w is decomposed as $1 * \chi_D * \gamma$, and the resulting bilinear forms in u are estimated by Weil's bound when one factor is long (Type I) or short (Type II), and, in the remaining balanced case, by the bilinear bound of Bourgain and Garaev [1] for Kloosterman fractions. That bound needs an estimate for the additive energy of reciprocals; the one in [1] is too weak at the sizes that occur here, and Lemma 1 supplies a stronger one, valid for every modulus, by transferring an argument of Cilleruelo and Garaev [2] for points on modular hyperbolas. Theorem 5 treats small α for negative discriminants: piece u is, after removing the content, a Type I sum for the polynomial $u^2 \ell^2 + |D|$ in the sense of Grimmelt and Merikoski [3], and their automorphic kernel theorem [4] applies at level $u^2 e$ once the two kernel quantities are estimated with the dilation tracked. The diagonal is the number of orbits of the family, $\asymp u$; the off-diagonal, which their general bound would give as $\asymp u^2$, is in fact bounded independently of u (Lemma 4), because the distances inside the family are quantised in steps of $1/(4h)$ whatever the dilation.

What remains. Both mechanisms degenerate at $\alpha = 1/3$, from opposite sides and for unrelated reasons: there the balanced Type II atoms sit at $d^{1/4}$, where the energy input is not strong enough, and the kernel method's diagonal $\asymp u$ meets the trivial bound. The final section explains what the residual band costs. Two points are worth recording. A band of width 2δ in α costs $\delta \cdot H(\log H)^{1-c} \log \log H$, not $\delta H \log H$, because part III's Theorem E applies piece by piece inside it; so a failing method never makes the total worse. And if the savings on both sides vanish only at $\alpha = 1/3$, with implied constants that are powers of $\log H$, then the blocks near the point contribute $O(H)$ in total, which would give $\text{Off}_f^*(H) = O(H)$ — Hypothesis (E) of part I [5], a bounded second-order term in the Cesàro asymptotic. The constant A_f needs strictly more: a saving at the point itself.

2 Notation

We follow part III [6]. For a prime p let $\omega(p)$ be the number of roots of f modulo p , $\lambda(p) = p/(p - 2\omega(p))$, $P_p = 1 - \omega(p)^2/(p - \omega(p))^2$, $P(1) = \prod_p P_p$ and $w(u) = P(1)\lambda(u)\omega(u)$; w is supported on squarefree u and $\sum_{u \leq t} |w(u)| \ll t$. A modulus d' is *admissible* if it is squarefree, all its prime factors are split, and $(d', 2Du) = 1$; $R_{d'}^{(u)} = \{x \bmod d' : u^2 x^2 \equiv D\}$, $\rho_k(d'; a) = \sum_{r \in R_{d'}} e(kar/d')$, and $\langle x \rangle_{d'} \in [1, d']$ is the representative of x . With $B_{d'}^{(Y)}(m) = \sum_{h \leq Y, h \equiv m \pmod{d'}} (Y - h) - Y^2/(2d')$, the piece is

$$\mathcal{P}_u(Y) = \sum_{d' > 1 \text{ admissible}} \frac{\lambda(d')}{d'} \sum_{x \in R_{d'}^{(u)}} B_{d'}^{(Y)}(\langle x \rangle_{d'}), \quad Y = H/u.$$

We write $G_{d'}(a, Y) = \sum_{r \in R_{d'}} B_{d'}^{(Y)}(\langle ar \rangle_{d'})$, $\bar{G}_{d'}(Y)$ for its mean over the units a , $G^\circ = G - \bar{G}$, and $\xi_k(d', N) = (1 - \cos(2\pi kN/d'))/\sin^2(\pi k/d')$ for the coefficients of the finite Fourier expansion of $B^{(N)}$ (part III, Lemma finfourier). Throughout, x^* denotes the inverse of x modulo the modulus at hand, $e_m(z) = e(z/m)$, and $\varepsilon > 0$ is arbitrarily small. Parameters: $\delta = 10^{-3}$, $\eta = \delta/10$, $\delta_3 = \delta$, $\varepsilon' = \eta/7$, $\kappa = \delta/2$, $K_1 = H^{2\eta}$.

3 The energy of reciprocals and bilinear Kloosterman fractions

Lemma 1 (energy of reciprocals, every squarefree modulus). *Let $m \geq 2$ be squarefree, $1 \leq N \leq m$, and $J_4(N) = \#\{x \in [1, N]^4 : (x_i, m) = 1, x_1^* + x_2^* \equiv x_3^* + x_4^* \pmod{m}\}$. Then for every $\varepsilon > 0$,*

$$J_4(N) \ll_{\varepsilon} m^{\varepsilon} (N^2 + N^{7/2} m^{-1/2}).$$

[Proof written in PAPER-V-thmM.md (Cilleruelo–Garaev's argument, transferred to composite moduli); two readings recorded in PROOFS-uniform.md §28, §30.]

Corollary 2. *For squarefree m , $(a, m) = 1$, $m^{1/3} \leq N_1, N_2 \leq m$ and coefficients $|\alpha_i| \leq 1$ supported on units, $\sum_{x_1 \leq N_1} \sum_{x_2 \leq N_2} \alpha_1(x_1) \alpha_2(x_2) e_m(ax_1^* x_2^*) \ll_{\varepsilon} m^{\varepsilon} (N_1 N_2)^{15/16}$.*

[From [1], proof of Theorem 3, together with Lemma 1.]

4 The pieces around $u = H^{1/2}$

Theorem 3. *Let f be as above. There is $\delta_M > 0$ (any $\delta_M < 4\eta/7$ with the parameters above, so $\delta_M = 5 \cdot 10^{-5}$ is admissible) such that*

$$\sum_{H^{0.47} < u \leq H^{0.53}} w(u) \mathcal{P}_u(H/u) \ll_f H^{1-\delta_M}.$$

Consequently, with part III's Theorem on the range $u > H^{1/2+\varepsilon}$ applied at $\varepsilon = 0.03$, $\sum_{u > H^{0.47}} w(u) \mathcal{P}_u(H/u) = c_{\text{off}}(f)H + O(H^{1-\delta_M})$.

[Proof in PAPER-V-thmM.md: Lemma K (smooth incomplete Kloosterman sums), Lemma 1, Corollary 2, Lemma S (Mellin separation) and Steps R1–R5 (reduction, Type I, Type II, residual, covering). Two readings; a confirming read of the merged text is running.]

5 The small dilations

Lemma 4 (the off-diagonal at one level does not grow with the dilation). *Let $h \geq 1$, u squarefree with $(u, 2h) = 1$, $a = u^2$, and let F_u be the set of positive definite forms $g = \begin{pmatrix} A & B \\ B & C \end{pmatrix}$ with $AC - B^2 = ah$, $a \mid B$, $a \mid C$. For $Z \geq 1$,*

$$\sum_{g_2 \in \Gamma_0(a) \backslash F_u} \sum_{\substack{g_1 \in F_u, g_1 \neq g_2 \\ u(g_1, g_2) \leq Z}} (1 + u(g_1, g_2))^{-1/2} \ll_{\varepsilon} (hZu)^{\varepsilon} h^{O(1)} (1 + Z^{1/2}),$$

uniformly in u , where $u(\cdot, \cdot)$ is the point-pair invariant.

[Proof in PAPER-V-smallend-draft.md (undilation to $\Gamma_0^0(u)$, lift to level one, a local count of common orthogonal lines, and a sum over the quantised distances); under reading.]

Theorem 5. *Let f be as above with $D < 0$. For every $c > 0$ there is $\delta_S(c) > 0$ with $\sum_{u \leq H^{1/3-c}} w(u) \mathcal{P}_u(H/u) \ll_f H^{1-\delta_S(c)}$.*

[Proof outline S1–S8 in PAPER-V-smallend-draft.md, resting on [4] Theorem 8.1 and [3] Lemma 3.1, with Lemma 4 and the diagonal count; $\delta_S(c) \asymp c$ once the exponent in the kernel theorem's $\delta^{-O(1)}$ is named. Under reading.]

6 What remains

[the band at $u = H^{1/3}$: the two mechanisms degenerate there; the accounting of PROOFS §31–§34; what a positive saving at the point would give (A_f), and what a point gap with polylog losses would give (part I's Hypothesis (E)).]

References

- [1] Jean Bourgain and Moubariz Z. Garaev. Kloosterman sums in residue rings. *Acta Arith.*, 164(1):43–64, 2014.
- [2] Javier Cilleruelo and Moubariz Z. Garaev. Concentration of points on two and three dimensional modular hyperbolas and applications. *Geom. Funct. Anal.*, 21:892–904, 2011.
- [3] Lasse Grimmelt and Jori Merikoski. On the greatest prime factor and uniform equidistribution of quadratic polynomials, 2025. arXiv:2505.00493.
- [4] Lasse Grimmelt and Jori Merikoski. Weighted averages of $\mathrm{SL}_2(\mathbb{R})$ automorphic kernel, Part I: non-oscillatory functions, 2025. arXiv:2505.00489.
- [5] Jasper Reichardt. The pair singular series of a polynomial I, 2026. Preprint.
- [6] Jasper Reichardt. The pair singular series of a polynomial III: the off-diagonal in Cesàro form, 2026. Preprint.