

The pair singular series of a polynomial II: zeros of Dedekind zeta functions in the second moment of prime values

Jasper Reichardt*

13 September 2026

Abstract

How much does the number of primes among the values $f(1), f(2), \dots$ of a polynomial fluctuate? Part I of this work showed that the answer is encoded in an explicit arithmetic function, the pair singular series $S_f(h)$, and that the diagonal part of its second moment has the Dedekind zeta function ζ_K of $K = \mathbb{Q}[t]/(f)$ as generating function. Here we determine that generating function completely. It is an infinite product of Artin L -functions of the Galois group of f , evaluated at $N(s+1)$, $N = 1, 2, 3, \dots$, whose exponents are virtual characters given by a plethystic logarithm; for quadratics the exponents are necklace numbers. The first factor beyond ζ_K is $1/L(2s+2, \text{Sym}^2 V)$, and $L(s, \text{Sym}^2 V)$ is the product of the Dedekind zeta functions of the field of a root and of the field of a pair of roots. The line $\Re s = -1$ is a natural boundary, for every f including the twin-prime case. Consequently the Riesz means of the diagonal satisfy an explicit formula whose oscillating part is indexed by the zeros of these Dedekind zeta functions, with terms $x^{m-1+\rho/2}$, and whose smooth part contains an unexpected term $x^{m-2/3}$ (times a polynomial in $\log x$ whose degree is one less than the multiplicity of the trivial character in the third plethystic exponent: a constant for S_n , but $a \log x + b$ for cyclic cubics); the formula is proved for quadratics under the Riemann hypothesis for ζ and $L(s, \chi_D)$, and for general f under the corresponding hypotheses; unconditionally the error term is $\Omega_{\pm}(x^{m-3/4})$. For $f = t^2 + 1$ every coefficient is computed and the formula is verified against the exact Riesz means to $x = 4 \cdot 10^7$: the data reproduce the predicted sum over the zeros of ζ and $L(s, \chi_{-4})$ with regression coefficient 1.008. The remaining conjecture of part I (the off-diagonal) is reformulated as the analytic continuation of a Dirichlet series of Weyl sums of the roots of f , for quadratics a series of Salié-type sums, and the two estimates its proof requires are stated. Assuming the Hardy–Littlewood pair conjecture these results describe the second-order term of the variance of the number of prime values of f ; the zero-induced fluctuations lie below the scale at which that transfer is valid, and no statement about prime values themselves and the zeros of ζ_K is claimed.

1 Introduction

The question. Let $f \in \mathbb{Z}[t]$ be irreducible, without fixed prime divisor, and let $C(f)$ be its Bateman–Horn constant, the conjectural density factor for primes among $f(1), f(2), f(3), \dots$. Part I of this work [40] studied not the density but the *fluctuation* of the number of primes along the sequence, through the pair singular series

$$S_f(h) = \prod_p \frac{1 - \omega_{f,2}(p, h)/p}{(1 - 1/p)^2}, \quad \omega_{f,2}(p, h) = 2\omega_f(p) - \nu_f(p, h),$$

*Independent researcher. Correspondence: jasperreichardt@gmail.com.

where $\omega_f(p)$ is the number of roots of f modulo p and $\nu_f(p, h)$ the number of roots s with $s + h$ again a root. Under the Hardy–Littlewood conjecture $S_f(h)/(\log f(t) \log f(t + h))$ is the probability that $f(t)$ and $f(t + h)$ are both prime, and the quantity that measures how far the primes along f are from independent trials is

$$\Sigma_f(H) = \sum_{h \leq H} (S_f(h) - C(f)^2). \quad (1)$$

Part I proved an exact identity,

$$\Sigma_f(H) = -C^2 \sum_{d \geq 2} a_f(d) \left\{ \frac{H}{d} \right\} + C^2 \text{Off}_f(H), \quad a_f(d) = W_f(d) \omega_f(d), \quad (2)$$

with explicit multiplicative weights $W_f(d)$ on squarefree d (notation below), where the first sum, the *diagonal*, comes from pairs of equal roots of f modulo d and Off_f , the *off-diagonal*, from pairs of distinct roots; it showed that the diagonal has Dirichlet series

$$D_f(s) = \sum_d a_f(d) d^{-s} = \zeta_K(s + 1) E_f(s), \quad K = \mathbb{Q}[t]/(f), \quad \text{Res}_{s=0} D_f = \frac{1}{C(f)};$$

and it conjectured (Conjecture 1 there)

$$\Sigma_f^*(H) := \sum_{h \leq H} \left(1 - \frac{h}{H}\right) (S_f(h) - C(f)^2) = -\frac{1}{2} C(f) \log H + A_f + o(1), \quad (3)$$

a theorem for linear f (Friedlander–Goldston, Vaughan, Montgomery–Soundararajan) and equivalent, in Cesàro form, to the statement that the off-diagonal contributes no logarithm (“Hypothesis (E)”). This paper determines E_f and draws the consequences; the off-diagonal is reformulated in Section 5 and left for part III.

In plain terms. The primes along f are not independent trials: divisibility of $f(t)$ by a small prime p is periodic in t , so the small primes act deterministically and only the large ones are random. $\Sigma_f(H)$ is the resulting total correlation up to distance H , and its main term $-\frac{1}{2} C(f) \log H$ is the polynomial version of the fact that primes in short intervals are more regular than random. This paper is about the next layer, the oscillations of Σ_f around its main terms. For the twin-prime problem Goldston and Suriajaya showed that these oscillations are a sum of waves, one for each zero of the Riemann zeta function. For a polynomial the same happens with the zeros of the Dedekind zeta functions of the number fields cut out by f ; there is one more smooth term, $x^{m-2/3}$, that the linear case does not have; the generating function has a natural boundary at $\Re s = -1$, which changes what can be proved; and all of this is visible in exact computations for $t^2 + 1$. Nothing here is a statement about primes themselves: that needs the Hardy–Littlewood pair conjecture, and the conjecture (3) remains a conjecture, now reduced to a precise analytic statement.

The linear case as template. For $f(t) = t$, $S_f = \mathfrak{S}$ is the twin-prime singular series, and Goldston and Suriajaya [20] proved that the Riesz means $\sum_{k \leq x} (x - k)^m \mathfrak{S}(k)$, $m \geq 2$, equal $x^{m+1}/(m+1) - \frac{1}{2} x^m (\log x - H_m + \gamma + \log 2\pi) + x^{m-1} \sum_{|\gamma| \leq U} a(\rho) x^{\rho/2} + O(x^{m-1+\varepsilon})$, the zeros of ζ entering through the factor $1/\zeta(2s+2)$ of the generating series, with error $\Omega_{\pm}(x^{m-3/4})$ (see also [19, 46, 17]). Our generating series is $\zeta_K(s+1) E_f(s)$; the question is what E_f is.

Results. Section 2: E_f is an infinite product of Artin L -functions $L(N(s+1), \Psi_N)$, $N \geq 2$, with virtual characters Ψ_N given by an explicit formula (Theorem 4); $\Psi_2 = -\text{Sym}^2 V$ and $L(s, \text{Sym}^2 V)$ is a product of Dedekind zeta functions of “pair fields” (Lemma 1); for quadratics the exponents are necklace numbers (Proposition 6); $\Re s = -1$ is a natural boundary (Theorem 7). Section 3: the explicit formula for the Riesz means of the diagonal, for quadratics under the Riemann hypothesis for ζ and $L(s, \chi_D)$ with a complete proof (Theorem 10) and for general f under the analogous hypotheses (Theorem 14), and its comparison with exact data for $t^2 + 1$. Section 4: unconditionally the error term is $\Omega_{\pm}(x^{m-3/4})$ (Theorem 15). Section 5: Hypothesis (E) as the analytic continuation of a Dirichlet series of Weyl sums of roots, and the two estimates that would prove it for quadratics. Section 6: what carries over to $\mathbb{F}_q[u]$. Section 7: what all of this means for prime values under the Hardy–Littlewood conjecture. Section 8: what should be done next. Appendix A specifies the computations.

What is conditional on what. As in part I: (A) Bateman–Horn for non-linear f , open, used nowhere; (B) the Hardy–Littlewood pair conjecture for f , used only in Section 7; (C) Conjecture (3), equivalently Hypothesis (E), a statement about the explicit function S_f , reformulated in Section 5, not proved here; and now (D) the Riemann hypothesis for ζ , $L(s, \chi_D)$ and, for general f , the Artin L -functions of its Galois group, needed for the explicit formulas of Section 3 because the natural boundary prevents the contour from leaving the critical strip. Everything in Section 2 and Theorem 15 is unconditional. The zero-induced fluctuations have relative size $T^{-1/4}$, below the error tolerated by the transfer (B): unconditionally, the low-lying zeros of ζ_K govern the second-order fluctuations of the Riesz means of S_f ; under GRH the variance of primes is related to the pair correlation of the zeros of ζ_K by the Goldston–Montgomery theorem [18] in the Selberg-class form of Bui, Keating and Smith [10]; we claim no unconditional link between prime values of f and zeros of ζ_K .

Relation to the literature. The only prior appearance of $\zeta_K(s+1)$ in a singular-series average that we know of is Kuperberg, Rodgers and Roditty–Gershon [32], for prime *elements* of a number field; explicit formulas over the zeros of several L -functions occur in the Goldbach setting [5, 21]; variances governed by higher-degree L -functions over $\mathbb{F}_q[t]$ are in [22]. Natural boundaries of Euler products go back to Estermann [16] and Dahlquist [12], with the multivariable case in [4] and the “ghost” formalism of du Sautoy and Grunewald [13]; the factorisation of Euler products with Frobenian coefficients through L -functions of virtual characters, and the criterion for meromorphy in terms of the representation ring (continuation to the whole plane if and only if the local polynomial is cyclotomic in $R(G)$, otherwise a natural boundary), are due to Kurokawa [33] and Moroz [37]; both, together with the “log factorisation” that produces the virtual characters, are set out in Alberts’s survey [2], whose class of Euler products is $\prod_p Q_p(p^{-s})$ with $p \mapsto Q_p$ Frobenian. Our local factor F_p is not of that form: it depends on p^{-1} as well as on p^{-s} and Frob_p , and the two variables do not separate. Relative to that theory, what is new here is this two-variable situation, the explicit characters Ψ_N for the specific series D_f , the location of the boundary, and its consequences for the explicit formula; a general natural-boundary criterion for products $\prod_p W(\text{Frob}_p; p^{-1}, p^{-s})$ presumably follows from Kurokawa–Moroz by the same accumulation argument, but we state only what we prove; we found no previous identification of $\text{Sym}^2 V$ or of a natural boundary in a singular-series average, including Goldston–Suriajaya’s series for $f(t) = t$, whose factor $G(s)$ is stated there as continuing to $\Re s > -1$ with no remark on the boundary.

Related work. The averaged and probabilistic forms of Bateman–Horn (Browning–Sofos–Teräväinen [8], Sofos [42], Banks–Ford [3], Bortolussi [7]) concern the main term and the count of primes, not the singular series of pairs; the singular-series averages of Kuperberg [30, 31] and of Kuperberg, Rodgers and Roditty–Gershon [32] are the linear and number-field cases, the latter with $\zeta_K(s+1)$ for the prime elements of K ; part I’s introduction places this paper among them.

Reader’s guide. The theorems are in Sections 2, 3 and 4; each is followed by a short paragraph “in words”. Section 5 is the foundation of part III. Appendix A lists every computation with inputs, outputs and cost.

Notation. χ_D denotes the *primitive* quadratic character attached to f , the Kronecker symbol of the fundamental discriminant D_0 with $\text{disc } f = D_0 m^2$; it is this character whose L -function has the functional equation, the Riemann hypothesis and the class-number value $L(1, \chi_D)$ used below. For $p \nmid \text{disc } f$ one has $\omega_f(p) = 1 + \chi_D(p)$; the finitely many primes dividing m change finitely many Euler factors, which are absorbed in $M_{f,K}$ (a second reading, 13 September 2026, noted that $f = t^2 + 3$, of discriminant -12 , needs this distinction). $f \in \mathbb{Z}[t]$ irreducible of degree n without fixed prime divisor; $G = \text{Gal}(f)$; $V = V_f$ its permutation representation on the roots, $\chi_V(g)$ the number of roots fixed by g ; $\omega_f(p) = \chi_V(\text{Frob}_p)$ for unramified p ; $K = \mathbb{Q}[t]/(f)$; $w = s + 1$; $u = 1/p$, $v = p^{-w}$; $P_p = 1 - \omega_f(p)^2/(p - \omega_f(p))^2$, $b(p) = (p - \omega_f(p))^{-2}$, $W_f(d) = db(d) \prod_{p \nmid d} P_p$ for squarefree d ; $a_f = W_f \omega_f$; $A_f = a_f * 1$; $D_f(s) = \sum_d a_f(d) d^{-s}$; $R_m(x) = \sum_{n \leq x} (x - n)^m A_f(n)$; $\rho = \beta + i\gamma$ a zero of the L -function in question; for a virtual character Ψ , $L(s, \Psi)$ is its Artin L -function.

2 The analytic structure of D_f

Throughout, $f \in \mathbb{Z}[t]$ is irreducible of degree n , without fixed prime divisor, $G = \text{Gal}(f)$, $V = V_f$ the permutation representation of G on the roots, $K = \mathbb{Q}[t]/(f)$, and for a prime p unramified in the splitting field, $g_p = \text{Frob}_p$ (a conjugacy class) with $\omega_f(p) = \chi_V(g_p)$ fixed points. We write $w = s + 1$, $u = u_p = p^{-1}$, $v = v_p = p^{-w}$. By part I, Theorem 3, $D_f(s) = \prod_p F_p(s)$ with

$$F_p(s) = P_p + p b(p) \omega_f(p) p^{-s} = \frac{1 - 2\omega_f(p)u + \omega_f(p)v}{(1 - \omega_f(p)u)^2} \quad (\omega_f(p) < p), \quad (4)$$

as one checks by writing $P_p = 1 - \omega^2 u^2 / (1 - \omega u)^2$ and $p b(p) \omega p^{-s} = \omega v / (1 - \omega u)^2$. The point of this section is that $F_p(s) \rightarrow 1 + \omega_f(p)v$ as $u \rightarrow 0$ with v fixed, and that $1 + \omega_f(p)v = 1 + \chi_V(g_p)v$ is *not* a finite product of Euler factors of Artin L -functions unless $\chi_V \leq 1$, i.e. unless f is linear.

2.1 The pair fields

Lemma 1 (pair fields). $\text{Sym}^2 V \cong V \oplus V^{(2)}$, where $V^{(2)}$ is the permutation representation of G on unordered pairs of distinct roots. Consequently

$$L(s, \text{Sym}^2 V) = \zeta_K(s) \prod_O \zeta_{K_O}(s),$$

the product over the G -orbits O on unordered pairs, K_O being the fixed field of the stabiliser of a pair in O . If G is 2-transitive there is one orbit, and $K^{(2)} = \mathbb{Q}(\alpha + \beta, \alpha\beta)$ has degree $n(n-1)/2$.

Proof. $\text{Sym}^2 V$ has the basis $e_\alpha e_\beta$ ($\alpha \leq \beta$ in any fixed order of the roots), which G permutes; the vectors e_α^2 span a copy of V and the vectors $e_\alpha e_\beta$, $\alpha \neq \beta$, span $V^{(2)}$. A permutation

representation $\text{Ind}_H^G \mathbf{1}$ has Artin L -function ζ_{L^H} , L the splitting field; the stabiliser of the pair $\{\alpha, \beta\}$ fixes $\alpha + \beta$ and $\alpha\beta$, and $\mathbb{Q}(\alpha + \beta, \alpha\beta)$ has the right degree when G is 2-transitive. $\square$

For quadratics, $\text{Sym}^2 V = \mathbf{1} \oplus \mathbf{1} \oplus \chi_D$ and $L(s, \text{Sym}^2 V) = \zeta(s)^2 L(s, \chi_D)$; for an S_3 -cubic $K^{(2)} \cong K$ and $L(s, \text{Sym}^2 V) = \zeta_K(s)^2$; for an S_4 - or A_4 -quartic $K^{(2)}$ is the resolvent sextic field.

In words. The symmetric square of the permutation representation, identified in part I as the source of the zeros, is the permutation action on *pairs* of roots. Every zero that appears below is therefore a zero of the Dedekind zeta function of an explicit number field: the field of one root, or the field of a pair of roots.

2.2 The plethystic exponents

Definition 2. For $N \geq 1$ and $g \in G$ put

$$\Psi_N(g) = \frac{1}{N} \sum_{j|N} \mu(j) (-1)^{N/j+1} \chi_V(g^j)^{N/j}.$$

Lemma 3. (i) Each Ψ_N is a virtual character of G , and $\Psi_1 = \chi_V$. (ii) For every $g \in G$, as formal power series in v ,

$$1 + \chi_V(g)v = \prod_{N \geq 1} \exp\left(\sum_{j \geq 1} \frac{\Psi_N(g^j)}{j} v^{Nj}\right) = \prod_{N \geq 1} \det(1 - g v^N \mid \Psi_N)^{-1},$$

where for a virtual character $\Psi = A - B$ we write $\det(1 - gX \mid \Psi)^{-1} = \det(1 - gX \mid A)^{-1} \det(1 - gX \mid B)$. (iii) $|\Psi_N(g)| \leq 2n^N/N$.

Proof. (ii) Taking logarithms, the claim is $\sum_{M \geq 1} c_M(g) v^M = \sum_{N,j} \Psi_N(g^j) v^{Nj}/j$ with $c_M(g) = (-1)^{M+1} \chi_V(g)^M/M$, i.e. $c_M(g) = \sum_{j|M} \Psi_{M/j}(g^j)/j$ for all M and g . Insert the definition: $\sum_{j|M} \frac{1}{j} \cdot \frac{j}{M} \sum_{i|M/j} \mu(i) (-1)^{M/(ij)+1} \chi_V(g^{ij})^{M/(ij)} = \frac{1}{M} \sum_{k|M} (-1)^{M/k+1} \chi_V(g^k)^{M/k} \sum_{i|k} \mu(i) = c_M(g)$, since only $k = 1$ survives. (i) Write $R(g, v) = 1 + \chi_V(g)v$; its coefficients lie in the representation ring $R(G)$. Suppose $\Psi_1, \dots, \Psi_{K-1}$ are virtual characters. The series $R(g, v) \prod_{N < K} \det(1 - g v^N \mid \Psi_N)$ has coefficients in $R(G)$ (products of the series $\det(1 - gX \mid A)^{\pm 1}$, whose coefficients are $\pm$ the characters of exterior or symmetric powers of A), and by (ii) it equals $1 + \Psi_K(g)v^K + O(v^{K+1})$; so $\Psi_K \in R(G)$. The case $N = 1$ is the definition. (iii) $|\chi_V(g^j)| \leq n$, and $\sum_{j|N} n^{N/j} \leq n^N + \sum_{j|N, j \geq 2} n^{N/j} \leq n^N + \sum_{k \leq N/2} n^k \leq n^N + 2n^{N/2} \leq 2n^N$ for $n \geq 2$; for $n = 1$ the claim is trivial. $\square$

For f linear, $\chi_V = 1$ and $1+v = (1-v^2)/(1-v)$: $\Psi_1 = 1$, $\Psi_2 = -1$, $\Psi_N = 0$ for $N \geq 3$, which is the factor $\zeta(s+1)/\zeta(2s+2)$ of the twin-prime case [20]. For $n \geq 2$ infinitely many Ψ_N are non-zero, because $1 + \chi_V(g)v$ with $\chi_V(g) \geq 2$ is not a finite product of cyclotomic polynomials in v .

In words. Let $p \rightarrow \infty$ in the local factor of D_f keeping $v = p^{-(s+1)}$ fixed: what remains is $1 + \omega v$, ω the number of roots of f modulo p . For $\omega = 1$ this is a ratio of two cyclotomic polynomials, which is why one factor $1/\zeta(2s+2)$ suffices in the twin-prime problem; for $\omega \geq 2$ no finite product of cyclotomic polynomials equals $1 + \omega v$, and infinitely many L -functions are needed, one for each Ψ_N . Lemma 3 says the Ψ_N are (virtual) characters, so these are Artin L -functions.

2.3 The structure theorem

For a virtual character Ψ of G let $L(s, \Psi)$ be its Artin L -function (a quotient of two Artin L -functions of genuine representations, hence meromorphic in $\mathbb{C}$), with local factor $L_p(s, \Psi) = \det(1 - g_p p^{-s} \mid \Psi)^{-1}$ at unramified p ; and let S be the finite set of primes ramified in the splitting field or dividing the leading coefficient of f .

Theorem 4 (structure of D_f). *For every $K \geq 1$,*

$$D_f(s) = \prod_{N=1}^K L(N(s+1), \Psi_N) \cdot M_{f,K}(s), \quad M_{f,K}(s) = \prod_p F_p(s) \prod_{N=1}^K L_p(N(s+1), \Psi_N)^{-1}, \quad (5)$$

where the Euler product $M_{f,K}$ converges absolutely and locally uniformly in $\Re s > -1 + 1/(K+1)$, so that $M_{f,K}$ is holomorphic there. In particular $L(s+1, \Psi_1) = \zeta_K(s+1)$, so that $E_f = \prod_{N=2}^K L(Nw, \Psi_N) M_{f,K}$; D_f is meromorphic in $\Re s > -1$; and in $\Re s > -1 + 1/(K+1)$ its poles are among the poles of $\prod_{N \leq K} L(Nw, \Psi_N)$, while its zeros that are not zeros of that product are exactly the zeros of the local factors F_p , $p \notin S$, together with those of the finitely many local factors at $p \in S$.

Proof. Fix K and a compact set in $\Re w > 1/(K+1)$, on which $\Re w \geq \eta > 1/(K+1)$. For $p \notin S$ the local factor of $M_{f,K}$ is, by Lemma 3(ii),

$$\begin{aligned} M_p(s) &= \frac{F_p(s)}{1 + \omega v} \cdot \frac{1 + \omega v}{\prod_{N \leq K} \det(1 - g_p v^N \mid \Psi_N)^{-1}} \\ &= \frac{1 - 2\omega u/(1 + \omega v)}{(1 - \omega u)^2} \cdot \exp\left(\sum_{M > K} d_M(g_p) v^M\right), \end{aligned}$$

with $\omega = \omega_f(p)$ and $d_M(g) = \sum_{N \mid M, N > K} \Psi_N(g^{M/N}) N/M$, so $|d_M| \leq 2n^M \tau(M)$ by Lemma 3(iii). For p so large that $n|v| \leq np^{-\eta} \leq \frac{1}{2}$ and $\omega u \leq \frac{1}{4}$ we get $|\sum_{M > K} d_M v^M| \leq \sum_{M > K} 2\tau(M)(n|v|)^M \ll_K (n|v|)^{K+1} \ll p^{-(K+1)\eta}$, and $\frac{1 - 2\omega u/(1 + \omega v)}{(1 - \omega u)^2} = 1 + \frac{2\omega^2 uv}{1 + \omega v} + O(u^2) = 1 + O(p^{-1-\eta}) + O(p^{-2})$. Since $(K+1)\eta > 1$, $\sum_p |M_p(s) - 1|$ converges uniformly on the compact set, which gives absolute and locally uniform convergence of $\prod_{p \notin S, p > p_0} M_p$; the finitely many remaining factors are rational functions of p^{-s} whose only possible poles are at $|p^{-Nw}| = 1$, i.e. on $\Re w = 0$. A uniformly convergent product of non-vanishing holomorphic factors is holomorphic and non-vanishing; hence $M_{f,K}$ vanishes exactly where a local factor does, and $M_p(s) = 0$ iff $F_p(s) = 0$ (the other factors of M_p are $\det(1 - g_p v^N \mid \Psi_N)$, non-zero for $\Re w > 0$). Letting $K \rightarrow \infty$ gives the continuation to $\Re s > -1$, the finitely many $L(Nw, \Psi_N)$ being meromorphic by Brauer's theorem. $\square$

In words. D_f is a product of finitely many Artin L -functions evaluated at $N(s+1)$, times a remainder that is harmless in a half-plane that grows with the number of factors extracted but never reaches $\Re s = -1$. Every singularity of D_f to the right of -1 is a zero or pole of an L -function of the Galois group of f , at a predictable place.

Remark 5. $M_{f,K}$ is not $1 + O(p^{-2})$ at the level of individual primes: the mixed term $2\omega^2 uv/(1 + \omega v)$ is of size $p^{-1-\Re w}$. In the numerical work these terms are extracted as well ($[\zeta(w+1)L(w+1, \chi_D)]^4$, $[\zeta(2w+1)L(2w+1, \chi_D)]^{-8}$, ... for quadratics) so that the remaining product converges like $\sum p^{-2}$; see `research/paper-II/scripts/explicit-diag.py`.

2.4 Quadratics: the exponents in closed form

For a quadratic f of discriminant D , $G = \{e, \tau\}$, $\chi_V = \mathbf{1} + \chi_D$, and every virtual character is $a\mathbf{1} + b\chi_D$ with $a, b \in \mathbb{Z}$. Write $\Psi_N = a_N\mathbf{1} + b_N\chi_D$, so that $L(Nw, \Psi_N) = \zeta(Nw)^{a_N} L(Nw, \chi_D)^{b_N}$, and let $M_N(x) = \frac{1}{N} \sum_{j|N} \mu(N/j) x^j$ be the necklace polynomial.

Proposition 6. *For every $N \geq 1$,*

$$a_N + b_N = \Psi_N(e) = -M_N(-2), \quad a_N - b_N = \Psi_N(\tau) = \frac{1}{N} \sum_{\substack{j|N \\ 2 \nmid j}} \mu(j) (-1)^{N/j+1} 2^{N/j}.$$

In particular $a_N = b_N = \frac{1}{2}M_N(2)$ for odd N , and for even N $a_N + b_N = -M_N(-2) < 0$; the first values are those of Table 1, and $a_1 = b_1 = 1$, i.e. $L(w, \Psi_1) = \zeta_K(w)$; $\Psi_2 = -\text{Sym}^2 V$; $a_3 = b_3 = 1$, so that $\zeta(3w)L(3w, \chi_D)$ appears in the numerator and D_f has a simple pole at $s = -\frac{2}{3}$; and $a_N > 0$ for all odd N , giving a pole of order $a_N = \frac{1}{2}M_N(2)$ at $s = -1 + 1/N$.

Proof. $\chi_V(e^j) = 2$ and $\chi_V(\tau^j) = 2$ or 0 according as j is even or odd; insert into Definition 2, using $(-1)^{k+1}2^k = -(-2)^k$ for the first formula. For odd N there is no even divisor, so $a_N = b_N = -\frac{1}{2}M_N(-2) = \frac{1}{2}M_N(2)$; $M_N(2) > 0$ counts binary Lyndon words. $\square$

Table 1: The exponents (a_N, b_N) of $\zeta(Nw)^{a_N} L(Nw, \chi_D)^{b_N}$ in the structure theorem for a quadratic f (Proposition 6; `research/paper-II/scripts/explicit-diag.py`). For odd N , $a_N = b_N = \frac{1}{2}M_N(2)$ is the number of binary Lyndon words of length N .

N	1	2	3	4	5	6	7	8	9	10	11	12
a_N	1	-2	1	-1	3	-6	9	-14	28	-54	93	-165
b_N	1	-1	1	-2	3	-5	9	-16	28	-51	93	-170

In words. For a quadratic the L -functions are ζ and $L(s, \chi_D)$ and the exponents are the necklace numbers of Table 1. Two features matter later: $\zeta(3s+3)$ in the numerator, which gives a real pole at $s = -\frac{2}{3}$ and hence a term $x^{m-2/3}$ in every average of S_f ; and $1/(\zeta(2s+2)^2 L(2s+2, \chi_D))$, whose poles at half the zeros give the oscillating terms $x^{\rho/2}$.

2.5 The natural boundary

Theorem 7 (natural boundary). *Let f be irreducible of degree $n \geq 1$ without fixed prime divisor. Then D_f has infinitely many zeros in $-1 < \Re s < 0$, accumulating at every point of the line $\Re s = -1$, and $\Re s = -1$ is a natural boundary of D_f and of $\sum_n A_f(n) n^{-s} = \zeta(s) D_f(s)$. The same holds for $f(t) = t$, i.e. for the generating Dirichlet series of the twin-prime singular series.*

Proof. By (4), for $p \notin S$ with $\omega = \omega_f(p) \geq 1$, $F_p(s) = 0$ iff $\omega v = -(1 - 2\omega u)$, i.e. iff $p^{-w} = -(1 - 2\omega/p)/\omega$. These are the points

$$s_{p,j} = -1 + \frac{\log \omega - \log(1 - 2\omega/p)}{\log p} + \frac{(2j+1)\pi i}{\log p}, \quad j \in \mathbb{Z},$$

with $\Re s_{p,j} \rightarrow -1^+$ as $p \rightarrow \infty$ (for $\omega = 1$, $\Re s_{p,j} = -1 + 2/(p \log p) + O(p^{-2})$). By the Chebotarev density theorem there are infinitely many p with $\omega_f(p) \geq 1$ (indeed with $\omega_f(p) = n$). Given $t_0 \in \mathbb{R}$ and $\varepsilon > 0$, choose such a p with $\log p > \pi/\varepsilon$ and $|\Re s_{p,j} + 1| < \varepsilon$; the spacing of the $\Im s_{p,j}$ is $2\pi/\log p < 2\varepsilon$, so some $s_{p,j}$ lies within 2ε of $-1 + it_0$. By Theorem 4 with K large, in a

neighbourhood of $s_{p,j}$ we have $D_f = \Lambda_K \cdot M_{f,K}$ with $\Lambda_K = \prod_{N \leq K} L(Nw, \Psi_N)$ meromorphic and $M_{f,K}(s_{p,j}) = 0$; so $s_{p,j}$ is a zero of D_f unless it is a pole of Λ_K . The poles of Λ_K form a discrete set, so all but finitely many of the zeros $s_{p,j}$ in any bounded region are zeros of D_f . Now suppose D_f continued meromorphically to a disc B around $-1 + it_0$. Zeros of a meromorphic function accumulate only at poles, so $-1 + it_0$ would be a pole; but in a punctured neighbourhood of a pole $|D_f|$ is large, whereas zeros $s_{p,j}$ lie arbitrarily close to $-1 + it_0$: a contradiction. For $\zeta(s)D_f(s)$ the same argument applies, since $\zeta(s) \neq 0$ on $\Re s = -1$. For $f(t) = t$, $F_p = (1 - 2u + v)/(1 - u)^2$ vanishes at $p^{-w} = -(1 - 2/p)$ and the argument is identical. $\square$




Figure 1: The analytic structure behind the explicit formula (Theorems 4–7, Proposition 6). The real poles at $s = -1 + 1/k$, k odd, have orders $1, 3, 9, 28, \dots$; the first zero family sits on $\Re s = -\frac{3}{4}$ at half the ordinates of ζ (double poles) and of $L(s, \chi_{-4})$ (simple), the second on $\Re s = -\frac{7}{8}$; the zeros of the local factors F_p (grey) march towards the dashed line $\Re s = -1$, which is a natural boundary. The contour of the explicit formula stops just left of $-\frac{3}{4}$.

Remark 8 (no asymptotic explicit formula). Bhowmik and Schlage-Puchta [6] proved that a Dirichlet series with a natural boundary at $\Re s = \sigma_0$ admits no asymptotic expansion of its summatory function by residue terms with error $O(x^{\sigma_0 - \varepsilon})$. Applied to $\zeta(s)D_f(s)$ (subject to checking their growth hypotheses in our setting), this says that for every $\varepsilon > 0$ the Riesz mean $R_m(x)$ minus the sum of *all* residues in $\Re s > -1 + \varepsilon$ is $\Omega(x^{m-1-\varepsilon})$: the error term $O(x^{m-1+\varepsilon})$ of the twin-prime explicit formula [20] is best possible in this strong sense, and so is ours.

In words. Each local factor vanishes on a lattice of points whose real parts tend to -1 and whose imaginary parts become dense, so D_f cannot be continued across $\Re s = -1$. This is why the explicit formula below is conditional while the twin-prime formula of Goldston and Suriajaya is not: they could move the contour up to the wall without crossing anything dangerous; for non-linear f the L -factors with $N \geq 3$ pile up in front of it.

Remark 9. For $f(t) = t$ the pure- v part of the local factor is $1 + v = (1 - v^2)/(1 - v)$, so the L -function part of D_f is just $\zeta(w)/\zeta(2w)$ and the contour in [20] can be moved to $\Re s = -1 + \varepsilon$, crossing the whole critical strip of $\zeta(2s + 2)$: this is why the linear case has an *unconditional* explicit formula with error $O(x^{m-1+\varepsilon})$, and why that error term cannot be improved by contour

shifting. For $n \geq 2$ the factors $L(Nw, \Psi_N)$ with $N \geq 3$ pile up between $\Re s = -\frac{3}{4}$ and $\Re s = -1$, and the contour can only be moved a little past the first zero family.

3 The explicit formula for the diagonal

Let $R_m(x) = \sum_{n \leq x} (x-n)^m A_f(n)$, so that for $\Re s > 1$

$$R_m(x) = \frac{1}{2\pi i} \int_{(2)} G_m(s) x^{s+m} ds, \quad G_m(s) = \frac{m! \zeta(s) D_f(s)}{s(s+1) \cdots (s+m)}. \quad (6)$$

From now on f is quadratic with discriminant D , so that by Theorem 4 and Proposition 6, for every $K \geq 4$,

$$D_f(s) = \frac{\zeta(w) L(w, \chi_D) \zeta(3w) L(3w, \chi_D)}{\zeta(2w)^2 L(2w, \chi_D)} \prod_{N=4}^K \zeta(Nw)^{a_N} L(Nw, \chi_D)^{b_N} M_{f,K}(s), \quad w = s+1. \quad (7)$$

3.1 Statement

Theorem 10. *Let f be an irreducible quadratic of discriminant D and $m \geq 2$. Assume the Riemann hypothesis for ζ and for $L(s, \chi_D)$. Then there is $\varepsilon_0 \in (0, \frac{1}{100})$ such that for $x \geq 2$ and some $U \in [x^{16}, 2x^{16}]$,*

$$R_m(x) = \frac{D_f(1)}{m+1} x^{m+1} + x^m \left(-\frac{\log x}{2C(f)} + c_{f,m} \right) + c'_{f,m} x^{m-2/3} + x^{m-1} \sum_{\substack{\rho=\beta+i\gamma \\ |\gamma| \leq 2U}} Q_{m,\rho}(\log x) x^{\rho/2} + O(x^{m-3/4-\varepsilon_0}), \quad (8)$$

where ρ runs over the zeros of $\zeta(s)^2 L(s, \chi_D)$ (all with $\beta = \frac{1}{2}$), $Q_{m,\rho}$ is the polynomial (of degree less than the order of the pole of G_m at $\rho/2 - 1$, i.e. of degree 0 at simple zeros of L and 1 at simple zeros of ζ) such that $Q_{m,\rho}(\log x) x^{\rho/2-1+m}$ is the residue of $G_m(s) x^{s+m}$ at $s = \rho/2 - 1$, and

$$c'_{f,m} = \operatorname{Res}_{s=-2/3} G_m(s) = \frac{m! \zeta(-\frac{2}{3}) \zeta(\frac{1}{3}) L(\frac{1}{3}, \chi_D) L(1, \chi_D)}{3 \zeta(\frac{2}{3})^2 L(\frac{2}{3}, \chi_D)} \times \prod_{N=4}^K \zeta(\frac{N}{3})^{a_N} L(\frac{N}{3}, \chi_D)^{b_N} \frac{M_{f,K}(-\frac{2}{3})}{(-\frac{2}{3})(\frac{1}{3}) \cdots (m-\frac{2}{3})},$$

$c_{f,m}$ being the constant term of the residue at $s = 0$. In $c'_{f,m}$ the truncation $K \geq 4$ is arbitrary: the finite product over $4 \leq N \leq K$ and the factor $M_{f,K}(-\frac{2}{3})$ each depend on K , their product does not, and the product over all $N \geq 4$ alone diverges, because $|\Psi_N(e)| \asymp 2^N/N$ makes the local factors at the small split primes grow (a second reading, 13 September 2026, found the product written without the truncation). The implied constant depends on f , m and ε_0 .

Remark 11 (the class number in the term $x^{m-2/3}$). The coefficient $c'_{f,m}$ contains the factor $L(1, \chi_D)$, which for $D < 0$ equals $2\pi h(D)/(w_D \sqrt{|D|})$ by Dirichlet's class number formula: the class number of $\mathbb{Q}(\sqrt{D})$ enters the second moment of the pair singular series of f through the real term $x^{m-2/3}$, alongside the values $\zeta(\frac{1}{3})$, $L(\frac{1}{3}, \chi_D)$ and the Euler product $M_{f,K}(-\frac{2}{3})$.

The coefficient of $x^m \log x$ is $-1/(2C(f))$ because $\operatorname{Res}_{s=0} D_f = 1/C(f)$ (part I, Theorem 3) and $\zeta(0) = -\frac{1}{2}$; explicitly $c_{f,m} = -\frac{1}{2C(f)} (\log 2\pi - H_m) - \frac{1}{2} \kappa_f$ with κ_f the constant term of the Laurent expansion of D_f at 0 and $H_m = \sum_{k \leq m} 1/k$.

In words. After the expected terms x^{m+1} , $x^m \log x$, x^m comes an unexpected real term $x^{m-2/3}$, then oscillating terms of size $x^{m-3/4}$ with frequencies $\gamma/2$ in $\log x$, γ the ordinates of the zeros of ζ and $L(s, \chi_D)$; the rest is smaller by a fixed power of x . The hypothesis is needed because the proof evaluates $1/\zeta$ and $1/L$ just left of the critical line.

3.2 Growth lemmas

We use the following consequences of the Riemann hypothesis for ζ and $L(s, \chi_D)$ ([44, Chapter 14] for ζ ; the proofs transfer verbatim to $L(s, \chi_D)$, see [27, §5.7]). Let $\Lambda \in \{\zeta, L(\cdot, \chi_D)\}$ and $t \geq 2$.

(G1) For every fixed $\sigma > \frac{1}{2}$: $\Lambda(\sigma + it)^{\pm 1} \ll t^\varepsilon$; for every fixed σ : $\Lambda(\sigma + it) \ll t^{\max(0, 1/2 - \sigma) + \varepsilon}$, and for fixed $\sigma < \frac{1}{2}$: $\Lambda(\sigma + it)^{-1} \ll t^\varepsilon$ (by the functional equation, $|\Lambda(\sigma + it)| \asymp t^{1/2 - \sigma} |\Lambda(1 - \sigma - it)|$).

(G2) Unconditionally, for $-1 \leq \sigma \leq 2$: $\frac{\Lambda'}{\Lambda}(\sigma + it) = \sum_{|\gamma - t| \leq 1} \frac{1}{\sigma + it - \rho} + O(\log t)$, the sum over zeros $\rho = \beta + i\gamma$ of Λ , and the number of zeros with $|\gamma - t| \leq 1$ is $\leq c_0 \log t$.

Lemma 12 (horizontal segments). *Assume RH for ζ and $L(s, \chi_D)$. For every $\delta > 0$ there is $\varepsilon_1(\delta) > 0$ such that for all $T \geq T_0(\delta)$ and $\varepsilon_0 \leq \varepsilon_1$, there exists $U \in [T, 2T]$ with*

$$|\zeta(\sigma + 2iU)^{-1}| + |L(\sigma + 2iU, \chi_D)^{-1}| \leq T^\delta \quad \text{uniformly for } \frac{1}{2} - 2\varepsilon_0 \leq \sigma \leq 2.$$

Proof. Let Λ be one of the two functions, $\sigma_1 = \frac{1}{2} + \varepsilon_0$, and $Z = [\frac{1}{2} - 2\varepsilon_0, \sigma_1]$ (the “zone”). For $\sigma \geq \sigma_1$ the claim is (G1) with $\delta/2$ in place of ε . For $\sigma \in Z$ and $t = 2U$ write, by (G2),

$$\begin{aligned} \log \frac{1}{|\Lambda(\sigma + it)|} &= \log \frac{1}{|\Lambda(\sigma_1 + it)|} + \Re \int_{\sigma}^{\sigma_1} \frac{\Lambda'}{\Lambda}(\sigma' + it) d\sigma' \\ &\leq \frac{\delta}{4} \log T + \sum_{|\gamma - t| \leq 1} \int_Z \frac{d\sigma'}{|\sigma' + it - \rho|} + 3\varepsilon_0 C \log t. \end{aligned}$$

Under RH $\beta = \frac{1}{2}$, so $|\sigma' + it - \rho| = \sqrt{(\sigma' - \frac{1}{2})^2 + (t - \gamma)^2}$ and, with $\theta = |t - \gamma|$, $\int_Z \leq \int_{-2\varepsilon_0}^{2\varepsilon_0} \frac{dy}{\sqrt{y^2 + \theta^2}} = 2 \operatorname{arsinh}(2\varepsilon_0/\theta)$, which is $\leq 4\varepsilon_0/\theta$ for $\theta \geq 2\varepsilon_0$ and $\leq 2 \log(4\varepsilon_0/\theta) + 2$ for $\theta < 2\varepsilon_0$. The zeros with $2\varepsilon_0 \leq \theta \leq 1$ contribute at most $4\varepsilon_0 T_\Lambda(t)$, $T_\Lambda(t) = \sum_{2\varepsilon_0 \leq |\gamma - t| \leq 1} |t - \gamma|^{-1}$, and those with $\theta < 2\varepsilon_0$ at most $2\Sigma_\Lambda(t) + 2\#\{\gamma : |\gamma - t| < 2\varepsilon_0\}$ with $\Sigma_\Lambda(t) = \sum_{|\gamma - t| < 2\varepsilon_0} \log(4\varepsilon_0/|t - \gamma|)$. (A pointwise bound for $T_\Lambda(t)$ from the zero count alone loses the factor ε_0 : a dyadic block $[\theta_0, 2\theta_0]$ of length $\theta_0 \ll 1/\log t$ may contain $\asymp \log t$ zeros, so the dyadic sum is $\ll \log t$ and not $\ll \varepsilon_0 \log(1/\varepsilon_0) \log t$; the first version of this proof made that mistake, found by a second reading on 13 September 2026. The average over t is what is needed, for all three quantities.) Average over $t \in [2T, 4T]$: each zero $\gamma \in [2T - 1, 4T + 1]$ contributes to $\int T_\Lambda dt$ at most $\int_{2\varepsilon_0 \leq |y| \leq 1} dy/|y| = 2 \log(1/2\varepsilon_0)$, to $\int \Sigma_\Lambda dt$ at most $\int_{-2\varepsilon_0}^{2\varepsilon_0} \log(4\varepsilon_0/|y|) dy = 4\varepsilon_0(1 + \log 2)$, and to $\int \# dt$ at most $4\varepsilon_0$, and there are $\leq c'_0 T \log T$ such zeros. Hence the averages over $[2T, 4T]$ of $4\varepsilon_0 T_\Lambda$, $2\Sigma_\Lambda$ and $2\#$ are each $\leq c_2 \varepsilon_0 \log(1/\varepsilon_0) \log T$ with c_2 absolute. By Markov's inequality the set of $t \in [2T, 4T]$ on which the sum of these six quantities (three for ζ , three for L) is at most $30c_2 \varepsilon_0 \log(1/\varepsilon_0) \log T$ has measure $\geq \frac{4}{5} \cdot 2T$; pick U with $2U$ in it. Altogether $\log(1/|\Lambda(\sigma + 2iU)|) \leq (\frac{\delta}{4} + c_1 \varepsilon_0 \log(1/\varepsilon_0)) \log T$ for $\sigma \in Z$, with c_1 absolute, which is $\leq \delta \log T$ once $\varepsilon_0 \leq \varepsilon_1(\delta)$. $\square$

3.3 Proof of Theorem 10

Fix $K = 15$ (any $K \geq 4$ with $(K + 1)(\frac{1}{4} - \varepsilon_0) > 1$ works), $\delta = \varepsilon_0$ and $\varepsilon_0 \leq \min(\varepsilon_1(\varepsilon_0), \frac{1}{100})$, which is possible since $\varepsilon_1(\delta) \gg \delta/\log(1/\delta)$. Let U be as in Lemma 12 with $T = x^{16}$, and let

$\mathcal{R}$ be the rectangle with vertices $2 \pm iU$ and $-\frac{3}{4} - \varepsilon_0 \pm iU$. By (6) (truncating the integral at height U costs $\ll x^{m+2}U^{-m}$, since $G_m(2+it) \ll t^{-m-1}$) and the residue theorem,

$$R_m(x) = \sum_{\text{poles in } \mathcal{R}} \text{Res}(G_m(s)x^{s+m}) + \frac{1}{2\pi i} \left(\int_{-3/4-\varepsilon_0-iU}^{-3/4-\varepsilon_0+iU} + \int_{\text{horiz.}} \right) G_m(s)x^{s+m} ds + O(x^{m+2}U^{-m}).$$

Poles. By (7) and Theorem 4, in $\Re s \geq -\frac{3}{4} - \varepsilon_0 > -\frac{4}{5}$ the poles of G_m are: $s = 1$ (simple, from $\zeta(s)$; $D_f(1) = \sum_d a_f(d)/d$), $s = 0$ (double: the pole of $\zeta(w)$ and the factor $1/s$), $s = -\frac{2}{3}$ (simple, from $\zeta(3w)$), and $s = \rho/2 - 1$ for the zeros ρ of $\zeta(2w)^2 L(2w, \chi_D)$ with $|\gamma| \leq 2U$; the factors $\zeta(Nw)^{a_N} L(Nw, \chi_D)^{b_N}$, $N \geq 4$, have $\Re(Nw) \geq 1 - 4\varepsilon_0 > \frac{1}{2}$ there, so under RH they have neither zeros nor poles except $\zeta(Nw)$ at $Nw = 1$, i.e. $s = -1 + 1/N \leq -\frac{3}{4}$, outside $\mathcal{R}$ for $N \geq 4$ (and $N = 4$ gives a zero of D_f); $M_{f,K}$ is holomorphic in $\Re s > -1 + \frac{1}{16}$; and $s = -1, \dots, -m$ are outside $\mathcal{R}$. The residues at 1, 0, $-\frac{2}{3}$ are the three explicit terms of (8), and the residue at $\rho/2 - 1$ is $Q_{m,\rho}(\log x)x^{\rho/2-1+m}$ by definition.

Vertical line. On $\Re s = -\frac{3}{4} - \varepsilon_0$ (so $\Re w = \frac{1}{4} - \varepsilon_0$, $\Re 2w = \frac{1}{2} - 2\varepsilon_0$, $\Re 3w = \frac{3}{4} - 3\varepsilon_0$, $\Re Nw \geq 1 - 4\varepsilon_0$ for $N \geq 4$), (G1) gives, for $|t| \geq 2$, $\zeta(s) \ll |t|^{5/4+\varepsilon_0+\varepsilon}$, $\zeta(w)L(w, \chi_D) \ll |t|^{1/2+2\varepsilon_0+\varepsilon}$, $\zeta(2w)^{-2}L(2w, \chi_D)^{-1} \ll |t|^\varepsilon$, $\zeta(3w)L(3w, \chi_D) \ll |t|^\varepsilon$, $\prod_{N \geq 4} \ll |t|^\varepsilon$, and $M_{f,K} \ll 1$; hence $G_m(s) \ll |t|^{7/4+3\varepsilon_0+\varepsilon-(m+1)}$, and for $m \geq 2$ the exponent is < -1 . The vertical integral is therefore $\ll x^{m-3/4-\varepsilon_0}$.

Horizontal segments. On $\Im s = \pm U$, $-\frac{3}{4} - \varepsilon_0 \leq \sigma \leq 2$: (G1) bounds $\zeta(s)$, $\zeta(w)L(w)$, $\zeta(3w)L(3w)$ and the $N \geq 4$ factors by $U^{7/4+3\varepsilon_0+\varepsilon}$ (worst case at the left end), Lemma 12 bounds $\zeta(2w)^{-2}L(2w, \chi_D)^{-1}$ by $U^{3\varepsilon_0}$, and $M_{f,K} \ll 1$; the kernel is $\ll U^{-m-1}$ and $|x^{s+m}| \leq x^{m+2}$. The two segments contribute $\ll x^{m+2}U^{7/4+6\varepsilon_0+\varepsilon-m-1} \leq x^{m+2}U^{-5/4+7\varepsilon_0}$ for $m \geq 2$, and the truncation of the original integral at height U contributes $\ll x^{m+2}U^{-m}$; with $U \geq x^{16}$ and $\varepsilon_0 \leq \frac{1}{100}$ both are $\ll x^{m-1}$.

Collecting, $R_m(x)$ equals the residue sum plus $O(x^{m-3/4-\varepsilon_0})$, which is (8). $\square$

Remark 13 (on the zero sum). The sum over $|\gamma| \leq 2U$ is not absolutely convergent as $U \rightarrow \infty$ with the polynomial growth of the coefficients; as in [20, Theorem 3], simplicity of the zeros and a bound $\sum_{|\gamma| \leq T} |\zeta'(\rho)|^{-2} \ll T$ (Gonek–Hejhal) would make $x^{m-3/4} \sum_\gamma Q_{m,\rho} x^{i\gamma/2}$ absolutely convergent for $m \geq 2$. Numerically the coefficients decay like $|\gamma|^{-5/4}$ for $m = 2$.

3.4 General f

Theorem 14 (general f). *Let f be irreducible of degree $n \geq 2$ and $m \geq 2$. Assume that the Artin L -functions of the irreducible constituents of $\Psi_1 = \chi_V$, $\Psi_2 = -\chi_{\text{Sym}^2 V}$, $\Psi_3 = \frac{1}{3}(\chi_V^3 - \psi^3 \chi_V)$ and $\Psi_4 = \frac{1}{4}(\psi^2 \chi_V^2 - \chi_V^4)$ are entire (automatic when G is an M -group, e.g. S_3, S_4, A_4, D_n) and satisfy the Riemann hypothesis. Let $m_N = \langle \Psi_N, \mathbf{1} \rangle$ be the multiplicity of the trivial character in Ψ_N , so that $L(N(s+1), \Psi_N)$ has at $s = -1 + 1/N$ a pole of order m_N if $m_N > 0$ and a zero of order $-m_N$ if $m_N < 0$. Then there is $\varepsilon_0 > 0$ such that, for $U \in [x^{16}, 2x^{16}]$ suitably chosen,*

$$R_m(x) = \frac{D_f(1)}{m+1} x^{m+1} + x^m \left(-\frac{\log x}{2C(f)} + c_{f,m} \right) + \sum_{N \in \{3,4\}} x^{m-1+1/N} P_{f,m,N}(\log x) + x^{m-1} \sum_{\substack{\rho \\ |\gamma| \leq 2U}} Q_{m,\rho}(\log x) x^{\rho/2} + O(x^{m-3/4-\varepsilon_0}),$$

where ρ runs over the zeros of $L(s, \text{Sym}^2 V) = \zeta_K(s) \prod_O \zeta_{K_O}(s)$, and $P_{f,m,N}$ is the polynomial (possibly zero) given by the residue at $s = -1 + 1/N$: if the integrand has there a pole of order $r_N \geq 1$, then $P_{f,m,N}$ has degree exactly $r_N - 1$, and $P_{f,m,N} = 0$ if $r_N \leq 0$. Under the hypotheses $r_3 = m_3$, since the other factors are finite and non-zero at $s = -\frac{2}{3}$; $r_4 = m_4 + k$,

where $k \geq 0$ is the order of vanishing of $L(\frac{1}{2}, \text{Sym}^2 V)$; and $s = -\frac{1}{2}$ is never a pole, since $m_2 = -\dim(\text{Sym}^2 V)^G \leq -1$. For $G = S_n$ one has $m_3 = 1$ and $m_4 = -2$ (-1 for $n = 2$), so the smooth part beyond x^m is a single term $c'_{f,m} x^{m-2/3}$, as for quadratics. For $G = C_3$ one has $m_3 = 2$, and the term is $x^{m-2/3}(a \log x + b)$ with $a \neq 0$ (Table 2).

Proof. Identical to the proof of Theorem 10, with $\zeta(w)L(w, \chi_D)$ replaced by $L(w, \Psi_1) = \zeta_K(w)$, $\zeta(2w)^{-2}L(2w, \chi_D)^{-1}$ by $L(2w, \Psi_2) = 1/L(2w, \text{Sym}^2 V)$, and $\zeta(3w)L(3w, \chi_D)$ by $L(3w, \Psi_3)$: (G1) and (G2) hold for entire Artin L -functions satisfying the Riemann hypothesis with the same proofs (Hadamard product and functional equation), the factors $L(Nw, \Psi_N)$, $N \geq 4$, are evaluated at $\Re(Nw) \geq 1 - 4\varepsilon_0$ where they are $\ll |t|^\varepsilon$ and free of zeros and poles, and $M_{f,K}$ is bounded by Theorem 4. The multiplicity of the trivial character in Ψ_3 is $\frac{1}{3}(\langle \chi_V^3, 1 \rangle - \langle \psi^3 \chi_V, 1 \rangle)$; for S_n , $n \geq 3$, $\langle \chi_V^3, 1 \rangle = 5$ (orbits of S_n on maps $\{1, 2, 3\} \rightarrow \{1, \dots, n\}$) and $\langle \psi^3 \chi_V, 1 \rangle$ is the mean number of fixed points of g^3 , i.e. $\mathbb{E}[c_1 + 3c_3] = 1 + 3 \cdot \frac{1}{3} = 2$ with c_k the number of k -cycles; for $n = 2$ the values are 4 and 1.

The residue at $s_0 = -1 + 1/N$ is the one place where the general case differs from the quadratic one. If the integrand has the Laurent expansion $\sum_{k=1}^r a_{-k}(s - s_0)^{-k} + O(1)$ with $a_{-r} \neq 0$, then the residue of its product with x^{s+m} is

$$x^{s_0+m} \sum_{k=1}^r \frac{a_{-k}}{(k-1)!} (\log x)^{k-1},$$

a polynomial of degree exactly $r - 1$ in $\log x$. At $s = -\frac{2}{3}$ the factors $\zeta_K(w)$, $L(2w, \Psi_2) = 1/L(\frac{2}{3}, \text{Sym}^2 V)$, $L(Nw, \Psi_N)$ for $N \geq 4$ (evaluated at $\Re(Nw) \geq \frac{4}{3}$) and $M_{f,K}$ are finite and non-zero under the hypotheses, so $r_3 = m_3$; at $s = -\frac{3}{4}$ the factor $L(2w, \Psi_2) = 1/L(\frac{1}{2}, \text{Sym}^2 V)$ may have a pole, whence $r_4 = m_4 + k$. That the multiplicity m_3 can exceed 1 was pointed out to us in an external assessment of this work (12 September 2026): for $f = t^3 - 3t - 1$, irreducible with discriminant 81 and Galois group C_3 , one has $\chi_V = (3, 0, 0)$ on the three group elements, $\Psi_3 = (8, -1, -1)$ and $m_3 = 2$, so $L(3w, \Psi_3)$ contains $\zeta(3w)^2$ and the pole at $s = -\frac{2}{3}$ is double. An earlier version of this theorem wrote the term as $c'_{f,m} x^{m-2/3}$ with a constant for every f , which is wrong for this polynomial and for every group in Table 2 other than S_n , A_n and A_4 . For $t^3 - 3t - 1$ the coefficient of $x^{m-2/3} \log x$ is computable from the residue machinery (`scripts/predict-cubic.py`, which reproduces $D_f(1)$ to $5 \cdot 10^{-12}$ at $s = 1$ as a check and is independent of the truncation K): $a = -0.00234$ at $m = 2$ and -0.00300 at $m = 3$, with $b = -0.0049$ and -0.0050 . It is four orders of magnitude below the zero terms $x^{m-3/4}$ at accessible x , so the correction is established by the multiplicity computation and cannot be seen in data to $x = 10^7$; for $t^2 + 1$ the corresponding constant is -0.455 . $\square$

In words. In any degree the same formula holds with the zeros of the Dedekind zeta functions of the root field and of the pair fields, under the corresponding Riemann hypotheses; the real term $x^{m-2/3}$ is universal, with a constant coefficient for the generic Galois group and a polynomial in $\log x$ of degree $m_3 - 1$ otherwise, already of degree 1 for cyclic cubics.

3.5 $f = t^2 + 1$: every coefficient

For $f = t^2 + 1$, $D = -4$, $\chi_D = \chi_{-4}$, $K = \mathbb{Q}(i)$, and every quantity in (8) is computable: $D_f(1) = 0.7507908137$, $C(f) = 1.372813462821$ (from $L(1, \chi_{-4}) = \pi/4$ and $M_{f,K}(0)$; part I quotes $1.3728134628\dots$), $c_{f,2} = -0.454974899864$, $c_{f,3} = -0.333569719058$, and the coefficients $Q_{m,\rho}$ for the zeros with $\gamma \leq 100$ are computed as Taylor coefficients of $(s - \rho/2 + 1)^{\text{ord}} G_m(s)$ by trapezoidal Cauchy integrals on circles of radius 0.015 (`scripts/explicit-diag.py`; $M_{f,K}$ with $K = 15$ and primes up to $4 \cdot 10^6$, with the tails of the mixed terms handled by the prime number theorem). The table of the first coefficients is item (C1) of Appendix A.

Table 2: The multiplicity $m_N = \langle \Psi_N, \mathbf{1} \rangle$ of the trivial character in the plethystic exponents Ψ_N of Lemma 3, for the transitive permutation groups of degree 2 to 5 and three of degree 6 and 7, computed by averaging $\Psi_N(g)$ over the group. A positive entry is the order of the pole of $L(N(s+1), \Psi_N)$ at $s = -1 + 1/N$, a negative one the order of its zero. In every case $m_2 < 0$ (a zero at $s = -\frac{1}{2}$), $m_3 \geq 1$ (a pole at $s = -\frac{2}{3}$, simple exactly for S_n , A_n and A_4), $m_4 < 0$ (a zero at $s = -\frac{3}{4}$) and $m_5 > 0$ (a pole at $s = -\frac{4}{5}$, inside the error term). The regular representations have the largest multiplicities.

G (degree)	$ G $	m_2	m_3	m_4	m_5	m_6
S_2 (2)	2	-2	1	-1	3	-6
C_3 (3)	3	-2	2	-6	16	-40
S_3 (3)	6	-2	1	-2	8	-22
C_4 (4)	4	-3	5	-14	51	-175
V_4 (4)	4	-4	5	-12	51	-180
D_4 (4)	8	-3	3	-6	27	-95
A_4 (4)	12	-2	1	-4	17	-58
S_4 (4)	24	-2	1	-2	10	-34
C_5 (5)	5	-3	8	-30	124	-524
D_5 (5)	10	-3	4	-12	62	-272
F_{20} (5)	20	-2	2	-6	31	-136
A_5 (5)	60	-2	1	-2	12	-51
S_5 (5)	120	-2	1	-2	10	-37
C_6 (6)	6	-4	11	-51	259	-1304
D_6 (6)	12	-4	6	-22	131	-672
A_6 (6)	360	-2	1	-2	10	-40
C_7 (7)	7	-4	16	-84	480	-2808
S_7 (7)	5040	-2	1	-2	10	-37

3.6 Numerical verification

The raw Riesz means $R_m(x)$, $m = 1, 2, 3$, were computed exactly (compensated summation, $A_f(n)$ from $\omega_f(p) = 1 + \chi_{-4}(p)$) at 4096 points equally spaced in $u = \log x \in [\log 10^4, \log 4 \cdot 10^7]$. From R_2 we subtract the exact terms at $s = 1$, $s = 0$, $-\frac{2}{3}$, $-\frac{4}{5}$ (order 3) and $-\frac{6}{7}$ (order 9); the only fitted quantity is a correction to the x^{m+1} coefficient, which comes out at $-1.0 \cdot 10^{-12}$ relative to $D_f(1)/(m+1)$ and is itself a check of $D_f(1)$ to twelve digits. The residual, divided by $x^{5/4}$, has root mean square 0.528. The predicted zero sum (zeros of $L(s, \chi_{-4})$ and of ζ with $\gamma \leq 100$) has root mean square 0.27. Regressing the residual on the prediction gives $\beta = 1.008$ (Figure 2); with a smooth nuisance basis $x^{m-8/9}(\log x)^j$, $j \leq 3$, absorbing the dropped real poles, $\beta = 0.991$ with $R^2 = 0.55$ and a remaining root mean square of 0.291; the two families separately give $\beta_\zeta = 1.006$ and $\beta_L = 1.020$. Controls in which all ordinates are shifted by $\delta \in [-0.6, 0.6]$ give β between -0.94 and 0.68 and negative R^2 . For $m = 3$ the zero terms (rms 0.11) lie below the smooth leftover (rms 0.6); with the nuisance basis $\beta = 0.998$. The remaining rms 0.29 is accounted for by the truncation at $\gamma \leq 100$ (the coefficients decay only like $\gamma^{-5/4}$) and the $O(x^{m-3/4-\varepsilon_0})$ term. We regard this as a confirmation of Theorem 10 with the predicted amplitudes and phases to about 1%; it does *not* resolve individual ordinates closer than $4\pi/8.3 \approx 1.5$. When the contour is pushed to $\Re s = -\frac{7}{8}$ and the second family of zero terms is added, the predicted sum has root mean square 13 (zeros of ζ) and 92 (zeros of L) in the same units and regression coefficient 0.001 against the data: the second family is real as a set of residues but cancels against the line integral, as explained after Theorem 7.

$f = t^2 + 1$, $m = 2$: data minus exact smooth terms (grey) and predicted zero sum (blue)




Figure 2: The residual of the diagonal Riesz mean of order 2 of $t^2 + 1$ after subtracting the exact terms at $s = 1$, $s = 0$ and the real poles $s = -2/3, -4/5, -6/7$, against the sum over the zeros of $L(s, \chi_{-4})$ and of ζ with $\gamma \leq 100$. Regression coefficient $\beta = 1.008$ (0.991 with a smooth nuisance basis); per family $\beta_\zeta = 1.006$, $\beta_L = 1.020$.

4 Ω -results and conditional bounds

Theorem 15. *Let f be an irreducible quadratic and $m \geq 1$, and let $E_m(x) = R_m(x) - \frac{D_f(1)}{m+1}x^{m+1} - x^m(-\frac{\log x}{2C(f)} + c_{f,m}) - c'_{f,m}x^{m-2/3}$. Suppose that for some zero $\rho_1 = \frac{1}{2} + i\gamma_1$ of $\zeta(s)^2 L(s, \chi_D)$ the function G_m has a genuine pole at $\rho_1/2 - 1$ (i.e. $Q_{m,\rho_1} \neq 0$), and that $L(s, \chi_D)$ has no real zero $\beta_0 > \frac{1}{2}$ (true under the Riemann hypothesis, and unconditionally for all but at most one D in any range $|D| \leq X$, by Landau's theorem on exceptional zeros). Then, with no further hypothesis,*

$$E_m(x) = \Omega_\pm(x^{m-3/4}).$$

For $f = t^2 + 1$ the hypothesis holds with $\gamma_1 = 6.0209\dots$, the first zero of $L(s, \chi_{-4})$ ($|Q_{2,\rho_1}|$ is computed in Section 3.5 and is non-zero).

Proof. Suppose $E_m(x) \leq \varepsilon x^{m-3/4}$ for all $x \geq x_0$. The function $F(x) = \varepsilon x^{m-3/4} - E_m(x)$ is then non-negative for $x \geq x_0$, and for $\Re s > 1$

$$\begin{aligned} \int_{x_0}^{\infty} F(x) x^{-s-m-1} dx &= \frac{\varepsilon x_0^{-s-1/4}}{s + \frac{1}{4}} - G_m(s) \\ &\quad + (\text{the Mellin transforms of the subtracted terms}) + (\text{entire}), \end{aligned}$$

where we used $\int_1^{\infty} R_m(x) x^{-s-m-1} dx = G_m(s)$ for $\Re s > 1$ (Mellin inversion of (6)) and the fact that $\int_1^{x_0}$ is entire; the Mellin transforms of x^{m+1} , $x^m \log x$, x^m , $x^{m-2/3}$ are $1/(s-1)$, $1/s^2$, $1/s$, $1/(s+\frac{2}{3})$ up to constants, and they cancel the poles of G_m at $1, 0, -\frac{2}{3}$ exactly by the definition of the coefficients. The right-hand side is therefore meromorphic in $\Re s > -\frac{4}{5}$ (Theorem 4), holomorphic on the real segment $(-\frac{4}{5}, \infty)$ (the remaining real singularities of D_f in this range are the zeros at $-\frac{1}{2}$ and $-\frac{3}{4}$; a real zero $\beta_0 > \frac{1}{2}$ of $L(s, \chi_D)$ would give $L(2w, \chi_D)^{-1}$ a real pole at $\beta_0/2 - 1 > -\frac{3}{4}$, which is why it is excluded — with such a zero the remainder would instead contain a term $\asymp x^{m-1+\beta_0/2}$ of constant sign), and has a pole at $\rho_1/2 - 1$, of real part $-\frac{3}{4}$. By Landau's theorem for integrals with non-negative integrand [36, Theorem 1.7], the abscissa of convergence of the left-hand side is a singularity of the function on the real axis; hence the integral converges for $\Re s > -\frac{4}{5}$, and its value is holomorphic there, contradicting the pole at $\rho_1/2 - 1$. Thus $E_m(x) > \varepsilon x^{m-3/4}$ for arbitrarily large x , i.e. $E_m = \Omega_+(x^{m-3/4})$; applying the argument to $-E_m$ gives Ω_- . $\square$

Remark 16. (i) No positivity of A_f is needed, and no hypothesis beyond the existence of one uncanceled pole on $\Re s = -\frac{3}{4}$; if $\zeta^2 L(\chi_D)$ had a zero with $\beta > \frac{1}{2}$ the same argument gives $\Omega_{\pm}(x^{m-1+\beta/2})$, provided that pole is uncanceled. (ii) Under RH for ζ and $L(s, \chi_D)$, Theorem 10 gives $E_m(x) \ll x^{m-3/4+\varepsilon}$ for $m \geq 2$ provided the zero sum is $O(x^\varepsilon)$ in the normalisation $x^{m-3/4}$, which follows from the simplicity and Gonek–Hejhal hypotheses of the remark above; so under those hypotheses the exponent $m - \frac{3}{4}$ is exact. (iii) For $f = t^2 + t + 41$ the first zero of $L(s, \chi_{-163})$ has $\gamma_1 = 0.2029$, so the Ω -oscillation has period $4\pi/\gamma_1 \approx 62$ in $\log x$: over any computable range the term looks like a constant multiple of $x^{m-3/4}$. The family $\{L(s, \chi_D)\}$ has symplectic symmetry [28], so such a low zero is the class-number-one exception, not the rule.

In words. Independently of any hypothesis, one uncanceled pole on $\Re s = -\frac{3}{4}$ forces the error term to be as large as $x^{m-3/4}$ in both directions infinitely often: the zeros of $L(s, \chi_D)$ are visible in the second moment of S_f unconditionally.

5 The off-diagonal: what part III must prove

This section turns Hypothesis (E) of part I into the analytic continuation of one Dirichlet series and identifies that series for quadratics.

5.1 The analytic reformulation

For d squarefree and roots $s \neq s'$ of f modulo d let $m_{ss'} \in [1, d-1]$ be the representative of $s' - s$. Define, for $\Re s > 1$,

$$O_f(s) = \sum_d W_f(d) \sum_{\substack{s, s' \bmod d \\ s \neq s'}} \left(d^{-s} \zeta\left(s, \frac{m_{ss'}}{d}\right) - \frac{\zeta(s)}{d} \right), \quad (9)$$

$\zeta(s, \alpha)$ the Hurwitz zeta function, so that $d^{-s} \zeta(s, m/d) = \sum_{h \equiv m \pmod{d}} h^{-s}$; each bracket is entire in s , the poles at $s = 1$ cancelling.

Proposition 17. (i) For $\Re s > 1$,

$$\sum_{h \geq 1} (S_f(h) - C(f)^2) h^{-s} = C(f)^2 \zeta(s) (D_f(s) - D_f(1)) + C(f)^2 O_f(s),$$

the left side converging absolutely and the double series (9) converging in the indicated order.

(ii) If O_f continues holomorphically to $\Re s > -\delta$ for some $\delta > 0$ with $O_f(\sigma + it) \ll (1 + |t|)^A$ there, then for some $\delta' > 0$

$$\sum_{h \leq H} \left(1 - \frac{h}{H}\right) (S_f(h) - C^2) = -\frac{1}{2} C(f) \log H + A_f^* + O(H^{-\delta'}), \quad A_f^* = C^2 (\kappa_f^* + O_f(0)),$$

κ_f^* the explicit constant produced by the diagonal. In particular Conjecture (3) holds in Cesàro form and Hypothesis (E) in Cesàro form, $\text{Off}_f^*(H) = O(H)$, holds.

Part III shows unconditionally that $\text{Off}_f^*(H) = c_{\text{off}}(f)H + \mathcal{W}(H) + O(H(\log H)^{1-c})$ with an explicit constant $c_{\text{off}}(f) < 0$ and a remainder $\mathcal{W}(H)$ (the “windows”) of trivial size $H \log H$; under the hypothesis of (ii), $\text{Off}_f^*(H)/H \rightarrow O_f(0)$, so that $O_f(0) = c_{\text{off}}(f)$ exactly when $\mathcal{W}(H) = o(H)$, which the numerics of part III support for $t^2 + 1$.

Proof. (i) By part I, Theorem 2, $S_f(h)/C^2 = \sum_q b(q) c_q^f(h)$ absolutely, and the sum over h in a residue class regroups, for each squarefree d , into pairs of roots modulo d , the subtraction of

the mean per pair being allowed because $\sum_{d|q} \mu(q/d) = 0$ for $q > 1$. With h^{-s} in place of the indicator of $h \leq H$, $\sum_{h \equiv m(d)} h^{-s} = d^{-s} \zeta(s, m/d)$ for $m \in [1, d-1]$ and $d^{-s} \zeta(s)$ for $m \equiv 0$, and the subtracted mean becomes $\zeta(s)/d$. The diagonal pairs give $\sum_d a_f(d) \zeta(s) (d^{-s} - d^{-1}) = \zeta(s) (D_f(s) - D_f(1))$ and the remaining pairs give $O_f(s)$. Absolute convergence of the left side for $\Re s > 1$ follows from $\sum_{h \leq H} S_f(h) = C^2 H + O(\log H)$ (part I, Theorem 2) by partial summation. (ii) By Perron's formula with the weight $1 - h/H$,

$$\sum_{h \leq H} \left(1 - \frac{h}{H}\right) (S_f(h) - C^2) = \frac{C^2}{2\pi i} \int_{(2)} [\zeta(s) (D_f(s) - D_f(1)) + O_f(s)] \frac{H^s}{s(s+1)} ds.$$

Move the contour to $\Re s = -\delta'$, $0 < \delta' < \min(\delta, \frac{1}{2})$. The diagonal factor $\zeta(s) (D_f(s) - D_f(1))$ is holomorphic there except at $s = 0$ (a simple pole of D_f times the kernel's $1/s$; $D_f(s) - D_f(1)$ vanishes at $s = 1$, so there is no pole at 1), where the residue is $-\frac{1}{2C} \log H + \kappa_f^*$ because $\zeta(0) = -\frac{1}{2}$ and $\text{Res}_0 D_f = 1/C$; the off-diagonal factor is holomorphic and contributes the residue $O_f(0)$ at the pole of the kernel. On $\Re s = -\delta'$ the diagonal factor has polynomial growth ($D_f = \zeta_K(w) \times$ an absolutely convergent product for $\Re s > -\frac{1}{2}$, Theorem 4 with $K = 1$), and so has O_f by hypothesis; the kernel makes the integral $O(H^{-\delta'})$. $\square$

In words. Conjecture (3) is the statement that one Dirichlet series, O_f , has no singularity at $s = 0$ and none slightly to its left. A pole of O_f at $s = 0$ would add a term $H \log H$ to the Cesàro sum and break the universality of $-\frac{1}{2}$; this is the danger for Galois groups that are not 2-transitive noted in part I, and (9) is where it would show.

5.2 Quadratics: a Dirichlet series of Salié-type sums

Let $f = at^2 + bt + c$ with discriminant D . For d coprime to $2aD$ the roots of f modulo d correspond to the square roots of $D' = D/a^2$ modulo d , and the differences of distinct roots modulo d are indexed by divisors $d' \mid d$, $d' > 1$, and square roots ν of D' modulo d' (part I, companion note, Proposition A1). Hurwitz's formula, valid for $\Re s < 0$,

$$\zeta(s, \alpha) = \frac{2\Gamma(1-s)}{(2\pi)^{1-s}} \sum_{k \geq 1} \frac{\sin(2\pi k\alpha + \pi s/2)}{k^{1-s}},$$

turns each term $d^{-s} \zeta(s, m/d)$ into a Fourier series in m/d , and hence $O_f(s)$, if it continues to $\Re s < 0$, into

$$O_f(s) = \frac{2\Gamma(1-s)}{(2\pi)^{1-s}} \sum_{k \geq 1} \frac{1}{k^{1-s}} \sum_d \frac{W_f(d)}{d^s} \sum_{\substack{d' \mid d \\ d' > 1}} 2^{k(d)-k(d')} \sum_{\substack{\nu^2 \equiv D' \pmod{d'} \\ (d')}} \sin\left(2\pi k \frac{\overline{(d/d')}}{d'} \nu + \frac{\pi s}{2}\right), \quad (10)$$

$k(\cdot)$ the number of prime factors and $\overline{(d/d')}$ the inverse of d/d' modulo d' . The innermost sum is a Weyl sum of quadratic roots, a Salié-type sum: for prime d' it is $\sum_{\nu^2 \equiv D' \pmod{d'}} e(k\nu/d')$ up to the phase. Dirichlet series in d of such sums, for fixed k and without the weights W_f , are the objects of Hooley [24], Bykovskii, Duke, Friedlander and Iwaniec [14, 15] and Tóth [45]: their continuation is the equidistribution of the roots of $\nu^2 \equiv D'$ modulo d , and its spectral source is the theory of Maass forms through the Kuznetsov formula and its Salié-sum analogue. The identity (10) is formal (it interchanges the sums over d and k); making it rigorous is the content of the following programme.

Proposition 18 (the two estimates). *Hypothesis (E) in Cesàro form holds for the quadratic f if the following two statements hold.*

(E1) (*Small moduli.*) For every $k \geq 1$ and every unit class, the weighted Weyl sums $\sum_{d \leq x} \lambda(d) \sum_{\nu^2 \equiv D'(d)} e(k\bar{u}\nu/d)$, with λ the multiplicative weight $\lambda(p) = p/(p-4)$ on split primes (so that $|W_f(d)| \asymp \lambda(d)/d$), are $\ll x(\log x)^{-1-\eta}$ uniformly in $k \leq x^c$, for some $\eta, c > 0$.

(E2) (*Large moduli.*) $\sum_{h \leq H} \sum_{\substack{e|Q(h) \\ e < Q(h)/H}} \frac{e \lambda(Q(h)/e)}{Q(h)} - H \sum_{d > H} \frac{W_f(d) \omega_f^\circ(d)}{d} = O(1)$, where $Q(h) = a^2 h^2 - D$ and $\omega_f^\circ = \omega_f^2 - \omega_f$.

Proof. This is Propositions A1–A3 of the companion note to part I: the moduli $d \leq H$ contribute a weighted sum of discrepancies of the roots at the points $\{H/d\}$, which (E1) controls through the Erdős–Turán inequality, and the moduli $d > H$ contribute the hyperbola sum of (E2), because a pair of distinct roots modulo $d > H$ has a difference $\leq H$ only if every prime factor of d divides $hQ(h)$ for some $h \leq H$. $\square$

5.3 What Hooley already did, and the route

The decomposition (2) is not new in spirit: Hooley’s 1963 asymptotic for $\sum_{n \leq x} \tau(n^2 + a)$ [24] splits the count, after the hyperbola reflection at $\sqrt{n^2 + a}$, into a Dirichlet-series part $\sum_{k \leq X} \rho(k)/k$ (our diagonal at $s = 1$) and a sum $\sum_k \{2\Psi_k(x) - \Psi_k(Y_k)\}$ of sawtooth values $\Psi_k(y) = \sum_{\nu^2 \equiv -a(k)} \psi((y - \nu)/k)$ at the roots (our Off_f , with Y_k the reflected argument giving our large-moduli part). He notes that his untwisted Weyl-sum bound does not apply to these sums because of the factors depending on k , and proves a twisted bound (his Lemma 5) for $\sum_{k \leq x} \rho(h, k) e(\pm hx/k)$ by building the phase into the parametrisation of the roots by binary quadratic forms and into the resulting incomplete Kloosterman sums. That twisted sum, with the phase $e(kH/d)$ and our weights, is what (E1) needs. The route to (E1) and (E2) is therefore:

- (A) a bound $\sum_{d \leq X, e|d} \rho(k, d) e(kH/d) \ll k^A e^B X^{1-\delta}$ for moduli in a progression, with polynomial dependence on k and e : for $d \asymp C \geq (kH)^{1/2+\varepsilon}$ the phase is smooth at scale C and the theorem of Duke, Friedlander and Iwaniec [15] (smooth weights, moduli $\equiv 0 \pmod q$, either sign of D) applies as it stands; for smaller C one must redo Hooley’s Lemma 5 with the phase, or treat $e(kH \cdot \Im \gamma i)$ as part of the test function of the Poincaré series of Bykovskii [11]; Iwaniec [26] gives the case $e \mid d$ for $D = -1$ with the uniformity needed for a sieve;
- (B) the weights: $W_f(d) = d^{-1} \sum_{e|d} g(e)$ with $g(e) \ll e^{-1+\varepsilon} \tau(e)^A$, so that only $e \leq (\log H)^A$ matter and polynomial dependence on e in (A) suffices; no theorem with genuinely multiplicative weights on composite moduli is needed, which is fortunate because none exists;
- (C) the large moduli: Hooley’s reflection turns them into the same structure with the divisors e of $Q(h)$ in place of d , and the level-of-distribution input for $\#\{h \leq H : e \mid Q(h)\}$ is in [26] and, uniformly in the shift, in the recent work of Grimmelt and Merikoski.

Expanding the sawtooth to length K (Vaaler) and using (A)–(C) gives $\text{Off}_f(H) \ll K^A + (\log H)/K$, i.e. $\text{Off}_f(H) = o(\log H)$: this is Hypothesis (E) in Cesàro form, hence Conjecture (3) in Cesàro form for every quadratic, since S_2 is 2-transitive on the two roots. The sharp form $\text{Off}_f(H) = O(1)$, which an earlier version treated as the stronger target, is false: $\text{Off}_f(H) - \text{Off}_f(H-1)$ is the off-diagonal part of $(S_f(H) - C^2)/C^2$, which is unbounded along h with many small split primes dividing $h^2 - D$ (part I, Remark on the sharp sum, v9); the sharp question is the order of the remainder, which needs decay in the frequency k that the $k^{1/4}$ of [15] does not provide; it is the spectral problem of the next remark.

Remark 19 (a second spectrum). The weighted counts of roots of quadratic congruences summed over moduli have an explicit formula over the Laplace eigenvalues of the modular surface: Bykovskii’s formula, in the form of Soundararajan and Young [43, Prop. 2.2], writes the prime geodesic counting function as $2 \sum_{n \leq X} \sqrt{n^2 - 4} L(1, n^2 - 4)$ with $L(s, \delta) = \frac{\zeta(2s)}{\zeta(s)} \sum_q \rho_q(\delta) q^{-s}$, $\rho_q(\delta)$ the number of roots of $x^2 \equiv \delta$ modulo $4q$, and its explicit formula runs over the Maass eigenvalues; Hejhal [23] treats the Weyl sums for $D > 0$ as integrals of automorphic functions over closed geodesics. In our setting the discriminant is fixed and the modulus varies, the regime of Poincaré series and long horocycles (Bykovskii; Marklof and Welsh [34]); $\text{Off}_f(H)$ is a Poincaré series with a y -dependent sawtooth test function, and its natural expansion is $\sum_j c_j(D) \hat{h}_H(t_j) + (\text{Eisenstein})$, the Eisenstein term reintroducing the zeros of ζ . The dichotomy “diagonal governed by zeros of ζ , off-diagonal by the discrete spectrum” is the one of the additive divisor problem and of Motohashi’s formula for the fourth moment [38]. Whether the discrete spectrum really contributes to Off_f , or decouples as in the non-dihedral cases of Templier and Tsimerman, is testable, and part IV of this series [41] addresses it. Two corrections to the test as we first posed it are worth recording. The observable is not $\text{Off}_f(H)$, which is bounded, but the Cesàro *pieces* $\mathcal{P}_u(Y)$ of part III at the scale $\sqrt{Y}$; and only the *even* Laplace parameters occur, the odd ones being annihilated because the Heegner points of a given discriminant are symmetric about the imaginary axis while an odd Maass form is antisymmetric. With those two corrections the lines are there: part IV derives the expansion for $u = 1$, with amplitudes the Katok–Sarnak periods, and confirms it numerically, the phases being predicted with no free parameter and agreeing to a median of 0.012 radians. (The proof given there has a gap at the passage from the restricted divisor sum to complete modular orbits, found by an external reader on 12 September 2026 and recorded in part IV and in the repository’s errata; the numerical evidence does not depend on it.) The guess recorded in this remark, that Off_f is a Poincaré series with a y -dependent sawtooth test function whose natural expansion is $\sum_j c_j(D) \hat{h}(t_j)$ plus an Eisenstein term, is exactly what that argument carries out.

In words. For quadratics the off-diagonal is a question about square roots modulo composite numbers: do the square roots of D modulo d , weighted by a divisor-like function of d , spread out evenly? Hooley proved this without weights, Duke, Friedlander and Iwaniec much more for prime moduli; the weights and the fractional-part test function are the new features; part III takes the off-diagonal apart and part IV identifies what the remaining pieces are. For Galois groups with three or more orbits on pairs of roots, the analogue of (10) has several independent families of Weyl sums, and a bias between them could produce a pole of O_f at $s = 0$; we do not know whether this happens.

5.4 Numerical evidence for Hypothesis (E), and a first look for the second spectrum

By Proposition 17(i) and the closed formula of part I, $\text{Off}_f(H)$ is exactly computable for $f = t^2 + 1$: $\text{Off}_f(H) = \Sigma_f(H)/C^2 + H D_f(1) - \sum_{n \leq H} A_f(n)$, where $\Sigma_f(H)/C^2 = \sum_{h \leq H} (S_f(h)/C^2 - 1)$ is a sieve over h and $h^2 + 4$ and $D_f(1) = 0.7507908137$ is known to twelve digits; the factor C^2 cancels. We computed it at 4096 points equally spaced in $u = \log H \in [\log 10^3, \log 10^7]$ (`scripts/offdiag-spectrum.ts`, four seconds). Over this range $\text{Off}_f(H)$ has mean -0.013 , standard deviation 0.51, extremes -1.40 and 1.49 , and a fitted drift of -0.0013 per unit of $\log H$, against the trivial bound $\asymp \log H/C$: Hypothesis (E) holds numerically to 10^7 with room to spare, and Conjecture (3) with it (Figure 3). The Hann-windowed spectrum of the detrended $\text{Off}_f(e^u)$ (resolution 0.68 in ω) shows no excess power either at the Laplace parameters t_j of $\text{SL}_2(\mathbb{Z})$ or at the half-ordinates $\gamma/2$ of ζ : in units of the local scatter of the log-spectrum, the mean height at the ten Maass lines in [5, 24] is 0.46 and at the eight ζ lines 0.69, against

0.50 ± 0.66 at random frequencies; the strongest peaks ($\omega = 2.38, 13.16, 5.36, 18.32$) are not at predicted positions. At this range the off-diagonal behaves as bounded broadband noise. This is compatible with the discrete spectrum decoupling, as in the non-dihedral cases of Templier and Tsimerman, and with spectral terms that are simply too small to resolve over 9.2 units of $\log H$; it is not evidence for the second spectrum. Item (C4) of Appendix A extends the range. Part III explains the value of the mean: the Cesàro off-diagonal has a main term $c_{\text{off}}(f)H$ with an explicit constant ($c_{\text{off}}(t^2 + 1) = -0.13439$), and the mean of the sharp $\text{Off}_f(t)$ is predicted to be $c_{\text{off}} + (1 - D_f(1))/2 = -0.00978$, in agreement with the exact data to 10^7 (-0.0098 ± 0.0007); so the boundedness of $\text{Off}_f(H)$ over this range is consistent with the data (though the sharp form $\text{Off}_f(H) = O(1)$ is false in the limit, its increments growing like $\log \log H$ along rare H , see §5), but the off-diagonal is not mean-zero, and its mean is not zero by accident but by a near-cancellation of two computable constants.




Figure 3: The exact off-diagonal of $t^2 + 1$ against $\log H$: bounded, mean -0.01 , no drift. A failure of Hypothesis (E) would appear as a slope of order $1/C(f) \approx 0.73$ per unit of $\log H$.

6 What carries over to $\mathbb{F}_q[u]$

Let $\mathcal{A} = \mathbb{F}_q[u]$, q odd, $f \in \mathcal{A}[t]$ irreducible and separable in t , and define ω_f , P_P , W_f , a_f , D_f as over $\mathbb{Z}$ with $|P| = q^{\deg P}$ in place of p ; put $T = q^{-s}$.

Theorem 20. *Theorem 4 holds verbatim over $\mathcal{A}$: $D_f(s) = \prod_{N \leq K} L(N(s+1), \Psi_N) M_{f,K}(s)$ with the same virtual characters Ψ_N of the Galois group of f over $\mathbb{F}_q(u)$, the L -functions being Artin L -functions over $\mathbb{F}_q(u)$ (rational functions of T^N by Weil), and $M_{f,K}$ an Euler product over the primes of $\mathcal{A}$ converging absolutely for $|T| < q^{1-1/(K+1)}$. The circle $|T| = q$ is a natural boundary; in particular D_f is not a rational function of T when $\deg_t f \geq 2$.*

Proof. The local computation (4) and Lemma 3 are identical with $u = |P|^{-1}$, $v = |P|^{-(s+1)}$; Chebotarev over $\mathbb{F}_q(u)$ supplies infinitely many primes with $\omega_f(P) \geq 1$, and the zeros of F_P have $|T| \rightarrow q$ with arguments $(2j+1)\pi/\deg P$, dense as $\deg P \rightarrow \infty$. $\square$

In words. The function field does not make the generating function rational, as one might hope; the structure and the wall are the same. What it gives is an *unconditional* explicit formula: the analogue of $R_m(x)$ is a coefficient extraction, a contour integral over a circle $|T| = r$, and Weil's theorem places every zero of every $L(T^N, \Psi_N)$ on an explicit circle, so the residue computation of Section 3 goes through with no hypothesis. Writing it out for $f = t^2 - D$ and comparing with the exact data of part I is item (C5) of Appendix A. (The companion note

to part I claimed that the remaining Euler product converges for $|T| < q$; this is wrong, for the same reason as over $\mathbb{Z}$.)

7 Consequences for prime values, conditionally

Everything in this section assumes the Hardy–Littlewood pair conjecture for f with a power-saving error term. Let $\mathbf{1}_t$ be the indicator that $f(t)$ is prime, $A_T = \sum_{t < T} \mathbf{1}_t$, and suppose, as in Montgomery–Soundararajan [35], that $\sum_{t < T} \mathbf{1}_t \mathbf{1}_{t+h} = S_f(h) \sum_{t < T-h} w_t w_{t+h} + O(T^{1/2+\varepsilon})$ uniformly for $h \leq T$, $w_t = 1/\log f(t)$. Then formula (6) of part I expresses $\text{Var}(A_T)$ through $\sum_{h < T} (S_f(h) - C^2) W_T(h)$ with a smooth weight W_T , and inserting the explicit formula gives, provided Conjecture (3) holds with the constant A_f ,

$$\begin{aligned} \frac{\text{Var}(A_T)}{\mathbb{E}(A_T)} &= 1 - \frac{\log T}{\log X} + \frac{1 + 2A_f/C(f)}{\log X} \\ &\quad + \frac{c_1(f) T^{-2/3} + \sum_{\rho} c_{\rho}(f, T) T^{-3/4}}{\log X} + O\left(\frac{T^{-1/2+\varepsilon}}{\log X}\right), \quad X \approx f(T), \end{aligned}$$

with $c_{\rho}(f, T)$ bounded and oscillating in $\log T$. Three remarks. (i) The first three terms are the polynomial Montgomery–Soundararajan law of part I; A_f consists of an explicit diagonal part and $C^2 O_f(0)$ (Proposition 17), the latter identified in part III as $C^2 c_{\text{off}}(f)$ up to the windows. (ii) The zero-induced terms have relative size $T^{-3/4}$, below the $T^{-1/2+\varepsilon}$ that the hypothesis tolerates: nothing about primes and zeros follows, and we do not claim it. (iii) Under GRH the variance side is related to pair correlation of zeros by the Goldston–Montgomery theorem; for $\zeta_K = \zeta L(\chi_D)$ the Selberg-class version [10] applies to each primitive factor, and Murty and Perelli [39] predict that the zeros of the two factors are uncorrelated; but this concerns prime *ideals* of K , for which a Dirichlet series exists, not prime values of f , for which none does.

8 Discussion and open problems

What we learned. (1) The object behind the second moment of prime values of f is not one L -function but the plethystic expansion of $1 + \chi_V v$: a canonical sequence of virtual characters Ψ_N of the Galois group, with $\Psi_1 = V$ and $\Psi_2 = -\text{Sym}^2 V$. (2) The linear case is degenerate: its expansion terminates, which is why the twin-prime explicit formula is unconditional and why the wall at $\Re s = -1$ was never in the way. (3) For non-linear f there is a term $x^{m-2/3}$ larger than the zero terms; a numerical search for the zeros that does not subtract it fails, which is what happened in part I. (4) Near a natural boundary an explicit formula is asymptotic, not convergent, in the position of the contour. (5) With the smooth terms subtracted exactly, the zeros of $L(s, \chi_{-4})$ appear in the second moment of $t^2 + 1$ with the predicted amplitude and phase.

Open problems. (a) Prove (E1) and (E2) for quadratics (part III). (b) The sign and size of $m_N = \langle \Psi_N, \mathbf{1} \rangle$ for general G and $N \geq 4$, hence the full list of real poles $-1 + 1/N$ and their orders; Table 2 has $m_4 < 0$ and $m_5 > 0$ in every case, so the next real pole after $-\frac{2}{3}$ would be at $-\frac{4}{5}$, inside the error term, but we have no proof of either sign in general. For S_n , m_N is $\frac{1}{N} \sum_{j|N} \mu(j) (-1)^{N/j+1} \mathbb{E}[\chi_V(g^j)^{N/j}]$, with moments of fixed-point counts of powers of random permutations. (c) The limiting logarithmic distribution of $E_m(x)/x^{m-3/4}$ (in the sense of Rubinstein–Sarnak) and whether it is biased. (d) Higher moments: the k -point singular series of Montgomery–Soundararajan for polynomial tuples and the virtual characters that appear. (e) The unconditional function-field explicit formula for $t^2 - D$ against the exact data.

(f) Whether Hypothesis (E) can fail for non-2-transitive Galois groups: each orbit O of ordered pairs gives a root sum of the difference resolvent $R_O(y) = \prod_O(y - (s' - s))$, quadratic for quadratics, reducible with quadratic factors for V_4 and D_4 quartics ($x^4 + 1$: $R_O = (y^2 - 2)(y^2 + 2)$), irreducible of degree 4 for the cyclic quartic, where only qualitative equidistribution is known (Hooley [25], Kowalski–Soundararajan [29]); the test is whether $\Sigma_f(H) + \frac{1}{2}C \log H$ converges for $x^4 + 1$, $x^4 - 2$, $x^4 + x^3 + x^2 + x + 1$ and a generic S_4 quartic. (g) By the general theorem of Akbary, Ng and Shahabi [1], the normalised error $E_m(x)/(x^{m-3/4} \log x)$ has a limiting logarithmic distribution under RH and a Gonek–Hejhal-type bound; under linear independence of the ordinates of $\zeta \cdot \zeta_{\mathbb{Q}(\sqrt{D})}$ it is symmetric, so E_m is negative with logarithmic density exactly $\frac{1}{2}$ despite the negative main term of (3): a “no bias” statement worth proving. (h) A general natural-boundary criterion for two-variable Frobenian Euler products $\prod_p W(\text{Frob}_p; p^{-1}, p^{-s})$, extending Kurokawa–Moroz; we have proved only the case at hand. (i) Lowering the Riesz order from $m \geq 2$ to all $m > 0$ by the double Mellin transform technique of Brüdern, Kaczorowski and Perelli [9], and a converse: does $E_m(x) \ll x^{m-3/4+\varepsilon}$ imply the Riemann hypothesis for $\zeta_{\mathbb{Q}(\sqrt{D})}$?

A Computational programme

Every number in this paper comes from `research/paper-II/scripts/riesz-raw.ts` (exact Riesz means; seconds) and `research/paper-II/scripts/explicit-diag.py` (exponents, constants, residues by Cauchy integrals, comparison; about 25 minutes with primes to $4 \cdot 10^6$ and ordinates to 100; residues cached). The following items are specified for execution elsewhere; none is needed for the theorems.

- (C1) Table of $Q_{2,\rho}$, $Q_{3,\rho}$ for the first 8 zeros of $L(s, \chi_{-4})$ and 6 of ζ , with an independent check by direct differentiation of $\log G_m$ (analytic derivatives of the local factors) instead of Cauchy integrals. Input: `data/explicit-diag-terms-*.pkl`. Output: a $\text{\LaTeX}$ table. Minutes.
- (C2) Zero sum to $\gamma \leq 300$ (`-analyse-only` reuses the cache; new poles about 2 hours), to see how the unexplained root mean square 0.29 decreases; a phase-randomised control (same ordinates, random phases) beside the shifted-ordinate control.
- (C3) Other polynomials: $t^2 + t + 41$ ($D = -163$; the zero at 0.2029 treated as a known non-oscillating component), $t^2 - 2$ ($D = 8$), an S_3 cubic ($t^3 - 2$; zeros of $\zeta_{\mathbb{Q}(2^{1/3})}$ from the LMFDB, $\omega_f(p)$ from the factorisation of f modulo p). Requires generalising the driver from χ_{-4} to arbitrary $\omega_f(p)$. A day of programming, hours of computing per polynomial.
- (C4) $\text{Off}_f(H)$ for $t^2 + 1$: done locally to $H = 10^7$ (Section 5.4, four seconds). Extend to 10^9 (memory-bound: two doubles per h ; segment the sieve) to gain 4.6 units of $\log H$ and a resolution of 0.45, and repeat for $t^2 + t + 41$ and $t^2 - 2$; then decide about the second spectrum.
- (C5) The function-field explicit formula for $f = t^2 - D$ over $\mathbb{F}_q[u]$, $q = 3, 5, 7$: L -polynomials, exponents (Table 1), residues on circles, comparison with `research/experiments/ff.ts`. A day.
- (C6) Raw Riesz means to $x = 10^9$ by a segmented sieve of $A_f(n)$, extending the $\log x$ range from 8.3 to 11.5 and separating the ordinates 12.99 and 14.13. Hours, ≥ 16 GB or a segmented rewrite.

Data and code

research/paper-II/scripts/riesz-raw.ts, research/paper-II/scripts/explicit-diag.py, research/paper-II/scripts/gen-wp0.py; results in research/paper-II/data/. Part I's code is in research/experiments/.

AI-assisted research: disclosure

This work was carried out with heavy use of large language models, the Anthropic models Claude Fable 5.1 and Claude Opus 5, used interactively by the author in September 2026 as research assistants. The author posed the questions, set the direction and the scope, and made every decision about what to pursue, claim and submit. The model was used to draft mathematical arguments, to write and run the software behind every number in the paper, to search the literature, and to draft the text; all of this was done in dialogue with the author, who verified the code and the numerical results, reviewed the proofs, and takes full responsibility for the content. The model is not an author. The complete record of the work, including the code, the data, a chronological knowledge base listing the errors made and corrected along the way, and the review passes, is in the public repository <https://github.com/gewure/ulamnd> (directory **research/**), so that the extent and the nature of the assistance can be checked in detail.

References

- [1] Amir Akbary, Nathan Ng, and Majid Shahabi. Limiting distributions of the classical error terms of prime number theory. *Q. J. Math.*, 65(3):743–780, 2014.
- [2] Brandon Alberts. Explicit analytic continuation of Euler products, 2024. arXiv:2406.18190; Essentials of Number Theory 5 (2026), 49–112.
- [3] W. Banks and K. Ford. Sets of integers satisfying Bateman–Horn statistics, 2026. arXiv:2605.01155.
- [4] Gautami Bhowmik, Driss Essouabri, and Ben Lichtin. Meromorphic continuation of multi-variable Euler products. *Forum Math.*, 19:1111–1139, 2007.
- [5] Gautami Bhowmik, Karin Halupczok, Kohji Matsumoto, and Yuta Suzuki. Goldbach representations in arithmetic progressions and zeros of Dirichlet L -functions. *Mathematika*, 65:57–97, 2019. arXiv:1704.06103.
- [6] Gautami Bhowmik and Jan-Christoph Schlage-Puchta. Natural boundaries of Dirichlet series. *Funct. Approx. Comment. Math.*, 37:17–29, 2007.
- [7] Giacomo Bortolussi. Worse than square-root cancellation in Bateman–Horn’s conjecture, 2026. arXiv:2604.02287.
- [8] Tim Browning, Efthymios Sofos, and Joni Teräväinen. Bateman–Horn, polynomial Chowla and the Hasse principle with probability 1, 2022. arXiv:2212.10373, revised 2026.
- [9] Jörg Brüdern, Jerzy Kaczorowski, and Alberto Perelli. Explicit formulae for averages of Goldbach representations. *Trans. Amer. Math. Soc.*, 372:6981–6999, 2019.
- [10] H. M. Bui, J. P. Keating, and D. J. Smith. On the variance of sums of arithmetic functions over primes in short intervals and pair correlation for L -functions in the Selberg class. *J. London Math. Soc. (2)*, 94(1):161–185, 2016. arXiv:1506.03741.

- [11] V. A. Bykovskii. Spectral expansions of certain automorphic functions and their number-theoretic applications. *Zap. Nauchn. Sem. LOMI*, 134:15–33, 1984. English transl.: J. Soviet Math. 36 (1987), 8–21.
- [12] G. Dahlquist. On the analytic continuation of Eulerian products. *Ark. Mat.*, 1:533–554, 1952.
- [13] Marcus du Sautoy and Fritz Grunewald. Zeta functions of groups: zeros and friendly ghosts. *Amer. J. Math.*, 124:1–48, 2002.
- [14] W. Duke, J. B. Friedlander, and H. Iwaniec. Equidistribution of roots of a quadratic congruence to prime moduli. *Annals of Mathematics*, 141:423–441, 1995.
- [15] W. Duke, J. B. Friedlander, and H. Iwaniec. Weyl sums for quadratic roots. *Int. Math. Res. Not. IMRN*, (11):2493–2549, 2012.
- [16] T. Estermann. On certain functions represented by Dirichlet series. *Proc. London Math. Soc. (2)*, 27:435–448, 1928.
- [17] J. B. Friedlander and D. A. Goldston. Some singular series averages and the distribution of Goldbach numbers in short intervals. *Illinois Journal of Mathematics*, 39:158–180, 1995.
- [18] D. A. Goldston and H. L. Montgomery. Pair correlation of zeros and primes in short intervals. In *Analytic Number Theory and Diophantine Problems (Stillwater, OK, 1984)*, volume 70 of *Progress in Mathematics*, pages 183–203. Birkhäuser, 1987.
- [19] D. A. Goldston and A. I. Suriajaya. The error term in the Cesàro mean of the prime pair singular series. *Journal of Number Theory*, 227:144–157, 2021.
- [20] D. A. Goldston and A. I. Suriajaya. A singular series average and the zeros of the Riemann zeta-function. *Acta Arithmetica*, 2021. arXiv:2007.16099.
- [21] D. A. Goldston and Ade Irma Suriajaya. On an average Goldbach representation formula of Fujii. *Nagoya Math. J.*, 250:511–532, 2023. arXiv:2110.14250.
- [22] Chris Hall, Jonathan P. Keating, and Edva Roditty-Gershon. Variance of arithmetic sums and L -functions in $\mathbb{F}_q[t]$, 2019. arXiv:1703.09190; Algebra Number Theory 13 (2019).
- [23] Dennis A. Hejhal. Roots of quadratic congruences and eigenvalues of the non-Euclidean Laplacian. In *The Selberg Trace Formula and Related Topics*, volume 53 of *Contemp. Math.*, pages 277–339. Amer. Math. Soc., 1986.
- [24] C. Hooley. On the number of divisors of quadratic polynomials. *Acta Mathematica*, 110:97–114, 1963.
- [25] C. Hooley. On the distribution of the roots of polynomial congruences. *Mathematika*, 11:39–49, 1964.
- [26] Henryk Iwaniec. Almost-primes represented by quadratic polynomials. *Invent. Math.*, 47:171–188, 1978.
- [27] Henryk Iwaniec and Emmanuel Kowalski. *Analytic Number Theory*, volume 53 of *American Mathematical Society Colloquium Publications*. American Mathematical Society, 2004.
- [28] Nicholas M. Katz and Peter Sarnak. Zeroes of zeta functions and symmetry. *Bull. Amer. Math. Soc. (N.S.)*, 36(1):1–26, 1999.

- [29] Emmanuel Kowalski and Kannan Soundararajan. Equidistribution from the Chinese remainder theorem. *Adv. Math.*, 385:107776, 2021. arXiv:2003.12965.
- [30] Vivian Kuperberg. Sums of singular series with large sets and the tail of the distribution of primes, 2022. arXiv:2210.09775, revised 2023.
- [31] Vivian Kuperberg. Odd moments in the distribution of primes. *Algebra Number Theory*, 19(4):617–666, 2025. arXiv:2109.03767.
- [32] Vivian Kuperberg, Brad Rodgers, and Edva Roditty-Gershon. Sums of singular series and primes in short intervals in algebraic number fields. *Ramanujan J.*, 58:291–317, 2022. arXiv:2001.09513.
- [33] Nobushige Kurokawa. On the meromorphy of Euler products (I). *Proc. London Math. Soc.* (3), 53(1):1–47, 1986.
- [34] Jens Marklof and Matthew Welsh. Fine-scale distribution of roots of quadratic congruences. *Duke Math. J.*, 172:2303–2364, 2023. arXiv:2105.02854.
- [35] H. L. Montgomery and K. Soundararajan. Primes in short intervals. *Communications in Mathematical Physics*, 252:589–617, 2004.
- [36] Hugh L. Montgomery and Robert C. Vaughan. *Multiplicative Number Theory I. Classical Theory*, volume 97 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, 2007.
- [37] B. Z. Moroz. On the analytic continuation of Euler products. *Manuscripta Math.*, 1988. cited after Alberts [2]; bibliographic details to be verified.
- [38] Yoichi Motohashi. An explicit formula for the fourth power mean of the Riemann zeta-function. *Acta Math.*, 170:181–220, 1993.
- [39] M. Ram Murty and A. Perelli. The pair correlation of zeros of functions in the Selberg class. *Int. Math. Res. Not.*, (10):531–545, 1999.
- [40] Jasper Reichardt. The pair singular series of a polynomial: an exact identity, its Dedekind zeta function, and the variance of prime values. Part I; preprint, 2026, <https://github.com/gewure/ulamnd>, 2026.
- [41] Jasper Reichardt. The pair singular series of a polynomial IV: the second spectrum. In preparation, 2026; <https://github.com/gewure/ulamnd>, 2026.
- [42] Efthymios Sofos. The Bateman–Horn conjecture on average for generalized von Mangoldt functions, 2026. arXiv:2606.15698.
- [43] Kannan Soundararajan and Matthew P. Young. The prime geodesic theorem. *J. Reine Angew. Math.*, 676:105–120, 2013.
- [44] E. C. Titchmarsh. *The Theory of the Riemann Zeta-Function*. Oxford University Press, second edition, 1986. Revised by D. R. Heath-Brown.
- [45] Á. Tóth. Roots of quadratic congruences. *International Mathematics Research Notices*, 2000(14):719–739, 2000.
- [46] R. C. Vaughan. On a variance associated with the distribution of primes in arithmetic progressions. *Proceedings of the London Mathematical Society*, 82:533–553, 2001.