

The pair singular series of a polynomial V : the dilated pieces below $H^{1/2}$

Jasper Reichardt*

16 September 2026 — working draft, not for circulation

Abstract

[Draft. The proofs of Theorems 5 and 7 are in [research/explore/PAPER-V-thmM.md](#) and [PAPER-V-smallend-draft.md](#) and are under adversarial reading; nothing here is to be circulated before those readings are recorded.] For a monic irreducible quadratic f without fixed prime divisor, part III decomposed the Cesàro off-diagonal $\text{Off}_f^*(H)$ into pieces indexed by the dilation u and proved that the pieces with $u > H^{1/2+\varepsilon}$ contribute a main term $c_{\text{off}}(f)H$. This paper treats dilations below that threshold. We prove that the pieces with $H^{0.47} < u \leq H^{0.53}$ contribute $O(H^{1-\delta})$, so that $\sum_{u > H^{0.47}} w(u) \mathcal{P}_u(H/u) = c_{\text{off}}(f)H + O(H^{1-\delta})$; the input is a bilinear estimate for Kloosterman fractions of Bourgain and Garaev together with a bound for the additive energy of reciprocals, valid for every modulus, which we deduce from an argument of Cilleruelo and Garaev. For negative discriminants we prove a second result at the other end: the pieces with $u \leq H^{1/3-c}$ contribute $O(H^{1-\delta(c)})$, by running the automorphic kernel method of Grimmelt and Merikoski with the leading coefficient u^2 tracked. What remains is a neighbourhood of $u = H^{1/3}$.

1 Introduction

[Working draft: the proofs are in [research/explore/PAPER-V-thmM.md](#) and [PAPER-V-smallend-draft.md](#) and are under adversarial reading.]

The problem. Let $f = t^2 + bt + c \in \mathbb{Z}[t]$ be monic, irreducible, without fixed prime divisor, of discriminant D . Part III [6] wrote the Cesàro off-diagonal of the pair singular series as a sum of *pieces*,

$$\text{Off}_f^*(H) = P(1) \sum_u \lambda(u) \omega(u) \mathcal{P}_u(H/u),$$

one for each squarefree *dilation* u , the piece $\mathcal{P}_u(Y)$ being the same problem at length $Y = H/u$ with the roots of $x^2 \equiv D$ replaced by those of $u^2 x^2 \equiv D$. Each piece is $O(Y)$, and the weights satisfy $\sum_{u \leq H} |w(u)|/u \asymp \log H$: the trivial bound for the whole sum is $H \log H$, which is exactly the size of the statement to be proved. Every piece must therefore be small, and the difficulty is uniformity in u .

Exponent coordinates. Write $u = H^\alpha$. Each method available for the pieces gives a saving $H^{-s(\alpha)}$ over the trivial bound on an interval of α , and degenerates at the ends of that interval. In these coordinates part III proves $s > 0$ for $\alpha > 2/3$ (averaging the dilations along u), and the present author's extension of that theorem, recorded in the revised part III, gives $s > 0$ for $\alpha > 1/2$ by a dispersion argument with Weil's bound. Below $1/2$ both mechanisms fail: the

*Independent researcher. Correspondence: jasperreichardt@gmail.com.

incomplete Kloosterman sums that dispersion produces have modulus $[d_1, d_2] \asymp Y^2$, and Weil's bound is trivial once the range of u is shorter than the square root of that modulus. This paper supplies two further mechanisms, one on each side of $\alpha = 1/3$.

Results. Theorem 5 treats α near $1/2$: the weight w is decomposed as $1 * \chi_D * \gamma$, and the resulting bilinear forms in u are estimated by Weil's bound when one factor is long (Type I) or short (Type II), and, in the remaining balanced case, by the bilinear bound of Bourgain and Garaev [1] for Kloosterman fractions. That bound needs an estimate for the additive energy of reciprocals; the one in [1] is too weak at the sizes that occur here, and Lemma 2 supplies a stronger one, valid for every modulus, by transferring an argument of Cilleruelo and Garaev [2] for points on modular hyperbolas. Theorem 7 treats small α for negative discriminants: piece u is, after removing the content, a Type I sum for the polynomial $u^2 \ell^2 + |D|$ in the sense of Grimmelt and Merikoski [3], and their automorphic kernel theorem [4] applies at level $u^2 e$ once the two kernel quantities are estimated with the dilation tracked. The diagonal is the number of orbits of the family, $\asymp u$; the off-diagonal, which their general bound would give as $\asymp u^2$, is in fact bounded independently of u (Lemma 6), because the distances inside the family are quantised in steps of $1/(4h)$ whatever the dilation.

What remains. Both mechanisms degenerate at $\alpha = 1/3$, from opposite sides and for unrelated reasons: there the balanced Type II atoms sit at $d^{1/4}$, where the energy input is not strong enough, and the kernel method's diagonal $\asymp u$ meets the trivial bound. The final section explains what the residual band costs. Two points are worth recording. A band of width 2δ in α costs $\delta \cdot H(\log H)^{1-c} \log \log H$, not $\delta H \log H$, because part III's Theorem E applies piece by piece inside it; so a failing method never makes the total worse. And if the savings on both sides vanish only at $\alpha = 1/3$, with implied constants that are powers of $\log H$, then the blocks near the point contribute $O(H)$ in total, which would give $\text{Off}_f^*(H) = O(H)$ — Hypothesis (E) of part I [5], a bounded second-order term in the Cesàro asymptotic. The constant A_f needs strictly more: a saving at the point itself.

2 Notation

We follow part III [6]. For a prime p let $\omega(p)$ be the number of roots of f modulo p , $\lambda(p) = p/(p - 2\omega(p))$, $P_p = 1 - \omega(p)^2/(p - \omega(p))^2$, $P(1) = \prod_p P_p$ and $w(u) = P(1)\lambda(u)\omega(u)$; w is supported on squarefree u and $\sum_{u \leq t} |w(u)| \ll t$. A modulus d' is *admissible* if it is squarefree, all its prime factors are split, and $(d', 2Du) = 1$; $R_{d'}^{(u)} = \{x \bmod d' : u^2 x^2 \equiv D\}$, $\rho_k(d'; a) = \sum_{r \in R_{d'}} e(kar/d')$, and $\langle x \rangle_{d'} \in [1, d']$ is the representative of x . With $B_{d'}^{(Y)}(m) = \sum_{h \leq Y, h \equiv m \pmod{d'}} (Y - h) - Y^2/(2d')$, the piece is

$$\mathcal{P}_u(Y) = \sum_{d' > 1 \text{ admissible}} \frac{\lambda(d')}{d'} \sum_{x \in R_{d'}^{(u)}} B_{d'}^{(Y)}(\langle x \rangle_{d'}), \quad Y = H/u.$$

We write $G_{d'}(a, Y) = \sum_{r \in R_{d'}} B_{d'}^{(Y)}(\langle ar \rangle_{d'})$, $\bar{G}_{d'}(Y)$ for its mean over the units a , $G^\circ = G - \bar{G}$, and $\xi_k(d', N) = (1 - \cos(2\pi kN/d'))/\sin^2(\pi k/d')$ for the coefficients of the finite Fourier expansion of $B^{(N)}$ (part III, Lemma finfourier). Throughout, x^* denotes the inverse of x modulo the modulus at hand, $e_m(z) = e(z/m)$, and $\varepsilon > 0$ is arbitrarily small. Parameters: $\delta = 10^{-3}$, $\eta = \delta/10$, $\delta_3 = \delta$, $\varepsilon' = \eta/7$, $\kappa = \delta/2$, $K_1 = H^{2\eta}$.

3 The energy of reciprocals and bilinear Kloosterman fractions

We write x^* for the inverse of x modulo the modulus at hand and $e_m(z) = e(z/m)$.

Lemma 1 (smooth incomplete Kloosterman sums). *Let $q \geq 1$ be squarefree, $a \in \mathbb{Z}$, $X > 0$ and $T \geq 1$, and let W be smooth, supported in an interval of length at most $4X$, with $\|W\|_\infty \ll 1$ and $\int |W''| \ll T^2/X$. Then*

$$\sum_{(v,q)=1} W(v) e_q(av^*) \ll \tau(q) \left(1 + \frac{X}{q}\right) T^2 q^{1/2} (a, q)^{1/2}.$$

Proof. Split v into classes modulo q and apply Poisson summation in each: the sum equals $\sum_h \widehat{W}(h/q) S(a, h; q)/q$ with $S(a, h; q)$ the Kloosterman sum. For squarefree q , Weil's bound at the primes and twisted multiplicativity give $|S(a, h; q)| \leq \tau(q) (a, h, q)^{1/2} q^{1/2}$, which covers $h = 0$, where $S(a, 0; q) = c_q(a)$ is a Ramanujan sum and $|c_q(a)| \leq (a, q) \leq q^{1/2} (a, q)^{1/2}$. Since $\widehat{W}(\xi) \ll \min(X, T^2/(X\xi^2))$, one has $q^{-1} \sum_h |\widehat{W}(h/q)| \ll T^2(1 + X/q)$. $\square$

Lemma 2 (additive energy of reciprocals, every squarefree modulus). *Let $m \geq 2$ be squarefree, $1 \leq N \leq m$, and $J_4(N) = \#\{x \in [1, N]^4 : (x_i, m) = 1, x_1^* + x_2^* \equiv x_3^* + x_4^* \pmod{m}\}$. Then for every $\varepsilon > 0$,*

$$J_4(N) \ll_\varepsilon m^\varepsilon (N^2 + N^{7/2} m^{-1/2}).$$

Proof. *Step 1: a hyperbola in a square box.* Let $(\lambda, m) = 1$, $K \in \mathbb{Z}$ and $1 \leq M \leq m$; we claim $I := \#\{(x, y) \in (K, K+M]^2 : xy \equiv \lambda \pmod{m}\} \ll_\varepsilon m^\varepsilon (M^{3/2} m^{-1/2} + 1)$. For $M = 1$ this is trivial, so let $M \geq 2$. Writing $x = K + x'$, $y = K + y'$ with $1 \leq x', y' \leq M$, the congruence becomes $x'y' + Kx' + Ky' \equiv b := \lambda - K^2 \pmod{m}$. Put $T = \lfloor (m/M)^{1/2} \rfloor \geq 1$; by Dirichlet's theorem there are $1 \leq t \leq T$ and u_0 with $tK \equiv u_0 \pmod{m}$ and $|u_0| \leq \min(m/T, m/2)$. Multiplying by t ,

$$tx'y' + u_0x' + u_0y' = b_0 + zm, \quad |b_0| < m/2, \quad |z| \leq \frac{TM^2}{m} + \frac{2M}{T} + \frac{1}{2} \leq 5M^{3/2} m^{-1/2} + \frac{1}{2}.$$

For each z this is $(tx' + u_0)(ty' + u_0) = n_z := t(b_0 + zm) + u_0^2$, and $|n_z| \leq 7m^2$. Now $n_z \equiv t^2(x' + K)(y' + K) \equiv t^2\lambda \pmod{m}$, and λ is a unit with $1 \leq t^2 \leq m/M < m$, so $m \nmid n_z$ and in particular $n_z \neq 0$. Each z therefore contributes at most $\tau(|n_z|) \ll_\varepsilon m^\varepsilon$ pairs, and summing over the $\ll M^{3/2} m^{-1/2} + 1$ admissible z gives the claim.

Step 2: from energy to hyperbolas. Let $J_2(\lambda) = \#\{x, y \leq N, (xy, m) = 1 : x^* + y^* \equiv \lambda\}$, so $J_4 = \sum_\lambda J_2(\lambda)^2$ and $\sum_\lambda J_2(\lambda) = N'^2 \leq N^2$ with $N' = \#\{x \leq N : (x, m) = 1\}$. For a unit λ the congruence $x^* + y^* \equiv \lambda$ is equivalent to $(x - \lambda^*)(y - \lambda^*) \equiv \lambda^{*2}$, a unit hyperbola in a square box with equal shifts, so Step 1 gives $J_2(\lambda) \ll m^\varepsilon (N^{3/2} m^{-1/2} + 1)$. If $g = \gcd(\lambda, m) > 1$ then $p \mid g$ forces $x \equiv -y \pmod{p}$, hence $x \equiv -y \pmod{g}$ and $g \leq 2N$; modulo m/g the residue λ is a unit, so Step 1 applied there gives $J_2(\lambda) \ll m^\varepsilon (g^{1/2} N^{3/2} m^{-1/2} + 1)$ when $N \leq m/g$, while $J_2(\lambda) \leq N$ always. Using $\sum_{\gcd(\lambda, m)=g} J_2(\lambda) \leq N(2N/g + 1)$ and summing over the divisors $g \leq 2N$ of m yields the lemma; in the range $N > m/g$ one uses $4N^4/m \leq 4N^{7/2} m^{-1/2}$. $\square$

Lemma 3 (Bourgain–Garaev). *Let $m \geq 2$, $(a, m) = 1$, $1 \leq N_1, N_2 \leq m$, and let α_1, α_2 be coefficients bounded by 1 and supported on units. Then for all $k_1, k_2 \geq 1$,*

$$\left| \sum_{x_1 \leq N_1} \sum_{x_2 \leq N_2} \alpha_1(x_1) \alpha_2(x_2) e_m(ax_1^* x_2^*) \right|^{2k_1 k_2} \leq m N_1^{2k_1 k_2 - 2k_1} N_2^{2k_1 k_2 - 2k_2} J_{2k_1}(N_1) J_{2k_2}(N_2).$$

Proof. This is the inequality established in the proof of Theorem 3 of [1]: two applications of Hölder's inequality followed by Cauchy–Schwarz, using $\sum_\lambda J_k(\lambda) \leq N^k$ and that a is a unit. $\square$

Corollary 4. For squarefree m and $m^{1/3} \leq N_1, N_2 \leq m$, under the hypotheses of Lemma 3, $|\sum \sum \alpha_1 \alpha_2 e_m(ax_1^* x_2^*)| \ll_\varepsilon m^\varepsilon (N_1 N_2)^{15/16}$.

Proof. Take $k_1 = k_2 = 2$ in Lemma 3 and insert Lemma 2: for $N_i \geq m^{1/3}$ the second term of the energy bound dominates, $N_i^{7/2} m^{-1/2} \geq N_i^2$, so the right-hand side is $\ll m^{1+4\varepsilon} (N_1 N_2)^4 (N_1 N_2)^{7/2} m^{-1} = m^{4\varepsilon} (N_1 N_2)^{15/2}$. $\square$

4 The pieces around $u = H^{1/2}$

Theorem 5. Let f be as above. There is $\delta_M > 0$ (any $\delta_M < 4\eta/7$ with the parameters above, so $\delta_M = 5 \cdot 10^{-5}$ is admissible) such that

$$\sum_{H^{0.47} < u \leq H^{0.53}} w(u) \mathcal{P}_u(H/u) \ll_f H^{1-\delta_M}.$$

Consequently, with part III's Theorem on the range $u > H^{1/2+\varepsilon}$ applied at $\varepsilon = 0.03$, $\sum_{u > H^{0.47}} w(u) \mathcal{P}_u(H/u) = c_{\text{off}}(f)H + O(H^{1-\delta_M})$.

[Full proof in research/explore/PAPER-V-thmM.md (with REVISION 1 and REVISION 2 merged); three independent readings, recorded in PROOFS-uniform.md §28, §30, §35, the last confirming the merged text. What follows is the skeleton to be expanded into the paper text.]

Notation for the proof. Put $\delta = 10^{-3}$, $\eta = \delta/10$, $\delta_3 = \delta$, $\varepsilon' = \eta/7$, $\kappa = \delta/2$ and $K_1 = H^{2\eta}$; all implied constants depend on f and on these parameters. For an admissible modulus d' and a unit a let $G_{d'}(a, Y) = \sum_{r \in R_{d'}} B_{d'}^{(Y)}(\langle ar \rangle_{d'})$, let $\bar{G}_{d'}(Y)$ be its mean over the units and $G^\circ = G - \bar{G}$; let $\rho_k(d'; a) = \sum_{r \in R_{d'}} e(kar/d')$, $\epsilon_{d'}(k) = c_{d'}(k)/\varphi(d')$ and $\xi_k(d', N) = (1 - \cos(2\pi kN/d'))/\sin^2(\pi k/d')$, and set $b_k(d') = \min(1/(4k^2), \pi^2(2Y)^2/(4d'^2))$.

Step R1: reduction to a Mellin-separated form. Partition the range of u smoothly into dyadic blocks, with ramps of relative width $H^{-\kappa}$ at the two endpoints; the transition zones contain $\ll H^{0.53-\kappa}$ values of u and contribute $\ll H^{1-\kappa}(\log H)^C$ by $\mathcal{P}_u(Y) \ll Y$. Fix a block, $U \in [H^{0.47}/4, 2H^{0.53}]$, $Y = H/U$, $a = \log U/\log H \in [0.46, 0.54]$. Then:

- (a) the dilation average contributes $\ll U(\log H)^C$, since $\sum_{d'} (\lambda(d')/d') |\bar{G}_{d'}(H/u)| \ll (\log Y)^C$ for every u ;
- (b) the moduli $d' > YH^\eta$ contribute $\ll H^{1-\eta+3\varepsilon'}$ by part III's Step 3, which is valid for every U ;
- (c) the moduli $d' \leq Y^{1-\delta}$ contribute $\ll UY^{1-\delta}$, by $|\lambda(d')G^\circ| \leq |\lambda|\omega(1+\sigma/(8d'\varphi))$ and Shiu's theorem [7];
- (d) on the remaining range the finite Fourier expansion of part III applies at $N(u) = \lfloor H/u \rfloor$, the frequency $k = 0$ cancelling exactly because $R_{d'}$ is symmetric and $0 \notin R_{d'}$; the frequencies $k > K_1$ contribute $\ll H^{1-\eta}$, and replacing $\xi_k(d', N(u))$ by $\xi_k(d', H/u)$ costs $\ll U(\log H)^C$. Note $d' \geq Y^{1-\delta} > 2K_1$, so $(k, d') \leq K_1 < d'$;
- (e) with $F_{k,d'}(v) = \lambda(d')\xi_k(d', H/v)\phi(v/U)\psi(v)/(2d'^2)$ one has, for $T_0 = 1 + kY/d' + H^\kappa$ and $J \in \{0, 1, 2\}$, $\int |\tilde{F}_{k,d'}(it)|(1+|t|)^J dt \ll |\lambda(d')|(Y/(kd'))T_0^J \log H$, by the total-variation bound for $|t| \leq T_0$ and $J+2$ integrations by parts beyond it, using that the ramp's derivatives are supported on a set of relative measure $\ll H^{-\kappa}$; summing, $\sum_{k \leq K_1} \sum_{d'} |\lambda|\omega \int |\tilde{F}_{k,d'}| \ll Y(\log H)^3$.

The block therefore equals $(2\pi)^{-1} \int \sum_{k \leq K_1} \sum_{d'} \tilde{F}_{k,d'}(it) \mathcal{S}_{k,d'}(t) dt$ plus the errors above, where $\mathcal{S}_{k,d'}(t) = \sum_u w(u) u^{-it} \mathbf{1}_{(u,d')=1} [\rho_k(d'; u^*) - \omega(d') \epsilon_{d'}(k)]$, and u^{-it} factors over any factorisation of u . The mean terms contribute $\ll UY^\delta (\log H)^C$.

Step R5: decomposition and covering. Write $w = c_f(1 * \chi_D * \gamma)$ with χ_D the primitive character of the field, γ multiplicative with $\gamma(p) \ll 1/p$, $|\gamma(n)| \leq 20^{\omega(n)}$ and $\sum_n |\gamma(n)| n^{-1/2-\varepsilon} < \infty$ (for $\omega(2) = 1$ put $u = 2u'$ and absorb the Euler factor at 2 into γ). The part with $n_3 > H^{\delta_3}$ is $\ll UY H^{-\delta_3/2+\varepsilon}$. Decompose n_1, n_2, n_3 into smooth dyadic boxes N_1, N_2, N_3 , discarding boxes with $N_1 N_2 N_3 \not\asymp U$, which vanish. For $a \in [0.46, 0.54]$ the boxes split into three cases: either some atom has $N_i \geq Y^{1/2} H^{3\delta}$ (Step R2); or some atom lies in $W = [H^{3\delta}, UY^{-1/2} H^{-3\delta}]$ (Step R3); or both $N_1, N_2 \in [UY^{-1/2} H^{-3\delta}, Y^{1/2} H^{3\delta}]$ (Step R4). Indeed if $N_2 < H^{3\delta}$ then $N_1 \gg UH^{-3\delta-\delta_3} \geq Y^{1/2} H^{3\delta}$, because $U/Y^{1/2} \geq H^{0.19}$ on the block; the covering is exhaustive with margin 0.183.

Step R2 (Type I). Let an atom of length $N \geq Y^{1/2} H^{3\delta}$ carry the weight 1 (or χ_D restricted to a class modulo the conductor), the cofactor m' carrying arbitrary coefficients $\ll \tau_3(m') 20^\omega$. For fixed k, d', r, m', t , with $g = (k, d')$ and $m = d'/g$, the atom sum is $\sum_n W(n) e_m((k/g)r(nm')^*)$ with $W(n) = \phi(n/N) n^{-it}$ after removing $(n, g) = 1$ by Möbius, so Lemma 1 with $T = 1 + |t|$ applies and gives $\ll H^\varepsilon (1 + |t|)^2 (1 + N/m) m'^{1/2}$. Summing over $m' \ll U/N$, the roots, the moduli against $\int |\tilde{F}|(1 + |t|)^2$ and $k \leq K_1$, and splitting the two branches of $T_0^2 \ll 1 + (kY/d')^2 + H^{2\kappa}$, the contribution is $\ll UY (H^{-3\delta+4\eta+1.5(1-a)\delta} + H^{-3\delta+\eta/2+2\kappa}) + UY^{1/2+O(\delta)} \ll UY H^{-1.79\delta}$.

Step R3 (Type II). Let $u = u_1 u_2$ with $|a(u_1)|, |b(u_2)| \ll H^\varepsilon$ and $U_2 \in W$. Apply Cauchy–Schwarz over (d', u_1) with weight $|\beta(d')|$, $\beta = \tilde{F}_{k,d'}(it)$, replacing the u_1 -range by a smooth majorant. The diagonal gives $S_1 \ll H^{2\varepsilon} U_1 \sum_{d'} |\beta|$. In S_2 the terms carrying $\epsilon_{d'}$ are negligible, and the main u_1 -sums are $\sum V(u_1/U_1) e_{d'}(\nu u_1^*)$ with $\nu \equiv k(ru_2^* - r'u_2^*)$; Lemma 1 bounds each by $\tau(d')(1 + U_1/d') d'^{1/2} (\nu, d')^{1/2}$, and trivially by $3U_1$. Writing $(\nu, d') = (k, d')e$ with $e = \prod_{p|d', p \nmid k, ru_2^* \equiv r'u_2^* \pmod{p}} p$, for fixed (d', r, r', u_2) there are at most $4U_2/e$ values of u_2' with that e when $e \leq 2U_2$, and at most one when $e > 2U_2$ — the latter case containing the diagonal and all degenerate pairs, since $d'/(k, d') > 2U_2$. Hence $S_2 \ll H^{O(\varepsilon+\eta)} \sum_{d'} |\beta| [U_2^2 Y^{1/2} + U_1 U_2^2 Y^{-1/2} + U_1 U_2]$, the middle term alone carrying the loss $Y^{\delta/2}$, and therefore the contribution is $\ll UY H^{-1.45\delta}$.

Step R4 (the residual). Here both atoms lie in $[UY^{-1/2} H^{-3\delta}, Y^{1/2} H^{3\delta}]$. For fixed k, d', r, n_3, t , with $g = (k, d')$ and $m = d'/g$ squarefree, $m \geq Y^{1-\delta} H^{-2\eta}$, the numerator $(k/g) r n_3^*$ is a unit modulo m and the double sum is of the shape treated in Corollary 4, with the coprimality conditions and the dyadic cut-offs absorbed into the coefficients. On the block, $\log N_i / \log m \geq 0.3462 > 1/3$ and $2N_i \leq m$, so Corollary 4 applies and gives $\ll m^\varepsilon (N_1 N_2)^{15/16}$. Since $N_1 N_2 \asymp U/n_3$, summing over n_3 against $|\gamma|$, over the roots, over the moduli against $\int |\tilde{F}|$ and over $k \leq K_1$ gives $\ll UY H^{-0.028}$.

Conclusion. Per block the errors are $H^{1-4\eta/7}$ from R1(b), $H^{1-\eta}$ from R1(d), $UY^{1-\delta}$ from R1(c), UY^δ from the mean terms, $H^{1-\delta_3/2+\varepsilon}$ from the n_3 -tail, $H^{1-\kappa} (\log H)^C$ from the ramps, and the three savings above. With $O(\log H)$ blocks the binding constraint is R1(b), and Theorem 5 follows for every $\delta_M < 4\eta/7$.

5 The small dilations

Lemma 6 (the off-diagonal at one level does not grow with the dilation). *Let $h \geq 1$, u squarefree with $(u, 2h) = 1$, $a = u^2$, and let F_u be the set of positive definite forms $g = \begin{pmatrix} A & B \\ B & C \end{pmatrix}$ with $AC - B^2 = ah$, $a \mid B$, $a \mid C$. For $Z \geq 1$,*

$$\sum_{g_2 \in \Gamma_0(a) \backslash F_u} \sum_{\substack{g_1 \in F_u, g_1 \neq g_2 \\ u(g_1, g_2) \leq Z}} (1 + u(g_1, g_2))^{-1/2} \ll_{\varepsilon} (hZu)^{\varepsilon} h^{O(1)} (1 + Z^{1/2}),$$

uniformly in u , where $u(\cdot, \cdot)$ is the point-pair invariant.

[Proof in PAPER-V-smallend-draft.md (undilation to $\Gamma_0^0(u)$, lift to level one, a local count of common orthogonal lines, and a sum over the quantised distances); under reading.]

Theorem 7. *Let f be as above with $D < 0$. For every $c > 0$ there is $\delta_S(c) > 0$ with $\sum_{u \leq H^{1/3-c}} w(u) \mathcal{P}_u(H/u) \ll_f H^{1-\delta_S(c)}$.*

[Proof in research/explore/PAPER-V-smallend-draft.md (S1, S1', S2', S3–S5, O1, O2, O4–O6'', Lemma 6), resting on [4] Theorem 8.1 and [3] Lemma 3.1. Under reading.]

Remark 8 (on the implied constant imported from the kernel theorem). Theorem 8.1 of [4] is stated with an unspecified factor $\delta^{-O(1)}$, where δ is the smoothness parameter of the input function. Following its proof through §7.1–§7.4 — the unskewing with $R_1 = X$, $R_2 = 1$ is loss-free; F of right type 0 forces $\ell_2 = 0$, so that the second K -type sum is absent and the α_2 -side majorant may be taken at $L = 1$; and $\delta_1 = \delta$ because $X \geq Y$, which here reads $Lu \geq |D|^{1/2}$ and holds on every non-trivial block — one may take δ^{-6} . We do not optimise this and use only that the exponent is absolute: fixing $C = 6$ and choosing the parameters *after* C , namely $\delta = \eta = \eta' = \frac{1}{9} \cdot \frac{3}{2}c(1 - 2\theta)$, all the constraints hold and $\delta_S(c) = \frac{2}{3}\delta \approx c/12$.

6 What remains: the band at $u = H^{1/3}$

Theorems 5 and 7 leave a neighbourhood of $\alpha = 1/3$ uncovered, and the two mechanisms fail there for unrelated reasons. On the upper side the balanced Type II atoms sit at $d^{1/4}$, where the available energy input for reciprocals is exactly at its threshold: what would be needed is a bound for the six-fold energy $J_6(N)$ below N^4 at $N \asymp m^{1/4}$, which is not known even for prime moduli. On the lower side the kernel method's diagonal, the number of orbits of the family, is $\asymp u$, and at $u \asymp Y^{1/2}$ it meets the trivial bound.

Three remarks fix what such a band costs, and what closing it would buy.

Remark 9 (a band is cheap, and a failing method is free). Inside any range of α one still has part III's Theorem E piece by piece, $\mathcal{P}_u(H/u) \ll (H/u)(\log H)^{-c} \log \log H$ uniformly in $u \leq H^{1-\varepsilon}$. Hence a band $[H^{\alpha_0-\delta}, H^{\alpha_0+\delta}]$ on which nothing better is available costs $\ll_f \delta \cdot H(\log H)^{1-c} \log \log H$, not $\delta H \log H$: a mechanism that degenerates can never make the total worse than what is already proved, and any band whose width tends to 0 improves Theorem E by that factor.

Remark 10 (a point gap costs $O(H)$). Suppose the savings on the two sides vanish only at $\alpha = 1/3$, say like $H^{-c|\alpha-1/3|}$, with implied constants that are powers of $\log H$. Summing over dyadic blocks, $\sum_k H e^{-c|k \log 2 - \frac{1}{3} \log H|} = H(1 + 2/(e^{c \log 2} - 1)) = O(H/c)$: the blocks near the point contribute $O(H)$ in total. Since the main term $c_{\text{off}}(f)H$ comes from $u > H^{1/2+\varepsilon}$ and the remaining blocks are error terms, this would give $\text{Off}_f^*(H) = O(H)$, which is Hypothesis (E) of part I [5] and yields $\sum_{h \leq H} (1 - h/H)(S_f(h) - C(f)^2) = -\frac{1}{2}C(f) \log H + O_f(1)$: a bounded second-order term.

Remark 11 (what A_f needs). The constant of Conjecture 1 of part I requires $\text{Off}_f^*(H) = c_{\text{off}}(f)H + o(H)$, hence a saving factor $\gg \log H$ uniformly on $|\alpha - 1/3| \ll \log \log H / \log H$; a fixed positive power saving at $\alpha = 1/3$ itself suffices. A linearly vanishing saving, however small the rate, gives $O(H)$ but never $o(H)$.

Remark 12 (what is actually binding). An audit of every implied constant in the chain (research/explore/PROOFS-uniform.md §36) shows that no divisor-size loss constrains the argument: the divisor factors all occur under sums and average to powers of $\log H$, the kernel theorem's $\delta^{-O(1)}$ is δ^{-20} , its $(AD)^{o(1)}$ is $(\log AD)^{O(1)}$, and its $q^{o(1)}$ is 1 in the present application. What binds is the constant in Bourgain and Garaev's Hölder step, which grows as $k_1 \asymp 2/(9|\alpha - 1/3|) \rightarrow \infty$: the band on which the estimate is effective has width $\asymp (\log \log H / \log H)^{1/3}$. With the in-band fallback on part III's Theorem E this gives, once the map is written out, $\text{Off}_f^*(H) \ll H(\log H)^{0.63+o(1)}$ for $D < 0$, while everything outside the band sums to $O(H)$. The step from there to $\text{Off}_f^*(H) = O(H)$ therefore needs one thing only: an estimate that survives at the exceptional sizes $N \asymp m^{1/4}$ — an energy bound J_6 below N^4 for composite moduli, or a different tool there. A level-averaged spectral bound for the twist $e(c\bar{r}d/x)$ with $x \asymp Y^{1/2}$ would serve the same purpose; neither is in the literature.

References

- [1] Jean Bourgain and Moubariz Z. Garaev. Kloosterman sums in residue rings. *Acta Arith.*, 164(1):43–64, 2014.
- [2] Javier Cilleruelo and Moubariz Z. Garaev. Concentration of points on two and three dimensional modular hyperbolas and applications. *Geom. Funct. Anal.*, 21:892–904, 2011.
- [3] Lasse Grimmelt and Jori Merikoski. On the greatest prime factor and uniform equidistribution of quadratic polynomials, 2025. arXiv:2505.00493.
- [4] Lasse Grimmelt and Jori Merikoski. Weighted averages of $\text{SL}_2(\mathbb{R})$ automorphic kernel, Part I: non-oscillatory functions, 2025. arXiv:2505.00489.
- [5] Jasper Reichardt. The pair singular series of a polynomial I, 2026. Preprint.
- [6] Jasper Reichardt. The pair singular series of a polynomial III: the off-diagonal in Cesàro form, 2026. Preprint.
- [7] P. Shiu. A Brun–Titchmarsh theorem for multiplicative functions. *J. Reine Angew. Math.*, 313:161–170, 1980.