

Dilated pieces, Hecke correspondences and the second spectrum: the level-one lines of the pieces $\mathcal{P}_u$

Jasper Reichardt*

Working draft of 15 September 2026 — not for circulation. Every displayed constant was tested numerically before it was written down; the proofs marked “sketch” are not complete.

Abstract

Part IV of this series [5] showed that the pieces $\mathcal{P}_u(Y)$ of the off-diagonal of the pair singular series of an irreducible quadratic f oscillate at the spectral parameters of the even Maass cusp forms, and that the piece with dilation u lives at level u^2 . It left open “the size, in u , of the restricted geometric factor”. This note computes that factor for the level-one lines and for the newforms of level u . The piece with dilation u is the level-one seed of the smooth-window theorem, at length uY , summed over the $\Gamma_0(u^2)$ -invariant family $\mathcal{W}_u$ of forms of discriminant $4u^2D$ with $u^2 \mid a$ and $2u^2 \mid b$. The coefficient of the line t_j of a level-one form u_j in piece u is therefore a projection onto the oldform space $\{u_j(z), u_j(uz), u_j(u^2z)\}$, and for an odd prime u it equals the coefficient in piece 1 times an explicit rational function of $\lambda_j(u)$, $\sqrt{u}$ and $\chi_D(u)$: the Gram matrix of the oldforms (Iwaniec–Luo–Sarnak) and two local period identities for the Hecke neighbours of Heegner points (Gross–Kohnen–Zagier). The ratio is $\lambda_j(u)/u + O(u^{-3/2})$ for $u \nmid D$ and $u^{-1/2}(1 + O(\lambda_j(u)/\sqrt{u}))$ for $u \mid D$, so the main terms of the level-one lines are uniformly summable over u . The same formula with a newform v of level u gives its line in piece u , and predicts that it vanishes when u is inert in $\mathbb{Q}(\sqrt{D})$. Both predictions carry no free parameter; they were tested on 21 pairs (u, D) for the first level-one line and on seven discriminants for the first newform of level 3, and hold to a few percent in amplitude and a few hundredths of a radian in phase. The error term uniform in u , the open core of the conjecture of part III, is not touched.

1 Introduction

The object. Notation as in part IV [5]. For a discriminant $D < 0$, a squarefree $u \geq 1$ and $Q_u(h) = u^2h^2 - D$, the unrestricted piece with dilation u and smooth window $w \in C_c^\infty((0, \infty))$ is

$$S_u^w(Y) = \sum_{h \geq 1} w\left(\frac{h}{Y}\right) (\sigma_{-1}(Q_u(h)) - \mathbb{E}_u), \quad \sigma_{-1}(n) = \sum_{d \mid n} \frac{1}{d}, \quad \mathbb{E}_u = \sum_d \frac{\rho_u(d)}{d^2},$$

$\rho_u(d)$ the number of roots of $u^2x^2 \equiv D \pmod{d}$. For $u = 1$ Theorem SMOOTH of part IV gives

$$\sqrt{Y} S_1^w(Y) = |D|^{-1/4} \sum_j \text{Per}_D(u_j) \tilde{L}_j[A_j(Y) + \overline{A_j(Y)}] + \mathcal{E}^w(Y) + O(Y^{-2+\varepsilon}), \quad A_j(Y) = \Gamma(-it_j) W_c\left(\frac{3}{2} + it_j\right) \left(\frac{\pi}{2} - \arg \lambda_j(u)\right) \quad (1)$$

the sum over an orthonormal basis of even Maass cusp forms of $\text{SL}_2(\mathbb{Z})$, with $\text{Per}_D(u_j) = \sum_Q u_j(z_Q)/|\Gamma_Q|$ over all classes of forms of discriminant $4D$, $\tilde{L}_j = \sum_k a_j(k)k^{-3/2}$, and W_c the

*Independent researcher. Correspondence: jasperreichardt@gmail.com. This work was carried out with heavy use of a large language model; see the disclosure at the end.

Mellin factor of the window. Part IV, Section 7, showed that piece u lives at level u^2 and named the one missing ingredient: how the coefficients depend on u . Its footnote on $u = 3, 5$ recorded that the level-one lines are weak there and that the lines of $\Gamma_0(9)$ take over.

Results. Write $\lambda_j(u)$ for the Hecke eigenvalue ($T_u u_j = \lambda_j(u) u_j$, $a_j(1) = 1$) and $\chi = \chi_D(u)$ for the Kronecker symbol, so $\chi = 0$ iff $u \mid D$.

Theorem 1 (the spectral formula for piece u). *Let $\mathcal{W}_u = \{[a, b, c] \text{ of discriminant } 4u^2 D : u^2 \mid a, 2u^2 \mid b\}$ (all contents). Then $\mathcal{W}_u$ is $\Gamma_0(u^2)$ -invariant and, for every $\varepsilon > 0$,*

$$\sqrt{Y} S_u^w(Y) = \sqrt{u} |D|^{-1/4} \sum_v \text{Per}_{\mathcal{W}_u}(v) \tilde{L}_v[A_v(uY) + \overline{A_v(uY)}] + \mathcal{E}_u^w(Y) + O(Y^{-2+\varepsilon}),$$

the sum over an orthonormal basis $\{v\}$ of the even Maass cusp forms of $\Gamma_0(u^2)$, with $\text{Per}_{\mathcal{W}_u}(v) = \sum_{Q \in \Gamma_0(u^2) \backslash \mathcal{W}_u} v(z_Q) / |\Gamma_0(u^2)_{z_Q}|$, $\tilde{L}_v$ formed with the Fourier coefficients of v at the cusp ∞ , and $\mathcal{E}_u^w$ the continuous spectrum.

Status: proved (Section 2): it is Theorem SMOOTH of part IV at level u^2 applied to an explicit identity between seeds; every constant checked numerically at $u = 2, 3, 5$.

Theorem 2 (the level-one lines). *Let u be an odd prime and u_j an even Hecke–Maass form of level one. The coefficient of the line t_j in $\sqrt{Y} S_u^w(Y)$ equals the coefficient in $\sqrt{Y} S_1^w(Y)$ multiplied by u^{-it_j} and by*

$$r_j(u; D) = \sqrt{u} P^\top (u(u+1) G)^{-1} L, \quad G = \begin{pmatrix} 1 & g & h \\ g & 1 & g \\ h & g & 1 \end{pmatrix}, \quad g = \frac{\lambda_j(u) \sqrt{u}}{u+1}, \quad h = \frac{\lambda_j(u)^2 - 1 - 1/u}{u+1}, \quad (2)$$

$$P = (p_0, p_1, p_0), \quad p_1 = \begin{cases} 2u & u \mid D, \\ u - \chi & u \nmid D, \end{cases} \quad p_0 = \lambda_j(u) \sqrt{u} + \begin{cases} u - 1 & u \mid D, \\ -(1 + \chi) & u \nmid D, \end{cases} \quad L = (1, u^{-1}, u^{-2}).$$

In particular $r_j(u; D) = \lambda_j(u)/u + O(u^{-3/2})$ for $u \nmid D$ and $r_j(u; D) = u^{-1/2}(1 + O(\lambda_j(u)u^{-1/2}))$ for $u \mid D$, the implied constants absolute.

Status: conditional on Lemmas 6, 7 and 8, each of which is stated with the values the numerics fix and proved to the extent marked. The values of (2) agree with the direct spectral computation to four digits and with the data to the noise (Table 1).

Theorem 3 (the newform lines). *Let u be an odd prime and v an even Hecke–Maass newform of level u with Atkin–Lehner eigenvalue ε ($a_v(u) = -\varepsilon u^{-1/2}$). The coefficient of its line in $\sqrt{Y} S_u^w(Y)$ is*

$$2\sqrt{u} |D|^{-1/4} \left[\sum_{k=1}^2 \text{Per}_{\mathcal{W}_u}(v_k) \tilde{L}(v_k) \right] \Gamma(-it_v) W_c\left(\frac{3}{2} + it_v\right) u^{-it_v},$$

$\{v_1, v_2\}$ the orthonormalisation of $\{v(z), v(uz)\}$ on $\Gamma_0(u^2) \backslash \mathbb{H}$, whose Gram matrix has off-diagonal entry $a_v(u) \sqrt{u}/u = -\varepsilon/u$ relative to $\langle v, v \rangle$. If u is inert in $\mathbb{Q}(\sqrt{D})$ then $\text{Per}_{\mathcal{W}_u}(v) = \text{Per}_{\mathcal{W}_u}(v(u \cdot)) = 0$ and the line is absent.

Status: the formula is Theorem 1 restricted to the two-dimensional oldspace; the Gram entry and the vanishing for inert u are observed numerically (to 10^{-17}) and not yet proved. Tested on seven discriminants (Table 2).

Corollary 4 (uniform summability of the level-one main terms). *For a fixed level-one form u_j and any bounded weights $w(u)$, $\sum_{u \leq U} |w(u)| (H/u)^{1/2} |r_j(u; D)| \ll H^{1/2}$ uniformly in U , the sum over odd primes u converging absolutely like $\sum_u |\lambda_j(u)| u^{-3/2}$.*

Remark 5 (what this does and does not settle). Part III's conjecture needs the pieces with $u \leq H^{2/3}$ to sum to $o(H \log H)$ with their *errors*. The corollary says the level-one *main* terms do so with room to spare; the newform lines of level u are new at every u (Theorem 3 gives their size for each u but not a bound uniform in the newform), the cycloidal-group forms of level u^2 [6] are not covered at all, and the error term of Remark GAPS of part IV is untouched. What has changed is that the main-term side of the u -problem is Hecke theory on Heegner divisors, and it decays.

Method and checks. Section 2 proves Theorem 1. Section 3 states the three lemmas behind Theorem 2 with what is proved of each. Section 4 reports the tests: the ratio (2) against a direct computation of the oldform projection with the Maass forms' values (no closed form used), and both against the pieces computed to $Y = 10^7$; then the newform of level 3; then the case $u = 2$ and the composite $u = 6$. Section 5 lists what is open. Scripts and data are in the repository (`research/explore/`, `hecke-oldform-predict.py`, `hecke-newform-u3.py`, `hecke-ratio-sharp.py`).

2 The seed identity and the family $\mathcal{W}_u$

The pairs (d, x) with $u^2 x^2 \equiv D \pmod{d}$ are the forms $Q = [d, 2ux, (u^2 x^2 - D)/d]$ of discriminant $4D$ with $2u \mid b$; write $\mathcal{H}^{(u)}$ for this sub-family of the set $\mathcal{H}$ of all forms of discriminant $4D$, and $z_Q = (-ux + i\sqrt{|D|})/d$. Poisson summation in h , exactly as in the proof of Theorem SMOOTH, writes $S_u^w(Y)$ as $\sum_{(d,x)} \frac{Y}{d^2} \sum_{k \neq 0} \hat{w}(kY/d) e(kx/d)$, and in the coordinates of $z = z_Q$ ($y = \sqrt{|D|}/d$, $x/d = -\Re z/u$) the summand is

$$\frac{Yy^2}{|D|} \sum_{k \neq 0} \hat{w}\left(\frac{kYy}{\sqrt{|D|}}\right) e\left(-\frac{k\Re z}{u}\right) = u \Psi_{uY}^w(z/u),$$

$\Psi_{Y'}^w(\tau) = \frac{Y'(\Im \tau)^2}{|D|} \sum_{k \neq 0} \hat{w}(kY'\Im \tau / \sqrt{|D|}) e(-k\Re \tau)$ the level-one seed of part IV at length Y' . The point $\tau = z_Q/u = (-2u^2x + i\sqrt{4u^2|D|})/(2u^2d)$ is the Heegner point of the form $Q' = [u^2d, 2u^2x, (u^2x^2 - D)/d]$ of discriminant $4u^2D$, and $Q \mapsto Q'$ is a bijection of $\mathcal{H}^{(u)}$ onto $\mathcal{W}_u$. For $\gamma \in \Gamma_0(u^2)$ the middle coefficient transforms as $b' = 2a\alpha\beta + b(\alpha\delta + \beta\gamma) + 2c\gamma\delta \equiv b\alpha\delta \equiv b \pmod{2u^2}$, since $u^2 \mid a$, $u^2 \mid \gamma$ and $\alpha\delta \equiv 1 \pmod{u^2}$; the condition $u^2 \mid a$ is $\Gamma_0(u^2)$ -stable by Lemma SUBFAMILY of part IV. So $\mathcal{W}_u$ is a finite union of $\Gamma_0(u^2)$ -orbits, the translation $\tau \mapsto \tau + 1$ is $x \mapsto x - d$, and $S_u^w(Y) = u \sum_{\tau \in \mathcal{W}_u/\mathbb{Z}} \Psi_{uY}^w(\tau)$. The proof of Theorem SMOOTH applies verbatim on $\Gamma_0(u^2) \backslash \mathbb{H}$ (its cusp at ∞ has width one, so the pairing with the Fourier expansions at ∞ is unchanged), with Y replaced by uY ; the factor u and $(uY)^{-1/2}$ combine to $\sqrt{u} Y^{-1/2}$, and $(\pi\sqrt{|D|}/(uY))^{it_v}$ carries the phase $-t_v \log u$. This proves Theorem 1. For $f_m(z) = u_j(u^m z)$ the coefficient at index $u^m n$ is $u^{m/2} a_j(n)$, so $\tilde{L}_{f_m} = u^{-m} \tilde{L}_j$, which is the vector L of Theorem 2.

3 The three lemmas

Throughout u is an odd prime and $\text{Per}_D(f) = \sum_{Q \in \mathcal{H}/\text{SL}_2(\mathbb{Z})} f(z_Q)/|\Gamma_Q|$.

Lemma 6 (multiplicity). *For every $\text{SL}_2(\mathbb{Z})$ -class C of forms of discriminant $4D$, $\sum_{Q' \in \Gamma_0(u^2) \backslash \mathcal{W}_u, uz_{Q'} \in C} |\Gamma_0(u^2)_{z_{Q'}}|^{-1} = m_u(D) |\Gamma_C|^{-1}$ with $m_u(D) = 2u$ if $u \mid D$ and $m_u(D) = u - \chi_D(u)$ if $u \nmid D$. Equivalently $\text{Per}_{\mathcal{W}_u}(u_j(u \cdot)) = m_u(D) \text{Per}_D(u_j)$.*

Sketch. $u_j(u\tau_{Q'}) = u_j(z_Q)$, so the identity is a count of the $\Gamma_0(u^2)$ -orbits of $\mathcal{W}_u$ above C , weighted by stabilisers. Transporting by $\tau = z/u$, these are the orbits of $\Gamma^0(u^2) = \{u^2 \mid \beta\}$ on

the forms $\gamma Q_0 \in C$ with $2u \mid b(\gamma Q_0)$. The condition depends on γ modulo u and singles out, in $\mathbb{P}^1(\mathbb{F}_u)$, the points at which a binary quadratic attached to Q_0 does not vanish; there are $u + 1 - (1 + \chi) = u - \chi$ of them when $u \nmid D$, and the lifts to $\mathbb{P}^1(\mathbb{Z}/u^2)$ account for the rest of the count. When $u \mid D$ the quadratic has a double root and the forms with $u \mid a$ contribute a second family of u orbits. [Complete the coset count; the identification of the excluded points with the roots of $Q_0 \bmod u$ is the step to write out. Numerically $m = 6.000, 10.00$ ($u = 3, 5$, $u \mid D$), $2.001, 4.000$ (split), $3.999, 6.000$ (inert).] $\square$

Lemma 7 (the period of the dilated family). $\text{Per}_{\mathcal{W}_u}(u_j) = \text{Per}_{\mathcal{W}_u}(u_j(u^2 \cdot)) = (\lambda_j(u)\sqrt{u} + c_u(D))\text{Per}_D(u_j)$ with $c_u(D) = u - 1$ if $u \mid D$ and $c_u(D) = -(1 + \chi_D(u))$ if $u \nmid D$.

Sketch. For $Q = [a, b, c]$ with $u \nmid a$ the $u + 1$ Hecke neighbours of z_Q are uz_Q , the point of $[a, ub, u^2c]$, and $(z_Q + k)/u$, the points of $[u^2a, u(b - 2ak), ak^2 - bk + c]$, $k \bmod u$, all of discriminant $4u^2D$, and $\sum_{\text{neighbours } \nu} u_j(\nu) = \lambda_j(u)\sqrt{u}u_j(z_Q)$. Exactly one neighbour lies in $\mathcal{W}_u$: $(z_Q + k)/u$ with $k \equiv (b/2)a^{-1} \pmod{u}$, since $2u^2 \mid u(b - 2ak)$ iff $b/2 \equiv ak \pmod{u}$. Among the others, when $u \nmid D$, exactly $1 + \chi$ have content divisible by u and are Heegner points of discriminant $4D$ again — the classical description of T_u on CM points [1]: the classes multiplied by the prime ideals above u — and summed over all classes of $\mathcal{H}$ these contribute $(1 + \chi)\text{Per}_D(u_j)$. The remaining neighbours of discriminant $4u^2D$ are $\Gamma_0(u^2)$ -equivalent to the selected neighbours of the other $m_u(D)$ representatives of the same class (Lemma 6); summing the Hecke relation over the classes and subtracting the same-discriminant terms gives the claim for $u \nmid D$. For $u \mid D$ the forms with $u \mid a$ add $u\text{Per}_D$, and the term -1 is the single ramified neighbour of discriminant $4D$. The identity for $u_j(u^2 \cdot)$: $u^2\tau_{Q'} = uz_Q$ is itself the first neighbour. [The bijection between non-selected neighbours and the selected neighbours of the other representatives is the step to write out; the ramified count too. Numerically $c = +2, +4$ ($u = 3, 5$, $u \mid D$), -2 (split), 0 (inert), each to four digits, and $\text{Per}(u_j) = \text{Per}(u_j(u^2 \cdot))$ exactly.] $\square$

Lemma 8 (the Gram matrix). On $\Gamma_0(u^2)\backslash\mathbb{H}$, with $f_m(z) = u_j(u^m z)$: $\langle f_0, f_0 \rangle = u(u + 1)\|u_j\|_{\text{SL}_2(\mathbb{Z})}^2$, $\langle f_1, f_0 \rangle = \langle f_2, f_1 \rangle = \frac{\lambda_j(u)\sqrt{u}}{u+1}\langle f_0, f_0 \rangle$ and $\langle f_2, f_0 \rangle = \frac{\lambda_j(u)^2 - 1 - 1/u}{u+1}\langle f_0, f_0 \rangle$.

Sketch. The first entry is the index; the second is [3, Lemma 2.4] (stated there for holomorphic forms; the unfolding is identical for Maass forms). The third follows by the same unfolding from $\langle f|B_{u^2}, f \rangle$ and the Hecke relation $\lambda(u^2) = \lambda(u)^2 - 1$; it reads $(\lambda_j(u^2) - 1/u)/(u + 1)$. [Write out or locate; numerically $0.3001, -0.3181, -0.1095$ at $u = 2, 3, 5$, matching the formula to four digits.] $\square$

Proof of Theorem 2 from the lemmas. Theorem 1 restricted to the span of f_0, f_1, f_2 gives the coefficient $\sqrt{u} \sum_k \text{Per}_{\mathcal{W}_u}(v_k) \tilde{L}(v_k)$ over an orthonormal basis v_k of that span, which is the basis-free quantity $\sqrt{u} \mathbf{P}^\top \mathbf{G}^{-1} \mathbf{L}$ with $\mathbf{P}_m = \text{Per}_{\mathcal{W}_u}(f_m)$, $\mathbf{G}_{mn} = \langle f_m, f_n \rangle$, $\mathbf{L}_m = \tilde{L}_{f_m}$. Dividing by the $u = 1$ coefficient $\text{Per}_D(u_j) \tilde{L}_j / \|u_j\|^2$ and inserting the lemmas gives (2); the phase u^{-it_j} is from Theorem 1. The asymptotics: as $u \rightarrow \infty$, $G \rightarrow I$, $L \rightarrow (1, 0, 0)$, and $p_0/(u(u + 1)) \rightarrow \lambda_j(u)u^{-1}$ or u^{-1} according as $u \nmid D$ or $u \mid D$. $\square$

4 Numerical confirmation

All data are the pieces of part IV computed to $Y = 10^7$ (`piece-divset.ts`, unrestricted object), sharp Riesz means on 4096 log-spaced points and smooth windows on 401; amplitudes and phases by joint least squares at the level-one lines with the dominant level- u^2 lines as nuisance frequencies (Strömberg's $\Gamma_0(9)$ eigenvalues at $u = 3$ [6], the six strongest data-driven peaks at $u = 5$). The predictions of Theorem 2 were computed twice: from (2), and directly by evaluating the Maass form at the orbit representatives of $\mathcal{W}_u$, integrating the Gram matrix over the cosets

of $\Gamma_0(u^2)$, and orthonormalising; the two agree to four digits, and the direct route also checks $\langle f_0, f_0 \rangle / \|u_j\|^2 = u(u+1)$ and the Iwaniec–Luo–Sarnak entry.

Table 1: The first even line $t_1 = 13.7798$: signed ratio of its amplitude in piece u to piece 1 (sign – when the phase is shifted by π beyond $-t_1 \log u$). “pred” from (2) for odd u and from the direct computation for $u = 2$; “obs” from the sharp grids. Predicted $|r| < 0.1$ is below the noise of these grids.

u , type of u in $\mathbb{Q}(\sqrt{D})$	pred		obs	D
2, $D \equiv 0 \pmod{4}$, $D/4 \equiv 2, 3 \pmod{4}$	+0.850	+0.852, +0.865, +0.899	–4, –8, –20	
2, $D \equiv 0 \pmod{4}$, $D/4 \equiv 1 \pmod{4}$	+0.945	(piece $u = 2$ not yet fitted)	–12	
2, split ($D \equiv 1 \pmod{8}$)	–0.341	–0.353, –0.379, –0.428	–7, –15, –23	
2, inert ($D \equiv 5 \pmod{8}$)	–0.068	–0.065, –0.061, –0.071	–3, –11, –19	
3, ramified	+0.695	+0.655, +0.726	–3, –15	
3, split	–0.311	–0.285, –0.320, –0.332	–8, –11, –20	
3, inert	+0.192	+0.196, (+0.341)	–4, –7	
5, ramified	+0.421	+0.431	–15	
5, split	–0.076	(noise)	–4, –11	
5, inert	+0.089	(noise)	–3, –8	

Table 2: The first even newform of level 3 ($t = 5.0987$, Fricke +1) in the smooth piece $u = 3$: predicted amplitude and phase from Theorem 3 with the LMFDB coefficients, against the fit (all $\Gamma_0(9)$ even lines and four level-one lines in the fit).

D	3 in $\mathbb{Q}(\sqrt{D})$	C_{pred}	C_{obs}	ratio	$\Delta\varphi$
–3	ramified	0.3200	0.3262	1.020	–0.008
–15	ramified	0.2016	0.1946	0.965	+0.024
–8	split	0.1404	0.1426	1.016	+0.072
–11	split	0.1371	0.1370	1.000	+0.050
–20	split	0.1945	0.1931	0.993	+0.032
–4	inert	0	0.019	—	—
–7	inert	0	0.006	—	—

The case $u = 2$. The local constants are not integers: $\text{Per}_{\mathcal{W}_2}(u_1(2\cdot))/\text{Per}_D(u_1) = 4.912$ for $D/4 \equiv 2, 3 \pmod{4}$ and 5.278 for $D/4 \equiv 1 \pmod{4}$, 0.321 for $D \equiv 1 \pmod{8}$, 1.373 for $D \equiv 5 \pmod{8}$; so the multiplicity of Lemma 6 is not class-independent at $u = 2$ and the closed form (2) does not apply. The direct computation does, and matches the data (Table 1). [\[2-adic local analysis.\]](#)

Composite u . For squarefree $u = u_1 u_2$ the family condition factors over the primes and the oldform space is $\{u_j(dz) : d \mid u^2\}$, of dimension 9 for $u = 6$. The direct computation at level 36 (Gram matrix over the 72 cosets, $\|f_0\|^2/\|u_j\|^2 = 72.000$) gives $r_1(6; -12) = +0.6573$ and $r_1(6; -8) = -0.2642$, against the products $r_1(2; -12)r_1(3; -12) = 0.945 \times 0.695 = 0.657$ and $r_1(2; -8)r_1(3; -8) = 0.850 \times (-0.311) = -0.264$; and the 9×9 Gram matrix is the tensor product of the 3×3 matrices at levels 4 and 9 entry by entry ($0.0781 = 0.7303 \times 0.1069$, $-0.2323 = 0.7303 \times (-0.3181)$, $-0.0955 = 0.3001 \times (-0.3181)$, ...). So the projection factors over the primes dividing u and the multiplicativity $r_j(u; D) = \prod_{p|u} r_j(p; D)$ holds on the theory side; its proof is the observation that the period vector and the Gram matrix are tensor products,

which follows from Lemmas 6–8 applied prime by prime. [Write out the tensor-product argument.] The data at $u = 6$ are noisy (the level-36 spectrum is dense): the five signs observed agree with the products and the magnitudes scatter by up to 0.2 ($D = -8, -20, -15, -4, -12$: observed $-0.449, -0.320, -0.295, +0.111, +0.549$ against $-0.264, -0.264, -0.237, +0.163, +0.657$).

The cycloidal lines. At $u = 3$ the pieces are dominated by the lines 3.536, 5.504, 6.647, 7.432, 8.698, which Strömberg attributes to forms of the cycloidal group Γ^3 [6]; their coefficients are not tabulated and their amplitudes in piece 3 cannot be predicted here.

5 What is open

1. The three lemmas, to be proved in full (Section 3); the vanishing of the newform period for inert u .
2. The 2-adic case; the tensor-product argument for composite u .
3. The lines of the cycloidal-group forms, and any bound for the newform lines uniform in the newform: the level-aspect equidistribution of [4, 2], route 1b of the knowledge base.
4. The error term uniform in u (part IV, Remark GAPS), which is the conjecture of part III itself.

AI-assisted research: disclosure

This work was carried out with heavy use of large language models, the Anthropic models Claude Fable 5.1 and Claude Opus 5, used interactively by the author in September 2026 as research assistants. The author posed the questions, set the direction and the scope, and made every decision about what to pursue, claim and submit. The model derived the identities, wrote and ran the software behind every number, searched the literature and drafted the text, in dialogue with the author, who verified the code and the numerical claims against the repository record. The repository (<https://github.com/gewure/ulamnd>) contains the scripts, the data, and the knowledge base with the record of errors.

References

- [1] Benedict Gross, Winfried Kohnen, and Don Zagier. Heegner points and derivatives of L -series. II. *Math. Ann.*, 278:497–562, 1987.
- [2] Peter Humphries and Asbjørn Christian Nordentoft. Sparse equidistribution of geometric invariants of real quadratic fields. *J. Eur. Math. Soc.*, 2025. to appear; arXiv:2211.05890.
- [3] Henryk Iwaniec, Wenzhi Luo, and Peter Sarnak. Low lying zeros of families of L -functions. *Publ. Math. Inst. Hautes Études Sci.*, 91:55–131, 2000.
- [4] Sheng-Chi Liu, Riad Masri, and Matthew P. Young. Subconvexity and equidistribution of Heegner points in the level aspect. *Compos. Math.*, 149:1150–1174, 2013. arXiv:1206.3208.
- [5] Jasper Reichardt. The pair singular series of a polynomial IV: the second spectrum, 2026. Working draft, <https://github.com/gewure/ulamnd>, research/paper-IV.
- [6] Fredrik Strömberg. Newforms and spectral multiplicities for $\Gamma_0(9)$. *Proc. Lond. Math. Soc.*, 2012. arXiv:1106.5741.