

50.007


Machine Learning

Lu, Wei



Hidden Markov Model (III)

Hidden Markov Model Parameterization



$$p(x_1, x_2, \dots, x_{n-1}, x_n, y_0, y_1, y_2, \dots, y_{n-1}, y_n, y_{n+1})$$


$$\underbrace{\prod_{j=0}^n a_{y_j, y_{j+1}}}_{\text{Transition probabilities}} \times \underbrace{\prod_{j=1}^n b_{y_j}(x_j)}_{\text{Emission probabilities}}$$

Transition probabilities

Emission probabilities

Hidden Markov Model

Supervised Learning



$$p(x_1, x_2, \dots, x_{n-1}, x_n, y_0, y_1, y_2, \dots, y_{n-1}, y_n, y_{n+1})$$

$$a_{u,v} = \frac{\text{count}(u,v)}{\text{count}(u)}$$

$$b_u(o) = \frac{\text{count}(u \rightarrow o)}{\text{count}(u)}$$


Hidden Markov Model Decoding

 $a_{u,v}$

| $u \setminus v$ | A | B | STOP |
|-----------------|-----|-----|------|
| START | 1.0 | 0.0 | 0.0 |
| A | 0.5 | 0.5 | 0.0 |
| B | 0.0 | 0.8 | 0.2 |

 $b_u(o)$

| $u \setminus o$ | “the” | “dog” |
|-----------------|-------|-------|
| A | 0.9 | 0.1 |
| B | 0.1 | 0.9 |




$\mathbf{x} = \text{the, dog, the}$

What is the most probable \mathbf{y} sequence for the given \mathbf{x} sequence?

Hidden Markov Model

Unsupervised Learning

We don't know the model parameters, but only know there are two possible states: A , B .



$\mathbf{x} = \text{the, dog, the}$

What is the most probable \mathbf{y} sequence for the given \mathbf{x} sequence?

Question

How to solve the unsupervised learning problem for HMM?

Expectation Maximization

E-Step

Find for each input its membership

M-Step

Update the model parameters

Hard EM for HMM

E-Step

For each input sequence,
find its most probable output sequence.



This is the decoding procedure!

Hard EM for HMM

M-Step

Update the model parameters,
based on the training sequence pairs.



This is the supervised learning procedure!

Soft EM for HMM

E-Step

Run Viterbi, and then collect counts from each instance

M-Step

$$a_{u,v} = \frac{\text{count}(u,v)}{\text{count}(u)} \quad b_u(o) = \frac{\text{count}(u \rightarrow o)}{\text{count}(u)}$$

Soft EM for HMM

E-Step

Run some algorithm to collect
fractional counts from each instance

M-Step

$$a_{u,v} = \frac{\text{count}(u,v)}{\text{count}(u)} \quad b_u(o) = \frac{\text{count}(u \rightarrow o)}{\text{count}(u)}$$

Soft EM for HMM

Finding the fractional count

A distribution over possible \mathbf{y} s



$$\text{count}(u, v) = \sum_{i=1}^m \sum_{\mathbf{y}} p(\mathbf{y} | \mathbf{x}^{(i)}) \text{count}(\mathbf{x}^{(i)}, \mathbf{y}, u \rightarrow v)$$

The number of times we see a transition from u to v in the sequence pair $(\mathbf{x}^{(i)}, \mathbf{y})$

Soft EM for HMM

$$\sum_{\mathbf{y}} p(\mathbf{y}|\mathbf{x}) \text{count}(\mathbf{x}, \mathbf{y}, u \rightarrow v)$$



Soft EM for HMM


$$\begin{aligned} & \sum_{\mathbf{y}} p(\mathbf{y}|\mathbf{x}) \text{count}(\mathbf{x}, \mathbf{y}, u \rightarrow v) \\ &= \sum_{\mathbf{y}} p(\mathbf{y}|\mathbf{x}) \sum_{j=0}^n \text{count}(\mathbf{x}, \mathbf{y}, u \rightarrow v, j) \end{aligned}$$

Soft EM for HMM

$$\begin{aligned} & \sum_{\mathbf{y}} p(\mathbf{y}|\mathbf{x}) \text{count}(\mathbf{x}, \mathbf{y}, u \rightarrow v) \\ &= \sum_{\mathbf{y}} p(\mathbf{y}|\mathbf{x}) \sum_{j=0}^n \text{count}(\mathbf{x}, \mathbf{y}, u \rightarrow v, j) \\ &= \sum_{j=0}^n \sum_{\mathbf{y}} p(\mathbf{y}|\mathbf{x}) \text{count}(\mathbf{x}, \mathbf{y}, u \rightarrow v, j) \end{aligned}$$

Soft EM for HMM

$$\begin{aligned} & \sum_{\mathbf{y}} p(\mathbf{y}|\mathbf{x}) \text{count}(\mathbf{x}, \mathbf{y}, u \rightarrow v) \\ &= \sum_{\mathbf{y}} p(\mathbf{y}|\mathbf{x}) \sum_{j=0}^n \text{count}(\mathbf{x}, \mathbf{y}, u \rightarrow v, j) \\ &= \sum_{j=0}^n \sum_{\mathbf{y}} p(\mathbf{y}|\mathbf{x}) \underbrace{\text{count}(\mathbf{x}, \mathbf{y}, u \rightarrow v, j)} \end{aligned}$$



This is an indicator function!

Soft EM for HMM

$$\sum_{\mathbf{y}} p(\mathbf{y}|\mathbf{x}) \text{count}(\mathbf{x}, \mathbf{y}, u \rightarrow v, j)$$



Soft EM for HMM

$$\begin{aligned} & \sum_{\mathbf{y}} p(\mathbf{y}|\mathbf{x}) \text{count}(\mathbf{x}, \mathbf{y}, u \rightarrow v, j) \\ &= \sum_{y_0, \dots, y_{n+1}} \left[p(y_0, \dots, y_j, y_{j+1}, \dots, y_{n+1} | \mathbf{x}) \right. \\ & \quad \left. \times \text{count}(\mathbf{x}, \mathbf{y}, u \rightarrow v, j) \right] \end{aligned}$$



Soft EM for HMM

$$\begin{aligned} & \sum_{\mathbf{y}} p(\mathbf{y}|\mathbf{x}) \text{count}(\mathbf{x}, \mathbf{y}, u \rightarrow v, j) \\ &= \sum_{y_0, \dots, y_{n+1}} \left[p(y_0, \dots, y_j, y_{j+1}, \dots, y_{n+1} | \mathbf{x}) \right. \\ & \quad \left. \times \text{count}(\mathbf{x}, \mathbf{y}, u \rightarrow v, j) \right] \\ &= \sum_{y_0, \dots, y_{j-1}, y_{j+2}, \dots, y_{n+1}} \left[p(y_0, \dots, y_{j-1}, \right. \\ & \quad \left. y_j = u, y_{j+1} = v, y_{j+2}, \dots, y_{n+1} | \mathbf{x}) \right] \end{aligned}$$

Soft EM for HMM

$$\begin{aligned} & \sum_{\mathbf{y}} p(\mathbf{y}|\mathbf{x}) \text{count}(\mathbf{x}, \mathbf{y}, u \rightarrow v, j) \\ &= \sum_{y_0, \dots, y_{n+1}} \left[p(y_0, \dots, y_j, y_{j+1}, \dots, y_{n+1} | \mathbf{x}) \right. \\ & \quad \left. \times \text{count}(\mathbf{x}, \mathbf{y}, u \rightarrow v, j) \right] \\ &= \sum_{y_0, \dots, y_{j-1}, y_{j+2}, \dots, y_{n+1}} \left[p(y_0, \dots, y_{j-1}, \right. \\ & \quad \left. y_j = u, y_{j+1} = v, y_{j+2}, \dots, y_{n+1} | \mathbf{x}) \right] \\ &= p(y_j = u, y_{j+1} = v | \mathbf{x}) \end{aligned}$$



Soft EM for HMM

$$\begin{aligned} & \sum_{\mathbf{y}} p(\mathbf{y}|\mathbf{x}) \text{count}(\mathbf{x}, \mathbf{y}, u, j) \\ &= \sum_{y_0, \dots, y_{n+1}} \left[p(y_0, \dots, y_j, y_{j+1}, \dots, y_{n+1} | \mathbf{x}) \right. \\ & \quad \left. \times \text{count}(\mathbf{x}, \mathbf{y}, u, j) \right] \\ &= \sum_{y_0, \dots, y_{j-1}, y_{j+1}, \dots, y_{n+1}} \left[p(y_0, \dots, y_{j-1}, \right. \\ & \quad \left. y_j = u, y_{j+1}, y_{j+2}, \dots, y_{n+1} | \mathbf{x}) \right] \\ &= p(y_j = u | \mathbf{x}) \end{aligned}$$

Inference in HMM

$$p(y_j = u | \mathbf{x}; \theta)$$



Model Parameters (α, β)

Inference in HMM

$$p(y_j = u | \mathbf{x}; \theta)$$

$$= \frac{p(y_j = u, \mathbf{x}; \theta)}{p(\mathbf{x}; \theta)}$$

Inference in HMM

$$\begin{aligned} & p(y_j = u | \mathbf{x}; \theta) \\ &= \frac{p(y_j = u, \mathbf{x}; \theta)}{p(\mathbf{x}; \theta)} \\ &= \frac{p(x_1, x_2, \dots, x_{j-1}, y_j = u, x_j, x_{j+1}, \dots, x_n; \theta)}{p(x_1, x_2, \dots, x_n; \theta)} \end{aligned}$$

Inference in HMM

$$\begin{aligned} & p(y_j = u | \mathbf{x}; \theta) \\ &= \frac{p(y_j = u, \mathbf{x}; \theta)}{p(\mathbf{x}; \theta)} \\ &= \frac{p(x_1, x_2, \dots, x_{j-1}, y_j = u, x_j, x_{j+1}, \dots, x_n; \theta)}{p(x_1, x_2, \dots, x_n; \theta)} \end{aligned}$$



What is the relation between the numerator and the denominator?

Inference in HMM

$$\begin{aligned} & p(y_j = u | \mathbf{x}; \theta) \\ &= \frac{p(y_j = u, \mathbf{x}; \theta)}{p(\mathbf{x}; \theta)} \\ &= \frac{p(x_1, x_2, \dots, x_{j-1}, y_j = u, x_j, x_{j+1}, \dots, x_n; \theta)}{p(x_1, x_2, \dots, x_n; \theta)} \\ &= \frac{p(x_1, x_2, \dots, x_{j-1}, y_j = u, x_j, x_{j+1}, \dots, x_n; \theta)}{\sum_v p(x_1, x_2, \dots, x_{j-1}, y_j = v, x_j, x_{j+1}, \dots, x_n; \theta)} \end{aligned}$$


Inference in HMM

$$\begin{aligned} & p(y_j = u | \mathbf{x}; \theta) \\ &= \frac{p(y_j = u, \mathbf{x}; \theta)}{p(\mathbf{x}; \theta)} \\ &= \frac{p(x_1, x_2, \dots, x_{j-1}, y_j = u, x_j, x_{j+1}, \dots, x_n; \theta)}{p(x_1, x_2, \dots, x_n; \theta)} \\ &= \frac{p(x_1, x_2, \dots, x_{j-1}, y_j = u, x_j, x_{j+1}, \dots, x_n; \theta)}{\sum_v p(x_1, x_2, \dots, x_{j-1}, y_j = v, x_j, x_{j+1}, \dots, x_n; \theta)} \end{aligned}$$

$$p(x_1, x_2, \dots, x_{j-1}, y_j = u, x_j, x_{j+1}, \dots, x_n; \theta)$$


Now let us take a closer look at this joint probability.

Inference in HMM




$$p(x_1, x_2, \dots, x_{j-1}, y_j = u, x_j, x_{j+1}, \dots, x_n; \theta)$$

Inference in HMM




$$p(x_1, x_2, \dots, x_{j-1}, y_j = u, x_j, x_{j+1}, \dots, x_n; \theta)$$

Inference in HMM




$$\begin{aligned} & p(x_1, x_2, \dots, x_{j-1}, y_j = u, x_j, x_{j+1}, \dots, x_n; \theta) \\ &= p(x_1, x_2, \dots, x_{j-1}, y_j = u; \theta) \\ & \quad p(x_j, x_{j+1}, \dots, x_n | y_j = u; \theta) \end{aligned}$$

Inference in HMM




$$\begin{aligned}
 & p(x_1, x_2, \dots, x_{j-1}, y_j = u, x_j, x_{j+1}, \dots, x_n; \theta) \\
 &= p(x_1, x_2, \dots, x_{j-1}, y_j = u; \theta) \\
 &\quad \times p(x_j, x_{j+1}, \dots, x_n | y_j = u; \theta)
 \end{aligned}$$

Inference in HMM




$$\begin{aligned}
 & p(x_1, x_2, \dots, x_{j-1}, y_j = u, x_j, x_{j+1}, \dots, x_n; \theta) \\
 &= p(x_1, x_2, \dots, x_{j-1}, y_j = u; \theta) \\
 &\quad \times p(x_j, x_{j+1}, \dots, x_n | y_j = u; \theta)
 \end{aligned}$$

Inference in HMM



$$\begin{aligned}
 & p(x_1, x_2, \dots, x_{j-1}, y_j = u, x_j, x_{j+1}, \dots, x_n; \theta) \\
 &= p(x_1, x_2, \dots, x_{j-1}, y_j = u; \theta) \alpha_u(j) \\
 &\quad \times p(x_j, x_{j+1}, \dots, x_n | y_j = u; \theta) \beta_u(j)
 \end{aligned}$$

Inference in HMM




$$p(x_1, \dots, x_{j-1}, y_j = u, x_j, \dots, x_n; \theta) = \alpha_u(j) \beta_u(j)$$

$$p(y_j = u | \mathbf{x}; \theta)$$

$$= \frac{p(x_1, x_2, \dots, x_{j-1}, y_j = u, x_j, x_{j+1}, \dots, x_n; \theta)}{\sum_v p(x_1, x_2, \dots, x_{k-1}, y_k = v, x_k, x_{k+1}, \dots, x_n; \theta)}$$


Inference in HMM



$$p(x_1, \dots, x_{j-1}, y_j = u, x_j, \dots, x_n; \theta) = \alpha_u(j) \beta_u(j)$$

$$p(y_j = u | \mathbf{x}; \theta) = \frac{\alpha_u(j) \beta_u(j)}{\sum_v \alpha_v(j) \beta_v(j)}$$


Inference in HMM



$$p(x_1, \dots, x_{j-1}, y_j = u, x_j, \dots, x_n; \theta) = \alpha_u(j) \beta_u(j)$$

$$p(y_j = u, y_{j+1} = v | \mathbf{x}; \theta) = \frac{p(x_1, x_2, \dots, x_{j-1}, y_j = u, x_j, y_{j+1} = v, x_{j+1}, \dots, x_n; \theta)}{\sum_v p(x_1, x_2, \dots, x_{j-1}, y_j = v, x_j, y_{j+1} = v, x_{j+1}, \dots, x_n; \theta)}$$


Inference in HMM




$$p(y_j = u, y_{j+1} = v | \mathbf{x}; \theta) = \frac{\alpha_u(j) \cdot b_u(x_j) \cdot a_{u,v} \cdot \beta_v(j+1)}{\sum_v \alpha_v(j) \beta_v(j+1)}$$

Question


How to find an efficient procedure to calculate forward and backward probabilities?




Viterbi Algorithm



Viterbi Algorithm



Inference in HMM




$$\alpha_{\text{STOP}}(n+1)$$

$$= p(x_1, \dots, x_{j-1}, x_j, \dots, x_n; \theta)$$

$$= \sum_{y_1, \dots, y_n} p(x_1, \dots, x_n, y_0, y_1, \dots, y_n, y_{n+1}; \theta)$$

Inference in HMM



$$\alpha_{\text{STOP}}(n+1)$$


$$= p(x_1, \dots, x_{j-1}, x_j, \dots, x_n; \theta)$$

$$= \sum_{y_1, \dots, y_n} p(x_1, \dots, x_n, y_0, y_1, \dots, y_n, y_{n+1}; \theta)$$

$$\max_{y_1, \dots, y_n} p(x_1, \dots, x_n, y_0, y_1, \dots, y_n, y_{n+1}; \theta)$$


Decoding
(Viterbi)

Inference in HMM




Inference in HMM

0 1 2 j $j+1$ $n-1$ n $n+1$




The sum of the scores of all paths from START to STOP.




$$\sum_{y_1, \dots, y_n} p(x_1, \dots, x_n, y_0, y_1, \dots, y_n, y_{n+1}; \theta)$$

Forward-Backward Algorithm



Forward-Backward Algorithm




The sum of the scores of all paths from this node to STOP

Forward-Backward Algorithm

0 1 2 j $j+1$ $n-1$ n $n+1$

$$\alpha_u(j)$$

The sum of the scores of all paths from START to this node.



Let us now work out the algorithm on our own

$$\beta_u(j)$$

The sum of the scores of all paths from this node to STOP