

Gradient magnetometry with various types of spin ensembles

Single atomic ensembles, chain of spins & two different ensembles

Iagoba Apellaniz¹ Iñigo Urizar-Lanz¹ Zoltán Zimborás^{1,2,3}
Philipp Hyllus¹ Géza Tóth^{1,3,4}

¹Department of Theoretical Physics, University of the Basque Country UPV/EHU, Bilbao, Spain

²Dahlem Center for Complex Quantum Systems, Freie Universität Berlin, Germany

³Wigner Research Centre for Physics, Hungarian Academy of Sciences, Budapest, Hungary

⁴IKERBASQUE, Basque Foundation for Science, Bilbao, Spain

DPG Meeting of the Atomic, Molecular, Quantum Optics and Photonics Section
(SAMOP)
- March 10, 2023 -





1 Multiparametric Quantum Metrology

- Cramér-Rao precision bound and quantum Fisher information
- Multiparametric qFI matrix and simultaneous estimation

2 System setup and precision bounds of the gradient parameter estimation for various states

- Gradient magnetometry and basic setup of the system
- Precision bounds for various systems and different spin states

3 Conclusions




1 Multiparametric Quantum Metrology

- Cramér-Rao precision bound and quantum Fisher information
- Multiparametric qFI matrix and simultaneous estimation

2 System setup and precision bounds of the gradient parameter estimation for various states


- Gradient magnetometry and basic setup of the system
- Precision bounds for various systems and different spin states

3 Conclusions



- The *quantum Cramér-Rao (qCR) bound* provides an upper bound for the precision


$$\frac{1}{(\Delta\theta)^2} \leq \mu \mathcal{F}_Q[\varrho, A].$$



- The *quantum Cramér-Rao (qCR) bound* provides an upper bound for the precision

$$\frac{1}{(\Delta\theta)^2} \leq \mu \mathcal{F}_Q[\varrho, A].$$

- **Objective:** Minimize $(\Delta\theta)^2$, or equivalently maximize $\mathcal{F}_Q[\varrho, A]$.



- The *quantum Cramér-Rao (qCR) bound* provides an upper bound for the precision

$$\frac{1}{(\Delta\theta)^2} \leq \mu \mathcal{F}_Q[\varrho, A].$$

- **Objective:** Minimize $(\Delta\theta)^2$, or equivalently maximize $\mathcal{F}_Q[\varrho, A]$.
- **Quantum Fisher information**

$$\mathcal{F}_Q[\varrho, A] = 2 \sum_{\lambda \neq \mu} \frac{(p_\lambda - p_\mu)^2}{p_\lambda + p_\mu} |\langle \lambda | A | \mu \rangle|^2$$

written on the eigenbasis of the state, $\varrho = \sum p_\lambda |\lambda\rangle\langle\lambda|$.

[M.G.A. Paris (2009), IJQI 7, 125]



$$\mathcal{F}_Q[\varrho, A] = 2 \sum_{\lambda \neq \mu} \frac{(p_\lambda - p_\mu)^2}{p_\lambda + p_\mu} |\langle \lambda | A | \mu \rangle|^2$$

Properties of the qFI for a single parameter estimation problem

- It is independent of the measurement. **An optimal measurement exists** though, which saturates the qCR bound.

[M G A Paris (2009), IJQI 7, 125]

[G Tóth *et al.* (2014), JPA:MT 47, 424006]

[L. Pezzé *et al.* (2018), RMP 90, 035005]



$$\mathcal{F}_Q[\varrho, A] = 2 \sum_{\lambda \neq \mu} \frac{(p_\lambda - p_\mu)^2}{p_\lambda + p_\mu} |\langle \lambda | A | \mu \rangle|^2$$

Properties of the qFI for a single parameter estimation problem

- 1 It is independent of the measurement. An optimal measurement exists though, which saturates the qCR bound.
- 2 It is convex over the set of quantum states. Hence, it is maximized by a pure state.

[M G A Paris (2009), IJQI 7, 125]

[G Tóth *et al.* (2014), JPA:MT 47, 424006]

[L. Pezzé *et al.* (2018), RMP 90, 035005]

$$\mathcal{F}_Q[\varrho, A] = 2 \sum_{\lambda \neq \mu} \frac{(p_\lambda - p_\mu)^2}{p_\lambda + p_\mu} |\langle \lambda | A | \mu \rangle|^2$$

Properties of the qFI for a single parameter estimation problem

- 1 It is independent of the measurement. An optimal measurement exists though, which saturates the qCR bound.
- 2 It is convex over the set of quantum states. Hence, it is maximized by a pure state.
- 3 For pure states $\mathcal{F}_Q[|\Psi\rangle, A] = 4(\Delta A)^2_\Psi$.

[M G A Paris (2009), IJQI 7, 125]

[G Tóth *et al.* (2014), JPA:MT 47, 424006]

[L. Pezzé *et al.* (2018), RMP 90, 035005]



Entanglement

- 1 Separable states can achieve at most the so called Shot-noise limit (SNL),

$$\mathcal{F}_Q[\varrho_{\text{sep}}, H] \sim N.$$

- 2 An ultimate limit is obtained maximizing the qFI over all pure states

$$\max_{|\Psi\rangle} \mathcal{F}_Q[|\Psi\rangle, H] = N^2,$$

which is called the Heisenberg limit.

Hence, entanglement is **needed** to overcome the SNL.

[V Giovannetti *et al.* (2004), *Sci.* **306** 1330]



Entanglement

- 1 Separable states can achieve at most the so called Shot-noise limit (SNL),

$$\mathcal{F}_Q[\varrho_{\text{sep}}, H] \sim N.$$

- 2 An ultimate limit is obtained maximizing the qFI over all pure states

$$\max_{|\Psi\rangle} \mathcal{F}_Q[|\Psi\rangle, H] = N^2,$$

which is called the Heisenberg limit.

Hence, entanglement is **needed** to overcome the SNL.

[V Giovannetti *et al.* (2004), *Sci.* **306** 1330]

E.g. entanglement criteria based on qFI


- Due to its tight relation with the variance, qFI has been used to improve some entanglement conditions.

[G Tóth (2022), *PRR* **4** 013075]

- Ion chains can be used to estimate the magnetic field as a function of position, $B(x)$.

[Matteo Fadel *et al.*, [arXiv.org:2201.11081](https://arxiv.org/abs/2201.11081)]

- Ion chains can be used to estimate the magnetic field as a function of position, $B(x)$.
[Matteo Fadel *et al.*, [arXiv.org:2201.11081](https://arxiv.org/abs/2201.11081)]
- States invariant to a global rotation of the system have been prepared in elongated traps.




[N Behbood *et al.* (2014), PRL **113** 093601]

[I Urizar-Lanz *et al.* (2013), PRA **88** 013626]

- Ion chains can be used to estimate the magnetic field as a function of position, $B(x)$.

[Matteo Fadel *et al.*, arXiv.org:2201.11081]

- States invariant to a global rotation of the system have been prepared in elongated traps.



[N Behbood *et al.* (2014), PRL **113** 093601]

[I Urizar-Lanz *et al.* (2013), PRA **88** 013626]


- Two distinguishable ensembles of atoms have been prepared with a highly entangled spin state.

[K Langle *et al.* (2018), Sci. **360** 6387]

- Ion chains can be used to estimate the magnetic field as a function of position, $B(x)$.

[Matteo Fadel *et al.*, arXiv.org:2201.11081]

- States invariant to a global rotation of the system have been prepared in elongated traps.



[N Behbood *et al.* (2014), PRL **113** 093601]

[I Urizar-Lanz *et al.* (2013), PRA **88** 013626]

- Two distinguishable ensembles of atoms have been prepared with a highly entangled spin state.

[K Langle *et al.* (2018), Sci. **360** 6387]


We assume that the magnetic field is pointing in the z -direction and its Taylor expansion around the origin is

$$B = (0, 0, B_0) + (0, 0, xB_1) + O(x^2).$$

- Ion chains can be used to estimate the magnetic field as a function of position, $B(x)$.

[Matteo Fadel *et al.*, arXiv.org:2201.11081]

- States invariant to a global rotation of the system have been prepared in elongated traps.



[N Behbood *et al.* (2014), PRL **113** 093601]

[I Urizar-Lanz *et al.* (2013), PRA **88** 013626]

- Two distinguishable ensembles of atoms have been prepared with a highly entangled spin state.


[K Langle *et al.* (2018), Sci. **360** 6387]

We assume that the magnetic field is pointing in the z -direction and its Taylor expansion around the origin is

$$B = (0, 0, B_0) + (0, 0, xB_1) + O(x^2).$$

In general, one cannot avoid a global rotation of the state.

- Ion chains can be used to estimate the magnetic field as a function of position, $B(x)$.
[Matteo Fadel *et al.*, arXiv.org:2201.11081]
- States invariant to a global rotation of the system have been prepared in elongated traps.



[N Behbood *et al.* (2014), PRL **113** 093601]

[I Urizar-Lanz *et al.* (2013), PRA **88** 013626]

- Two distinguishable ensembles of atoms have been prepared with a highly entangled spin state.

[K Langle *et al.* (2018), Sci. **360** 6387]

We assume that the magnetic field is pointing in the z -direction and its Taylor expansion around the origin is

$$B = (0, 0, B_0) + (0, 0, xB_1) + O(x^2).$$

We want to estimate B_1 .



Consider the following evolution for the state

$$\varrho_{\theta} = e^{-i \sum_k A_k \theta_k} \varrho e^{+i \sum_k A_k \theta_k}.$$

- In this case the CR bound is a matrix inequality for the covariance matrix

$$\text{Cov}[\theta_i, \theta_j] \geq \frac{1}{\mu} (\mathcal{F}_Q^{-1})_{i,j},$$

where $\text{Cov}[\theta_i, \theta_j] = \langle \theta_i \theta_j \rangle - \langle \theta_i \rangle \langle \theta_j \rangle$.



Consider the following evolution for the state

$$\varrho_{\theta} = e^{-i \sum_k A_k \theta_k} \varrho e^{+i \sum_k A_k \theta_k}.$$

- In this case the CR bound is a matrix inequality for the covariance matrix

$$\text{Cov}[\theta_i, \theta_j] \geq \frac{1}{\mu} (\mathcal{F}_Q^{-1})_{i,j},$$

where $\text{Cov}[\theta_i, \theta_j] = \langle \theta_i \theta_j \rangle - \langle \theta_i \rangle \langle \theta_j \rangle$.

- The qFI matrix elements are

$$\mathcal{F}_Q[\varrho, A_i, A_j] := (\mathcal{F}_Q)_{i,j} = 2 \sum_{\lambda \neq \mu} \frac{(p_{\lambda} - p_{\mu})^2}{p_{\lambda} + p_{\mu}} \langle \lambda | A_i | \mu \rangle \langle \mu | A_j | \lambda \rangle.$$



Consider the following evolution for the state

$$\varrho_{\theta} = e^{-i \sum_k A_k \theta_k} \varrho e^{+i \sum_k A_k \theta_k}.$$

- In this case the CR bound is a matrix inequality for the covariance matrix

$$\text{Cov}[\theta_i, \theta_j] \geq \frac{1}{\mu} (\mathcal{F}_Q^{-1})_{i,j},$$

where $\text{Cov}[\theta_i, \theta_j] = \langle \theta_i \theta_j \rangle - \langle \theta_i \rangle \langle \theta_j \rangle$.

- The qFI matrix elements are

$$\mathcal{F}_Q[\varrho, A_i, A_j] := (\mathcal{F}_Q)_{i,j} = 2 \sum_{\lambda \neq \mu} \frac{(p_{\lambda} - p_{\mu})^2}{p_{\lambda} + p_{\mu}} \langle \lambda | A_i | \mu \rangle \langle \mu | A_j | \lambda \rangle.$$

- For pure states we have $\mathcal{F}_Q[|\Psi\rangle, A_i, A_j] = 4(\langle A_i A_j \rangle_{\Psi} - \langle A_i \rangle_{\Psi} \langle A_j \rangle_{\Psi})$.



Consider the following evolution for the state

$$\varrho_{\theta} = e^{-i \sum_k A_k \theta_k} \varrho e^{+i \sum_k A_k \theta_k}.$$

- In this case the CR bound is a matrix inequality for the covariance matrix

$$\text{Cov}[\theta_i, \theta_j] \geq \frac{1}{\mu} (\mathcal{F}_Q^{-1})_{i,j},$$

where $\text{Cov}[\theta_i, \theta_j] = \langle \theta_i \theta_j \rangle - \langle \theta_i \rangle \langle \theta_j \rangle$.

- The qFI matrix elements are

$$\mathcal{F}_Q[\varrho, A_i, A_j] := (\mathcal{F}_Q)_{i,j} = 2 \sum_{\lambda \neq \mu} \frac{(p_{\lambda} - p_{\mu})^2}{p_{\lambda} + p_{\mu}} \langle \lambda | A_i | \mu \rangle \langle \mu | A_j | \lambda \rangle.$$

- For pure states we have $\mathcal{F}_Q[|\Psi\rangle, A_i, A_j] = 4(\langle A_i A_j \rangle_{\Psi} - \langle A_i \rangle_{\Psi} \langle A_j \rangle_{\Psi})$.
- When $[A_i, A_j] = 0$, the bounds can be saturated.




1 Multiparametric Quantum Metrology

- Cramér-Rao precision bound and quantum Fisher information
- Multiparametric qFI matrix and simultaneous estimation

2 System setup and precision bounds of the gradient parameter estimation for various states


- Gradient magnetometry and basic setup of the system
- Precision bounds for various systems and different spin states

3 Conclusions



- The system is elongated in one of the spatial directions. The quantum state is a product state between position and spin states,

$$\varrho = \varrho^{(x)} \otimes \varrho^{(s)}.$$



- The system is elongated in one of the spatial directions. The quantum state is a product state between position and spin states,

$$\varrho = \varrho^{(x)} \otimes \varrho^{(s)}.$$

- In this work we assume that the position state is an statistical mixture of point-like particles

$$\varrho^{(x)} = \int \frac{P(x)}{\langle x|x \rangle} |x\rangle\langle x|.$$



- The atoms interact only with the magnetic field, $h^{(n)} = \gamma B_z^{(n)} \otimes j_z^{(n)}$, where $\gamma = g\mu_B$. The collective Hamiltonian is

$$H = \gamma \sum B_z^{(n)} \otimes j_z^{(n)}.$$



- The atoms interact only with the magnetic field, $h^{(n)} = \gamma B_z^{(n)} \otimes j_z^{(n)}$, where $\gamma = g\mu_B$. The collective Hamiltonian is

$$H = \gamma \sum B_z^{(n)} \otimes j_z^{(n)}.$$

- The two unknown parameters are B_0 and B_1 are encoded in b_0 and b_1 acting onto the state with the following unitary operator

$$U = e^{-i(b_0 H_0 + b_1 H_1)},$$

where

$$H_0 := J_z = \sum_n j_z^{(n)} \quad \text{and} \quad H_1 = \sum_n x^{(n)} \otimes j_z^{(n)}.$$



- The atoms interact only with the magnetic field, $h^{(n)} = \gamma B_z^{(n)} \otimes j_z^{(n)}$, where $\gamma = g\mu_B$. The collective Hamiltonian is

$$H = \gamma \sum B_z^{(n)} \otimes j_z^{(n)}.$$

- The two unknown parameters are B_0 and B_1 are encoded in b_0 and b_1 acting onto the state with the following unitary operator

$$U = e^{-i(b_0 H_0 + b_1 H_1)},$$

where

$$H_0 := J_z = \sum_n j_z^{(n)} \quad \text{and} \quad H_1 = \sum_n x^{(n)} \otimes j_z^{(n)}.$$

- Since $[H_0, H_1] = 0$, the precision bounds can be saturated.



- The atoms interact only with the magnetic field, $h^{(n)} = \gamma B_z^{(n)} \otimes j_z^{(n)}$, where $\gamma = g\mu_B$. The collective Hamiltonian is

$$H = \gamma \sum B_z^{(n)} \otimes j_z^{(n)}.$$

- The two unknown parameters are B_0 and B_1 are encoded in b_0 and b_1 acting onto the state with the following unitary operator

$$U = e^{-i(b_0 H_0 + b_1 H_1)},$$

where

$$H_0 := J_z = \sum_n j_z^{(n)} \quad \text{and} \quad H_1 = \sum_n x^{(n)} \otimes j_z^{(n)}.$$

- Since $[H_0, H_1] = 0$, the precision bounds can be saturated.

In the following we are interested on the precision bound for b_1 , the gradient parameter.

Gradient magnetometry and basic setup



Precision bounds for states **insensitive** to the homogeneous B_0

For states that commute with the homogeneous field, $[\varrho, J_z] = 0$, the precision bound is

$$\frac{1}{(\Delta b_1)^2} \leq \mathcal{F}_Q[\varrho, H_1],$$

and it is saturable.

Precision bounds for states **insensitive** to the homogeneous B_0

For states that commute with the homogeneous field, $[\varrho, J_z] = 0$, the precision bound is

$$\frac{1}{(\Delta b_1)^2} \leq \mathcal{F}_Q[\varrho, H_1],$$

and it is saturable.

- For statistical mixtures of point-like particles

$$\frac{1}{(\Delta b_1)^2} \leq \sum_{n,m} \int x_n x_m P(x) dx \mathcal{F}_Q[\varrho^{(s)}, j_z^{(n)}, j_z^{(m)}]$$

Gradient magnetometry and basic setup

Precision bounds for states **insensitive** to the homogeneous B_0

For states that commute with the homogeneous field, $[\varrho, J_z] = 0$, the precision bound is

$$\frac{1}{(\Delta b_1)^2} \leq \mathcal{F}_Q[\varrho, H_1],$$

and it is saturable.

- For statistical mixtures of point-like particles

$$\frac{1}{(\Delta b_1)^2} \leq \sum_{n,m} \int x_n x_m P(x) dx \mathcal{F}_Q[\varrho^{(s)}, j_z^{(n)}, j_z^{(m)}]$$

Precision bounds for states **sensitive** to the homogeneous B_0

For states sensitive to global rotations of the spin state, the precision bound is

$$\frac{1}{(\Delta b_1)^2} \leq (\mathcal{F}_Q)_{1,1} - \frac{(\mathcal{F}_Q)_{0,1}^2}{(\mathcal{F}_Q)_{0,0}}.$$

Gradient magnetometry and basic setup

Precision bounds for states **insensitive** to the homogeneous B_0

For states that commute with the homogeneous field, $[\varrho, J_z] = 0$, the precision bound is

$$\frac{1}{(\Delta b_1)^2} \leq \mathcal{F}_Q[\varrho, H_1],$$

and it is saturable.

- For statistical mixtures of point-like particles

$$\frac{1}{(\Delta b_1)^2} \leq \sum_{n,m} \int x_n x_m P(x) dx \mathcal{F}_Q[\varrho^{(s)}, j_z^{(n)}, j_z^{(m)}]$$

Precision bounds for states **sensitive** to the homogeneous B_0

For states sensitive to global rotations of the spin state, the precision bound is

$$\frac{1}{(\Delta b_1)^2} \leq (\mathcal{F}_Q)_{1,1} - \frac{(\mathcal{F}_Q)_{0,1}^2}{(\mathcal{F}_Q)_{0,0}}.$$

- For statistical mixtures of point-like particles

$$\frac{1}{(\Delta b_1)^2} \leq \sum_{n,m} \int x_n x_m P(x) dx \mathcal{F}_Q[\varrho^{(s)}, j_z^{(n)}, j_z^{(m)}] - \frac{\left(\sum_n \int x_n P(x) dx \mathcal{F}_Q[\varrho^{(s)}, j_z^{(n)}, J_z] \right)^2}{\mathcal{F}_Q[\varrho^{(s)}, J_z]}.$$



1 Multiparametric Quantum Metrology

- Cramér-Rao precision bound and quantum Fisher information
- Multiparametric qFI matrix and simultaneous estimation

2 System setup and precision bounds of the gradient parameter estimation for various states

- Gradient magnetometry and basic setup of the system
- Precision bounds for various systems and different spin states


3 Conclusions

$$P(x) = \prod_n \delta(x_n - na).$$

Chain of qubits

$$P(x) = \prod_n \delta(x_n - na).$$

Totally polarized $|0\rangle_y^{\otimes N}$ state under a magnetic field pointing towards the z-direction



$$\frac{1}{(\Delta b_1)^2} \leq \sum_{n,m} n m a^2 \mathcal{F}_Q[|0\rangle_y^{\otimes N}, j_z^{(n)}, j_z^{(m)}] - \frac{\left(\sum_n n a \mathcal{F}_Q[|0\rangle_y^{\otimes N}, j_z^{(n)}, J_z]\right)^2}{\mathcal{F}_Q[|0\rangle_y^{\otimes N}, J_z]}$$

$$P(x) = \prod_n \delta(x_n - na).$$


- Mean particle position:

$$\mu = a \frac{N+1}{2}$$

- Variance of the particle positions:

$$\sigma^2 = a^2 \frac{N^2 - 1}{12}$$

Totally polarized $|0\rangle_y^{\otimes N}$ state under a magnetic field pointing towards the z-direction




$$\begin{aligned} \frac{1}{(\Delta b_1)^2} &\leq \sum_{n,m} n m a^2 \mathcal{F}_Q[|0\rangle_y^{\otimes N}, j_z^{(n)}, j_z^{(m)}] - \frac{\left(\sum_n n a \mathcal{F}_Q[|0\rangle_y^{\otimes N}, j_z^{(n)}, J_z] \right)^2}{\mathcal{F}_Q[|0\rangle_y^{\otimes N}, J_z]} \\ &= a^2 \frac{N^2 - 1}{12} N = \sigma^2 N. \end{aligned}$$

Single ensemble of point-like spin- $\frac{1}{2}$ atoms

Permutationally invariant PDF

$$P(\mathbf{x}) = \frac{1}{N!} \sum_{k \in S_N} \mathcal{P}_k[P(\mathbf{x})]$$

- $\mu = \int x_n P(\mathbf{x}) d\mathbf{x}$.
- $\sigma^2 = \int x_n^2 P(\mathbf{x}) d\mathbf{x}$, if the origin is at 0.
- $\eta = \int x_n x_m P(\mathbf{x}) d\mathbf{x}$ for $n \neq m$.
 $\eta \in [-\sigma^2/(N-1), \sigma^2]$.



[N Behbood *et al.* (2014), PRL **113** 093601]

Single ensemble of point-like spin- $\frac{1}{2}$ atoms


Permutationally invariant PDF

$$P(\mathbf{x}) = \frac{1}{N!} \sum_{k \in S_N} \mathcal{P}_k[P(\mathbf{x})]$$

- $\mu = \int x_n P(\mathbf{x}) d\mathbf{x}.$
- $\sigma^2 = \int x_n^2 P(\mathbf{x}) d\mathbf{x},$ if the origin is at 0.
- $\eta = \int x_n x_m P(\mathbf{x}) d\mathbf{x}$ for $n \neq m.$
 $\eta \in [-\sigma^2/(N-1), \sigma^2].$

Precision CR bound

$$\frac{1}{(\Delta b_1)^2} \leq (\sigma^2 - \eta) \sum_n \mathcal{F}_Q[\varrho^{(s)}, j_z^{(n)}] \\ + \eta \mathcal{F}_Q[\varrho^{(s)}, J_z]$$



[N Behbood *et al.* (2014), PRL **113** 093601]

Single ensemble of point-like spin- $\frac{1}{2}$ atoms


Permutationally invariant PDF

$$P(\mathbf{x}) = \frac{1}{N!} \sum_{k \in S_N} \mathcal{P}_k[P(\mathbf{x})]$$


- $\mu = \int x_n P(\mathbf{x}) d\mathbf{x}$.
- $\sigma^2 = \int x_n^2 P(\mathbf{x}) d\mathbf{x}$, if the origin is at 0.
- $\eta = \int x_n x_m P(\mathbf{x}) d\mathbf{x}$ for $n \neq m$.
 $\eta \in [-\sigma^2/(N-1), \sigma^2]$.

Precision CR bound

$$\frac{1}{(\Delta b_1)^2} \leq (\sigma^2 - \eta) \sum_n \mathcal{F}_Q[\varrho^{(s)}, j_z^{(n)}] + \eta \mathcal{F}_Q[\varrho^{(s)}, J_z]$$



[N Behbood *et al.* (2014), PRL **113** 093601]





Single ensemble of point-like spin- $\frac{1}{2}$ atoms

Precision bounds for various spin states

Singlet states

$$\varrho^{(s)} = \sum_{\lambda} p_{\lambda} |0, 0, i\rangle \langle 0, 0, i|$$

Its precision bound is

$$\frac{1}{(\Delta b_1)^2} \leq (\sigma^2 - \eta)N.$$



Single ensemble of point-like spin- $\frac{1}{2}$ atoms

Precision bounds for various spin states

Singlet states

$$\varrho^{(s)} = \sum_{\lambda} p_{\lambda} |0, 0, i\rangle \langle 0, 0, i|$$

Its precision bound is

$$\frac{1}{(\Delta b_1)^2} \leq (\sigma^2 - \eta)N.$$

Best separable state

$$\mathcal{F}_Q[|\psi\rangle_{\text{sep}}, j_z^{(n)}, j_z^{(m)}] = \begin{cases} 4(\Delta j_z^{(n)})^2 & \text{if } n = m \\ 0 & \text{otherwise.} \end{cases}$$

Then, the precision is

$$\frac{1}{(\Delta b_1)^2} \leq \sigma^2 N.$$



Single ensemble of point-like spin- $\frac{1}{2}$ atoms

Precision bounds for various spin states

Singlet states

$$\varrho^{(s)} = \sum_{\lambda} p_{\lambda} |0, 0, i\rangle \langle 0, 0, i|$$

Its precision bound is

$$\frac{1}{(\Delta b_1)^2} \leq (\sigma^2 - \eta)N.$$

Best separable state

$$\mathcal{F}_Q[|\psi\rangle_{\text{sep}}, j_z^{(n)}, j_z^{(m)}] = \begin{cases} 4(\Delta j_z^{(n)})^2 & \text{if } n = m \\ 0 & \text{otherwise.} \end{cases}$$

Then, the precision is

$$\frac{1}{(\Delta b_1)^2} \leq \sigma^2 N.$$

|GHZ> states

$$\mathcal{F}_Q[|\text{GHZ}\rangle, j_z^{(n)}] = 1$$

and

$$\mathcal{F}_Q[|\text{GHZ}\rangle_x, J_z] = N^2.$$

Hence,

$$\frac{1}{(\Delta b_1)^2} \leq (\sigma^2 - \eta)N + \eta N^2.$$



Single ensemble of point-like spin- $\frac{1}{2}$ atoms

Precision bounds for various spin states

Singlet states

$$\varrho^{(s)} = \sum_{\lambda} p_{\lambda} |0, 0, i\rangle \langle 0, 0, i|$$

Its precision bound is

$$\frac{1}{(\Delta b_1)^2} \leq (\sigma^2 - \eta)N.$$

Best separable state

$$\mathcal{F}_Q[|\psi\rangle_{\text{sep}}, j_z^{(n)}, j_z^{(m)}] = \begin{cases} 4(\Delta j_z^{(n)})^2 & \text{if } n = m \\ 0 & \text{otherwise.} \end{cases}$$

Then, the precision is

$$\frac{1}{(\Delta b_1)^2} \leq \sigma^2 N.$$

|GHZ> states

$$\mathcal{F}_Q[|\text{GHZ}\rangle, j_z^{(n)}] = 1$$

and

$$\mathcal{F}_Q[|\text{GHZ}\rangle_x, J_z] = N^2.$$

Hence,

$$\frac{1}{(\Delta b_1)^2} \leq (\sigma^2 - \eta)N + \eta N^2.$$


More in PRA 97, 053603 (2018)

$$P(x) = \prod_{n=1}^{N/2} \delta(x_n + a) \prod_{n=N/2+1}^N \delta(x_n - a)$$

The contribution of the position of the particles:

$$\int x_n P(x) dx = \begin{cases} -a \\ +a \end{cases} \quad \text{and} \quad \int x_n x_m P(x) dx = \begin{cases} +a^2 \\ -a^2 \end{cases}$$

- In this case the mean position is $\mu = 0$ and the variance is $\sigma^2 = a^2$.



[K Langle *et al.* (2018), *Sci.* **360** 6387]

$$P(x) = \prod_{n=1}^{N/2} \delta(x_n + a) \prod_{n=N/2+1}^N \delta(x_n - a)$$


The contribution of the position of the particles:

$$\int x_n P(x) dx = \begin{cases} -a \\ +a \end{cases} \quad \text{and} \quad \int x_n x_m P(x) dx = \begin{cases} +a^2 \\ -a^2 \end{cases}$$

- In this case the mean position is $\mu = 0$ and the variance is $\sigma^2 = a^2$.

For spin- $\frac{1}{2}$ system, the state that maximizes the bound is

$$|\psi\rangle = \frac{|\overbrace{0, \dots, 0}^{N/2}, \overbrace{1, \dots, 1}^{N/2}\rangle + |1, \dots, 1, 0, \dots, 0\rangle}{\sqrt{2}}, \quad \text{and} \quad \frac{1}{(\Delta b_1)^2} \leq \sigma^2 N^2.$$



[K Langle *et al.* (2018), *Sci.* **360** 6387]



Product of two equal spin states

For states of the type $|\psi\rangle^{(L)} \otimes |\psi\rangle^{(R)}$, we have that

$$\mathcal{F}_Q[|\psi\rangle^{(L)} \otimes |\psi\rangle^{(R)}, j_z^{(n)}, j_z^{(m)}] = \begin{cases} \mathcal{F}_Q[|\psi\rangle, j_z^{(n)}, j_z^{(m)}] & \text{if } n \text{ and } m \text{ same well} \\ 0 & \text{otherwise} \end{cases}$$



Product of two equal spin states

For states of the type $|\psi\rangle^{(L)} \otimes |\psi\rangle^{(R)}$, we have that

$$\mathcal{F}_Q[|\psi\rangle^{(L)} \otimes |\psi\rangle^{(R)}, j_z^{(n)}, j_z^{(m)}] = \begin{cases} \mathcal{F}_Q[|\psi\rangle, j_z^{(n)}, j_z^{(m)}] & \text{if } n \text{ and } m \text{ same well} \\ 0 & \text{otherwise} \end{cases}$$

Hence, the precision bounds can be simply computed for $N/2$ particles at one of the wells,

$$\frac{1}{(\Delta b_1)^2} \leq 2\sigma^2 \mathcal{F}_Q[|\psi\rangle, J_z^{(N/2)}] \leq \sigma^2 N^2 / 2.$$



If we assume $a = 1$, we have that $H_0 = J_z^{(L)} + J_z^{(R)}$ and $H_1 = J_z^{(L)} - J_z^{(R)}$.

$$\mathcal{F}_Q[\rho, H_0] + \mathcal{F}_Q[\rho, H_1] = 2\mathcal{F}_Q[\rho, J_z^{(L)}] + 2\mathcal{F}_Q[\rho, J_z^{(R)}]$$

Separable states

$$\mathcal{F}_Q[\rho, H_0] + \mathcal{F}_Q[\rho, H_1] = 2N_L + 2N_R = 2N.$$

Heisenberg limit for evenly split systems

$$\mathcal{F}_Q[\rho, H_0] + \mathcal{F}_Q[\rho, H_1] = 2N_L^2 + 2N_R^2 = N^2.$$

Examples

$$|\text{GHZ}\rangle \rightarrow \mathcal{F}_Q[|\psi\rangle, H_0] = N^2 \quad \text{and} \quad \mathcal{F}_Q[|\psi\rangle, H_1] = 0.$$

$$|\psi\rangle = \frac{\overbrace{|0, \dots, 1, \dots\rangle}^{N/2} + \overbrace{|1, \dots, 0, \dots\rangle}^{N/2}}{\sqrt{2}} \rightarrow \mathcal{F}_Q[|\psi\rangle, H_1] = N^2 \quad \text{and} \quad \mathcal{F}_Q[|\psi\rangle, H_0] = 0.$$



Conclusions

- In principle, the **effect of an unknown global rotation** has to be considered.
 - For a **single ensemble** with localized particles, a method with a huge practical advantage, the shot-noise limit can be surpassed if and only if there is a **strong statistical correlation between the particle positions**.
 - There is a **trade-off** between homogeneous and gradient magnetometry if one wants to estimate both parameters at the same time.
-



Conclusions

- In principle, the **effect of an unknown global rotation** has to be considered.
 - For a **single ensemble** with localized particles, a method with a huge practical advantage, the shot-noise limit can be surpassed if and only if there is a **strong statistical correlation between the particle positions**.
 - There is a **trade-off** between homogeneous and gradient magnetometry if one wants to estimate both parameters at the same time.
-

Thank you for your attention!