

# Activating hidden metrological usefulness

Géza Tóth<sup>1,2,3,4</sup>, Tamás Vértesi<sup>5</sup>, Paweł Horodecki<sup>6,7</sup>,  
Ryszard Horodecki<sup>8</sup>

<sup>1</sup>Theoretical Physics, University of the Basque Country (UPV/EHU), Bilbao, Spain

<sup>2</sup>Donostia International Physics Center (DIPC), San Sebastián, Spain

<sup>3</sup>IKERBASQUE, Basque Foundation for Science, Bilbao, Spain

<sup>4</sup>Wigner Research Centre for Physics, Budapest, Hungary

<sup>5</sup>Institute for Nuclear Research, Hungarian Academy of Sciences, Debrecen, Hungary

<sup>6</sup>International Centre for Theory of Quantum Technologies, University of Gdansk, Poland

<sup>7</sup>Faculty of Applied Physics and Mathematics, National Quantum Information Centre,  
Gdansk University of Technology, Gdansk, Poland

<sup>8</sup>Institute of Theoretical Physics and Astrophysics, National Quantum Information Centre,  
Faculty of Mathematics, Physics and Informatics, University of Gdansk, Poland

Torun, Poland,  
8 September 2021 (online).

# Photos



Tamás Vértesi  
Hungary



Paweł Horodecki  
Poland



Ryszard Horodecki  
Poland

# What are entangled states useful for?

- Entanglement is needed for beating the shot-noise limit in quantum metrology.
- Not all entangled states are more useful than separable states.
- Question:
  - Can a quantum state become useful metrologically, if an ancilla or a second copy is added?
  - How to find the local Hamiltonian, for which a quantum state is the most useful compared to separable states?

# Metrological usefulness

- Metrological gain optimized over all local Hamiltonians

$$g(\varrho) = \max_{\text{local } \mathcal{H}} \frac{\mathcal{F}_Q[\varrho, \mathcal{H}]}{\mathcal{F}_Q^{(\text{sep})}(\mathcal{H})} \quad \begin{array}{l} \leftarrow \text{metrological performance of } \varrho \\ \leftarrow \text{best metrological performance of} \\ \text{separable states} \end{array}$$

- A state  $\varrho$  is useful if  $g(\varrho) > 1$ .

G. Toth, T. Vertesi, P. Horodecki, R. Horodecki, PRL 2020.

- We would like to determine  $g$ .

# Method for finding the optimal Hamiltonian II

Maximization of  $\mathcal{F}_Q[\varrho, \mathcal{H}]$  over  $\mathcal{H}$  is difficult: it is convex in  $\mathcal{H}$ .


The maximum over local Hamiltonians can be obtained as

$$\max_{\text{local } \mathcal{H}} \mathcal{F}_Q[\varrho, \mathcal{H}] = \max_{\text{local } \mathcal{H}} \max_M \frac{\langle i[M, \mathcal{H}] \rangle_{\varrho}^2}{(\Delta M)^2}.$$

Similar idea for optimizing over the state, rather than over  $\mathcal{H}$ :

[K. Macieszczak, arXiv:1312.1356; K. Macieszczak, M. Fraas, and R. Demkowicz-Dobrzański, New J. Phys. 16, 113002 (2014); Tóth and Vértesi, Phys. Rev. Lett. (2018).]

# See-saw algorithm



The precision  
cannot get worse  
with the iteration!

Note that  $\mathcal{H}_1, \mathcal{H}_2$  fulfill


$$c_n \mathbb{1} \pm \mathcal{H}_n \geq 0.$$

# Numerical results for isotropic state

- The  $3 \times 3$  isotropic state is useful if

$$p < \frac{25 - \sqrt{177}}{32} \approx 0.3655.$$

- Then, we have the following results for activation.



	Analytic example	Numerics
Ancilla	0.3752	0.3941
Second copy	0.4164	0.4170

# Pure states

- All bipartite pure entangled states are useful.
- In the infinite copy limit, all bipartite pure entangled states are maximally useful.



# Summary

See:

Géza Tóth, Tamás Vértesi, Paweł Horodecki, Ryszard Horodecki,

Activating hidden metrological usefulness,  
[Phys. Rev. Lett. 125, 020402 \(2020\). \(open access\)](#)

THANK YOU FOR YOUR ATTENTION!

