

Entanglement measures (How much is it entangled?) (Lecture of the Quantum Information class of the Master in Quantum Science and Technology)

Géza Tóth

Theoretical Physics, University of the Basque Country (UPV/EHU), Bilbao, Spain
Donostia International Physics Center (DIPC), San Sebastián, Spain
IKERBASQUE, Basque Foundation for Science, Bilbao, Spain
Wigner Research Centre for Physics, Budapest, Hungary

UPV/EHU, Leioa
5 February, 2026

- 1 **Entanglement measures (How much is it entangled?)**
 - Motivation
 - A. General quantum operation
 - B. Local operations and classical communication (LOCC)
 - C. Entanglement of formation
 - D. Concurrence

Entanglement measures

- After detecting entanglement, we have to ask how entangled the state is.
- It will turn out that entanglement is a resource.

- 1 **Entanglement measures (How much is it entangled?)**
 - Motivation
 - A. General quantum operation
 - B. Local operations and classical communication (LOCC)
 - C. Entanglement of formation
 - D. Concurrence

General quantum operation

- The general quantum operation is defined as

$$\varrho' = \sum_k E_k \varrho E_k^\dagger$$

with

$$\sum_k E_k^\dagger E_k = 1.$$

- E_k are Kraus operators.
- Generalized measurements, POVM (positive operator-valued measure).
- Special case: von Neumann measurements, when E_k are pairwise orthogonal projectors.
- Naimark's dilation theorem:
general operation =
von Neumann measurement on system+ancilla.

- 1 **Entanglement measures (How much is it entangled?)**
 - Motivation
 - A. General quantum operation
 - B. Local operations and classical communication (LOCC)
 - C. Entanglement of formation
 - D. Concurrence

Local operations and classical communication (LOCC)

- LOCC are
 - local unitaries,
 - local von Neumann or POVM measurements,
 - local unitaries or measurements conditioned on measurement outcomes on the other party.
- Mathematical description of LOCC. Separable operations are a somewhat larger set, however, this set can easily be described.

$$\varrho' = \sum_k E_k^{(1)} \otimes E_k^{(2)} \varrho \left(E_k^{(1)} \otimes E_k^{(2)} \right)^\dagger$$

with

$$\sum_k \left(E_k^{(1)} \otimes E_k^{(2)} \right)^\dagger \left(E_k^{(1)} \otimes E_k^{(2)} \right) = 1.$$

Local operations and classical communication (LOCC) II

- Stochastic Local Operations and Classical Communication (SLOCC):

$$|\Psi\rangle' \leftarrow E_k^{(1)} \otimes E_k^{(2)} |\Psi\rangle$$

It happens with some probability, not deterministic.

- LOCC cannot create entanglement. Separable states remain separable under LOCC.
- LOCC can create correlations.

- 1 **Entanglement measures (How much is it entangled?)**
 - Motivation
 - A. General quantum operation
 - B. Local operations and classical communication (LOCC)
 - **C. Entanglement of formation**
 - D. Concurrence

Entropy of entanglement

- The von Neumann entropy is defined as

$$S(\varrho) = -\text{Tr}(\varrho \log_2 \varrho).$$

- It can be written with the eigenvalues of the density matrix as

$$S(\varrho) = -\sum_{k=1}^d \lambda_k \log_2 \lambda_k.$$

- For a pure state we have $\lambda_k = \{1, 0, 0, \dots, 0\}$, and thus it is zero.
- Its maximal is for the completely mixed state for which $\lambda_k = \{\frac{1}{d}, \frac{1}{d}, \frac{1}{d}, \dots, \frac{1}{d}\}$, and its value is $\log_2 d$.
- For a bipartite pure state, the **entropy of entanglement** is

$$E_E(|\Psi\rangle) = S(\text{Tr}_1(|\Psi\rangle\langle\Psi|)).$$

That is, it is the von Neumann entropy of the reduced state is an entanglement measure.

Entropy of entanglement II

- Comments
 - It is one for two-qubit singlet states.
 - It is zero for product states.
 - It is invariant under $U_1 \otimes U_2$.

Entanglement of formation

- For mixed states, the entanglement of formation is the convex roof of the von Neumann entropy of the reduced state.

$$E_F = \min_{|\Psi_k\rangle, p_k} \sum_k p_k E_E(|\Psi_k\rangle),$$

- The optimization is over all decompositions of the state of the type

$$\varrho = \sum_k p_k |\Psi_k\rangle\langle\Psi_k|.$$

- E_F tells us, in the asymptotic limit, how many singlets we need to create the state.
- Is it easy to compute? No. For 2×2 systems, there is an explicit formula with the concurrence. For larger systems, there is not a general method.

- 1 **Entanglement measures (How much is it entangled?)**
 - Motivation
 - A. General quantum operation
 - B. Local operations and classical communication (LOCC)
 - C. Entanglement of formation
 - D. Concurrence

Entanglement of formation

- For two qubits, E_F can be calculated explicitly (Wootters, 1997).
- Special case: for pure states the concurrence is

$$C(|\Psi\rangle) = |\langle\Psi|\tilde{\Psi}\rangle| = 2|a_{11}a_{22} - a_{12}a_{21}|,$$

where

$$|\Psi\rangle = \begin{pmatrix} a_{11} \\ a_{12} \\ a_{21} \\ a_{22} \end{pmatrix}.$$

- It is related to the linear entropy of the reduced state.

$$C = \sqrt{2(1 - \text{Tr}(\rho_{\text{red}}^2))}, \quad (1)$$

where

$$\rho_{\text{red}} = \text{Tr}_2(|\Psi\rangle\langle\Psi|). \quad (2)$$

Entanglement of formation II

- Now we have to compute E_F from C .
- We also need that

$$\epsilon(c) = H_2\left(\frac{1 + \sqrt{1 - c^2}}{2}\right).$$

Here

$$H_2 = -x \log_2 x - (1 - x) \log_2(1 - x).$$

- Then, E_F can be obtained as

$$E_F(\varrho) = \epsilon(C(\varrho)).$$

Entanglement of formation III

- For mixed states, the **concurrence** is defined as

$$C(\varrho) = \max(0, \lambda_1 - \lambda_2 - \lambda_3 - \lambda_4),$$

where λ_k 's are, in a decreasing order, the eigenvalues of

$$R = \sqrt{\sqrt{\varrho} \tilde{\varrho} \sqrt{\varrho}},$$

and

$$\tilde{\varrho} = (\sigma_y \otimes \sigma_y) \varrho^* (\sigma_y \otimes \sigma_y).$$