

# Uncertainty relations with the variance and the quantum Fisher information

## — The Cramér-Rao bound as a convex roof

Géza Tóth<sup>1,2,3,4</sup> and Florian Fröwis<sup>5</sup>



<sup>1</sup>Theoretical Physics and EHU Quantum Center,

University of the Basque Country (UPV/EHU), Bilbao, Spain

<sup>2</sup>Donostia International Physics Center (DIPC), San Sebastián, Spain

<sup>3</sup>IKERBASQUE, Basque Foundation for Science, Bilbao, Spain

<sup>4</sup>Wigner Research Centre for Physics, Budapest, Hungary

<sup>5</sup>Group of Applied Physics, University of Geneva, CH-1211 Geneva, Switzerland

Benasque, 19 May 2022

# Outline

## 1 Motivation

- The Cramér-Rao bound

## 2 The derivation

- The Cramér-Rao bound based on a convex roof

# Proving the Cramér-Rao bound

- The Cramér-Rao bound is a fundamental relation in metrology.
- It is an expression with the quantum Fisher information (QFI), which is a complicated function of the state and the Hamiltonian.
- We will look for a simple proof of the Cramér-Rao bound based on fundamental uncertainty relations.
- We will exploit the fact that the QFI is the convex roof of the variance.

# Outline

## 1 Motivation

- The Cramér-Rao bound

## 2 The derivation

- The Cramér-Rao bound based on a convex roof

# Cramér-Rao bound

- Error propagation formula

$$(\Delta\theta)_A^2 = \frac{(\Delta A)^2}{|\partial_\theta \langle A \rangle|^2} = \frac{(\Delta A)^2}{|\langle i[A, B] \rangle|^2}.$$

- The precision of the estimation is bounded as

$$(\Delta\theta)^2 \geq \frac{1}{m} \min_A (\Delta\theta)_A^2,$$

where  $m$  is the number of independent repetitions.

- Let us consider a decomposition of the density matrix

$$\varrho = \sum_k p_k |\psi_k\rangle \langle \psi_k|.$$

- The Heisenberg uncertainty for the components is

$$(\Delta A)_{\psi_k}^2 (\Delta B)_{\psi_k}^2 \geq \frac{1}{4} |\langle i[A, B] \rangle_{\psi_k}|^2.$$

# Cramér-Rao bound II

- Let us consider the inequality

$$\left( \sum_k p_k a_k \right) \left( \sum_k p_k b_k \right) \geq \left( \sum_k p_k \sqrt{a_k b_k} \right)^2,$$

where  $a_k, b_k \geq 0$ .

- Hence, we arrive at

$$\left[ \sum_k p_k (\Delta A)^2_{\psi_k} \right] \left[ \sum_k p_k (\Delta B)^2_{\psi_k} \right] \geq \frac{1}{4} \left[ \sum_k p_k |\langle i[A, B] \rangle_{\psi_k}| \right]^2.$$

- We can choose the decomposition such that

$$\sum_k p_k (\Delta B)^2_{\psi_k} = F_Q[\varrho, B]/4.$$

- Due to the concavity of the variance we also know that

$$\sum_k p_k (\Delta A)^2_{\psi_k} \leq (\Delta A)^2.$$

# Cramér-Rao bound III

- Hence, it follows that

$$(\Delta A)^2_{\varrho} \left[ 4 \min_{\rho_k, \psi_k} \sum_k \rho_k (\Delta B)^2_{\psi_k} \right] \geq |\langle i[A, B] \rangle_{\varrho}|^2.$$

- Then,

$$\frac{(\Delta A)^2_{\varrho}}{|\langle i[A, B] \rangle_{\varrho}|^2} \geq \frac{1}{\left[ 4 \min_{\rho_k, \psi_k} \sum_k \rho_k (\Delta B)^2_{\psi_k} \right]}.$$

- Finally, for the precision of estimation, if we measure  $A$  and the Hamiltonian is  $B$ , we have

$$(\Delta \theta)^2 \geq \frac{1}{m} \min_A (\Delta \theta)^2_A \geq \frac{1}{m} \times \underbrace{\frac{1}{4 \min_{\rho_k, \psi_k} \sum_k \rho_k (\Delta B)^2_{\psi_k}}}_{F_Q[\varrho, B], \text{ the QFI!}}.$$

# Summary

- We showed how to derive the Cramér-Rao bound with the convex roof of the variance.

See:

Géza Tóth and Florian Fröwis,

Uncertainty relations with the variance and the quantum Fisher information based on convex decompositions of density matrices,

[Phys. Rev. Research 4, 013075 \(2022\)](#).

THANK YOU FOR YOUR ATTENTION!