



JYVÄSKYLÄN YLIOPISTO



High precision decay energy measurements of low Q-value beta decays with JYFLTRAP

Zhuang Ge


University of Jyväskylä

Determination of neutrino mass from single β^\pm/EC decay

Current direct neutrino mass probes: Ground-state to ground-state (gs-to-gs) decays
(β^- : Tritium, ^{187}Re ; EC: ^{163}Ho)


- Lower Q-value, higher sensitivity to neutrino mass
- Model independent method

Tritium (β^- -decay)



$$E_0 = Q_0 - E_{\text{rec}} \text{ (recoil corrections: 1.72 eV)}$$

Endpoint energy $E_0 \sim 18.57 \text{ keV}$




Our Purpose: Search for low Q-value decays

$Q \rightarrow 0$, and $Q < 1 \text{ keV}$ (ultra-low)

^{163}Ho (Electron Capture)

$Q_0 \sim 2.83 \text{ keV}$



Situation with gs-to-gs Q-value precision



Ground-to-ground state decays

- Measured with high precision already
- Beta/EC spectrum endpoint measurements already ongoing

1. β -decay of ^3H : Q-value = 18.59201(7) keV

E. G. Myers et al, Phys. Rev. Lett. 114 (2015)

**gs-to-gs decay
Project:**

**KATRIN,
Project8**

recent upper limit of antineutrino mass: 0.8 eV (90% C. L.) from β -decay

2. β -decay of ^{187}Re : Q-value = 2.4709(13) keV

P. Filianin et al., Phys. Rev. Lett. 127, 072502 (2021)

MARE

stopped

3. EC in ^{163}Ho : Q-value = 2.8632(6) keV

Schweiger C., Braß M., Debierre V. et al. Nat. Phys.(2024)

**EChO,
HOLMES**

recent upper limit of neutrino mass: 150 eV (95% C. L.) from EC

Low Q-value decays for neutrino mass determination




We search for low Q-value ground state to nuclear excited state decays.

- Low Q-value (Q^*): < 1 keV

J. Suhonen, Phys. Scr. 89, 054032 (2014)

N. D. Gamage et al., Hyp. Int. 240, 43 (2019)



$$Q^* = Q_{gs} - E^*$$

E^* From gamma spectroscopy

- Typical uncertainty ~ 100 eV
- Potentially ~ 10 eV

1. β -decay of $^{115}\text{In}(9/2^+) \rightarrow ^{115}\text{Sn}^*(9/2^+)$: Q^* -value = 0.147(10) keV

E^* improvement: *V. A. Zheltonozhsky et al. 2018 EPL 121 12001*

2. β -decay of $^{135}\text{Cs}(7/2^-) \rightarrow ^{135}\text{Ba}^*(11/2^+)$: Q^* -value = 0.44(31) keV

Q_{gs} improvement: *A. De Roubin, J. Kostensalo, T. Eronen et al., Phys. Rev. Lett., 124 (22), 222503.*

2. EC of $^{75}\text{Se}(5/2^+) \rightarrow ^{75}\text{As}^*({3/2^-, 5/2^-})$: Q^* -value = 0.64(51) keV

Q_{gs} improvement: *M. Ramalho, Z. Ge, T. Eronen et al., Phys. Rev. C 106, 015501 (2022)*

Our work: Q_{gs} measurements

- Penning trap mass spectrometry (JYFLTRAP)
- Q_{gs} through $E = mc^2$

Nuclear theory:

- Partial half-life based on Q^*



Summary of measured Q-values of potential candidates at JYFLTRAP



List of measured promising low Q-value decay candidates for neutrino mass determination

	Parent	T1/2	Daughter	E* (keV)	decay type	Q* (keV)	Decay	Q ₀ (keV)	dQ ₀ (keV)
	146Pm(3-)	5.53(5) y	146Nd(2+)	1470.63(6)	1st FNU	1.3(4.2)	EC	1472.000	4.000
Z. Ge T. Eronen, et al., Phys. Rev. C 00, 005500 (2022)	149Gd(7/2-)	9.28(10) dy	149Eu(5/2+)	1312(4)	1st FNU	2(6.4)	EC	1314.100	4.000
	155Tb(3/2+)	5.32(6) dy	155Gd{3/2+}	815.731(3)	Allowed{?}	4.2(10.1)	EC	820.000	10.000
	159Dy(3/2-)	144.4(2) dy	159Tb(5/2-)	363.5449(14)	Allowed	1.7(1.2)	EC	365.200	1.200
Z. Ge, T. Eronen, et al., Phys. Rev. Lett. 127, 272301 (2021)			159Tb(11/2+)	362.050(40)	3rd FU	3.2(1.2)	EC	365.200	1.200
	161Ho(5/2-)	18.479(4) hr	161Dy{7/2+}	858.502(7)	1st FNU	1.0(2.2)	EC	858.500	2.200
			161Dy{3/2-}	858.7919(18)	Allowed	-0.3(2.2)	EC	858.500	2.200
	72As(2-)	26.0(1)h	72Ge{1}	4358.7(3)	Allowed{?}	-2.8(4.0)	EC	4356.000	4.000
Z. Ge, T. Eronen et al., PHYSICAL REVIEW C 103, 065502 (2021)			72Ge(3-)	3325.01(3)	Allowed	8.9(4.0)	β+	4356.000	4.000
			72Ge(2+)	3327(3)	1st FNU	6.9(5.0)	β+	4356.000	4.000
			72Ge{1+}	3338.0(3)	1st FNU{?}	-4.1(4.0)	β+	4356.000	4.000
			72Ge{2-}	3341.76(4)	Allowed{?}	-7.9(4.0)	β+	4356.000	4.000
	159Gd(3/2-)	26.24(9) h	159Tb{1/2+}	971	1st FNU{?}	0.0(1.8)	β-	970.900	0.800
Z. Ge T. Eronen, et al., EPJA (2024)	77As(3/2-)	38.79(5) h	77Se(5/2+)	680.1035(17)	1st FNU	3.1(1.7)	β-	683.200	1.700
	76As(2-)	26.24(9) h	76Se{2-}	2968.4(7)	Allowed{?}	-7.8(1.1)	β-	2960.600	0.900
	153Tb(5/2+)	2.34(1)dy	153Gd(5/2-)	548.7645(18)	1st FNU	-1.2(4.0)	β+	1569.000	4.000
			153Gd{5/2}	551.092(19)	Allowed{?}	-3.5(4.0)	β+	1569.000	4.000
Z. Ge, T. Eronen, et al., Physics Letters B 832 (2022) 137226	111In(9/2+)	3dy	111Cd(3/2+)	864.8(3)	2nd FU	-6.6(3.0)	EC	860.2	3.4
			111Cd(3/2+)	864.8(3)	2nd FU	-4.6(3.0)	EC	860.2	3.4
			111Cd(3/2+)	855.6(1.0)	2nd FU	4.6(3.2)	EC	860.2	3.4
T. Eronen, Z. Ge et al., Physics Letters B 830 (2022) 137135			111Cd(7/2+)	853.94(7)	Allowed	6.3(3.0)	EC	860.2	3.4
	131I(7/2+)	8dy	131Xe{9/2+}	971.22(13)	Allowed{?}	-0.42(0.61)	β-	970.80	0.60
			131Xe(7/2+)	973.11(14)	Allowed	-2.31(0.62)	β-	970.80	0.60
	155Eu(5/2+)	5yr	155Gd(9/2-)	251.7056(10)	1st FU	0.1(1.8)	β-	252.00	2.40

Z. Ge T. Eronen, et al., Phys. Rev. C 108, 045502 (2023)

Q₀ from: M. Wang et al. , Chinese Physics C 45, 030003 (2021)


E* from: National nuclear data center, Available at <https://www.nndc.bnl.gov>

One more long table list will not be shown in this talk

The Ion Guide Isotope Separator On-Line facility (IGISOL)



J. Ärje, J. Äystö et al., PRL 54 (1985) 99





T. Eronen et al., Eur. Phys. J. A 48 (2012) 46

JYFLTRAP double Penning trap



JYFLTRAP double Penning trap



Eronen et al., EPJA 48 (2012) 46

Cyclotron frequency

$$\nu_c = \frac{1}{2\pi} \frac{q}{m} B$$

Penning trap eigenfrequencies:

$$\nu_z = \frac{1}{2\pi} \sqrt{\frac{U_0}{d^2} \frac{q}{m}}$$


$$\nu_{\pm} = \frac{1}{2} \left(\nu_c \pm \sqrt{\nu_c^2 - 2\nu_z^2} \right)$$

Invariance theorem

$$\nu_c^2 = \nu_-^2 + \nu_+^2 + \nu_z^2$$

TOF – ICR and PI – ICR:



$$\nu_c = \nu_+ + \nu_-$$





TOF-ICR method



one-pulse radio-frequency (rf) field (400 ms)




two-pulse rf field (25-350-25 ms)



Interleaved measurements of Ion-of-interest and reference (^{111}In and ^{111}Cd)

Phase-imaging Ion-Cyclotron-Resonance (PI-ICR)

Delay-Line Microchannel Channel Plate (MCP) Detector from Roentdek GmbH




Position sensitive detector

Angle between cyclotron and magnetron motion phases with respect to the center spot:

$$\alpha_c = \alpha_- + \alpha_+$$

cyclotron frequency:



$$\nu_c = \nu_+ + \nu_- = \frac{\alpha_c + 2\pi n}{2\pi t}$$



S. Eliseev et al., Phys. Rev. Lett. 110, 082501 (2013).


D. A. Nesterenko, et al., European Physical Journal A 54, 0 (2018).

Phase-imaging Ion-Cyclotron-Resonance (PI-ICR)



- / Dipolar excitation at ν_+
- Direct conversion to magnetron motion
→ Magnetron phase
 - Accumulation then conversion to magnetron motion
→ Modified cyclotron phase

Schematic of PI-ICR for ^{159}Dy - ^{159}Tb Q-value measurements




Typical systematic uncertainties

D.A. Nesterenko, T. Eronen, Z. Ge, A. Kankainen, M. Vilen, *Eur. Phys. J. A* (2021) 57:302

• JYFLTRAP (TOF-ICR)


- 7.8×10^{-10} per u, residual: 1.2×10^{-8}
- If $m - m_{ref} < 24u$:
 - 7.5×10^{-10} per u, residual: 7.9×10^{-9}





magnetic field fluctuations:
 $\delta B/B = 2.01(25) \times 10^{-12} / \text{min } \delta t$

• JYFLTRAP (PI-ICR)

- $-2.35(81) \times 10^{-10}$ per u, residual: 9×10^{-9}
- If $m - m_{ref} < 12u$:
 - $-2.3(21) \times 10^{-10}$ per u, residual: 5.3×10^{-9}






To reach the precision at the level of 10^{-10} or better, need:
 $|m_{ref} - m_{ioi}| \leq 2 u$ or ideally A/q doublets.



Cleaning methods coupled with PI-ICR method



Z. Ge, T. Eronen, A. de Roubin et al., Phys. Rev. C **108**, 045502 (2023)




Z. Ge, M. Reponen, T. Eronen et al., arXiv:2401.07976v2 (PRL, accepted)

Conatminant-free ion sample preparation:

Coupling of Ramsey cleaning&Buffer gas cleaning&laser frequency scan and PI-ICR method for unambiguous cleaning contaminants of 90 keV ($A=136$) away from ion of interest easy to clean, more than 10^6 resolving power to clean 2 or more closely lying contaminants

Q-value measurement of ^{159}Dy




Gs-to-GS Q value ($Q_{\text{EC}}^{\text{gs}}$)

Obtained frequency ratio r with a precision of 1.3×10^{-9}



Q-value precision: **190 eV**
now **6.3 times** more precise
and 0.47 keV smaller than
literature value

Level scheme of ^{159}Dy with refined Q-value



GS-to-GS Q value (Q_{EC}^{gs})

E_i^*	Binding energy e_x (allowed atomic shell x of the EC)
5/2- 363.5449(14)	+
11/2+ 362.050(40)	




With the refined Q_{EC}^{gs} :

- Captures to 5/2-
 - only from N1 or higher orbitals
 - M2 and M1 captures forbidden at $> 4\sigma$ level
- captures to 11/2+
 - from M1 and higher orbitals

Z. Ge, T. Eronen, K. S. Tyrin et al., Phys. Rev. Lett. 127, 272301(2021)

M. Wang et al., Chinese Physics C 45, 030003 (2021) -> AME2020
National nuclear data center, Available at <https://www.nndc.bnl.gov>

EC spectrum of ^{159}Dy ($3/2^- \rightarrow 5/2^-$) compared to ^{163}Ho
(Dirac-Hartree-Fock atomic many-body calculations)



$^{159}\text{Dy}(3/2^-) \rightarrow ^{159}\text{Tb}^*(5/2^-), Q_{\text{EC}}^* = 1.18(19) \text{ keV}$

⇒ lower than the GS-to-GS Q_{EC} of ^{163}Ho (running)


- allowed transition
 - known branching ratio $1.9(5) \times 10^{-6}$
 - ultra-low distance to the atomic line M1: 0.79(19) keV
- ⇒ the most promising gs-to-excited state transition for future calorimetric experiment

Z. Ge, T. Eronen, K. S. Tyrin et al., Phys. Rev. Lett. 127, 272301(2021)

Puzzles in potential candidates ^{72}As , ^{111}In




State, i	E^* (keV)	J^π	Decay type	Q (keV)
1	3325.01(3)	3^-	β^+ : Allowed	8.9(40)
2	3327(3)	2^+	β^+ : 1st FNU	6.9(50)
3	3338.0(3)	1^{+}	β^+ : 1st FNU{?}	-4.1(40)
4	3341.76(4)	$\{2\}^-$	β^+ : Allowed{?}	-7.9(40)
5	4358.7(3)	1	EC: Allowed{?}	-2.8(40)
gs	0	0^+		4356(4)



Z. Ge, T. Eronen et al., *PHYSICAL REVIEW C* 103, 065502 (2021)

similarly




Z. Ge, T. Eronen, et al., *Physics Letters B* 832 (2022) 137226


Q_0 from M. Wang et al. , *Chinese Physics C* 45, 030003 (2021)

E^* from National nuclear data center, Available at <https://www.nndc.bnl.gov>

Q-value of potential candidate ^{72}As , ^{111}In



Large discrepancy from the AME2020




Good agreement with the AME2020

GS-to-GS Q-values to $\sim 100 \text{ eV}$ precision to determine whether the cases are suitable for neutrino mass determination

Required to be measured directly with high accuracy and precision

Ruling out ^{72}As as potential candidate



Q -values (in keV) for the decay candidate to the excited states of the daughter nucleus ^{72}Ge

E^*	Q -value (AME2020)	Q -value (This work)	$Q/\delta Q$ (This work)
3325.01(3)	8.9(40)	-3.42(8)	43
3327(3)	6.9(50)	-5.4(30)	1.8
3338.0(3)	-4.1(40)	-16.41(31)	53
3341.76(4)	-7.9(40)	-20.17(8)	238
4358.7(3)	-2.8(40)	-15.11(31)	49

five potential ultra-low Q -value β^+ -decay or electron capture transitions are energetically forbidden, precluding all the transitions as possible candidates for the electron neutrino mass determination




However, the discovery of *small negative Q -values* opens up the possibility to use ^{72}As for the study of virtual β - γ transitions

Potential cases of ^{111}In , ^{131}I and ^{95}Tc



Normalized distribution of the released energy (EC) and partial half-lives:
atomic self-consistent many-electron Dirac–Hartree–Fock–Slater method and NSM





- ✓ Neutrino mass, Neutrino capture, relic ne

- ⁹⁵Tc --- submitted (PLB)

- ❖ More than 10 other Cases to be characterized for

- The JYFLTRAP wanna some rest now --- 0-T s

- ❖ More low Q value cases to be measured

- 14 days of beam time with Penning trap is left

- **More ...**

- ❖ To do: N~Z mass measurements (recently Funded Academy project)


- Make JYFLTRAP great again (PRL accepted: masses 95-97Ag, Trap+Laser+Hot cavity+HIGISOL/M)

- **IGISOL MR-TOF (RIKEN MR-TOF?)**

- **RIKEN storage ring + Bro-TOF**

- ✓ (9 days of beam time approved: 100Sn) x3 to measure all N=Z nuclei

- **RIKEN storage ring + Bro-TÖF (9 days of beam time) x3 to measure all N=Z**



Let us see how productive for this project in next PLATAN



Theory
experiment

COLLABORATION LIST



T. Eronen,¹ Z. Ge,¹ A. de Roubin,² D. A. Nesterenko,¹ M. Hukkanen,^{1, 2} O. Beliuskina,¹ R. de Groote,¹ C. Delafosse,¹ S. Geldhof,¹ W. Gins,¹ A. Kankainen,¹ Á. Koszorús,⁷ I. D. Moore,¹ H. Penttilä,¹ A. Raggio,¹ S. Rinta-Antila,¹ M. Stryczyk,¹ V. Virtanen,¹ A. P. Weaver,⁸ A. Zadvornaya,¹ A. Jokinen¹ and the IGISOL collaboration

J. Suhonen,¹ K. S. Tyrin,³ M. I. Krivoruchenko,^{3,4} J. Kostensalo,¹ J. Kotila,^{5,6} P. Pirinen,¹ M. Ramalho¹

¹ Department of Physics, University of Jyväskylä, Finland

² Centre d'Etudes Nucléaires de Bordeaux Gradignan, UMR 5797 CNRS/IN2P3 - Université de Bordeaux, France

³ National Research Centre "Kurchatov Institute", Russia

⁴ Institute for Theoretical and Experimental Physics, NRC "Kurchatov Institute", Russia

⁵ Finnish Institute for Educational Research, University of Jyväskylä, Finland

⁶ Yale University, USA

⁷ University of Liverpool, United Kingdom

⁸ University of Brighton, United Kingdom

Theoretical Collaborators before 2021

Financial Support

Academy of Finland projects

No. 306980, 312544, 275389, 284516, 295207, 314733, 315179, 327629, 320062, 354589, 345869 and 354968.

EU Horizon 2020 research and innovation program

under grant

No. 771036 (ERC CoG MAIDEN)

No. 861198–LISA–H2020-MSCA-ITN-2019

Zhuang Ge^{a,*}, Tommi Eronen^a, Vasile Alin Sevestrean^{b,c,d,*}, Ovidiu Nițescu^{b,d}, Sabin Stoica^b, Marlom Ramalho^a, Jouni Suhonen^{a,b,*}, Antoine de Roubin^{e,f}, Dmitrii Nesterenko^a, Anu Kankainen^a, Pauline Ascher^f, Samuel Ayet San Andres^g, Olga Beliuskina^a, Pierre Delahaye^h, Mathieu Flayol^f, Mathias Gerbaux^f, Stéphane Grévy^f, Marjut Hukkanen^{a,i}, Arthur Jaries^a, Ari Jokinen^a, Audric Husson^f, Daid Kahl^{j,2}, Joel Kostensalo^k, Jenni Kotila^{b,l,m}, Iain Moore^a, Stylianos Nikas^a, Marek Stryczyk^a and Ville Virtanen^a

New theoretical collaborators

^aDepartment of Physics, University of Jyväskylä, P.O. Box 35, FI-40014, Jyväskylä, Finland

^bInternational Centre for Advanced Training and Research in Physics (CIFRA), POB MG-12, RO-077125, Bucharest-Măgurele, Romania

^cFaculty of Physics, University of Bucharest, 405 Atomistilor, POB MG-11, RO-077125, Bucharest-Măgurele, Romania

^d"Horia Hulubei" National Institute of Physics and Nuclear Engineering, 30 Reactorului, POB MG-6, RO-077125, Bucharest-Măgurele, Romania

^eKU Leuven, Instituut voor Kern- en Stralingsfysica, B-3001, Leuven, Belgium

^fUniversité de Bordeaux, CNRS/IN2P3, UMR 5797, F-33170, Gradignan, France

^gInstituto de Física Corpuscular, CSIC-UV, 46980, Gradignan, Spain

^hGANIL, CEA/DSM-CNRS/IN2P3, Bd Henri Becquerel, 14000, Caen, France

ⁱUniversité de Bordeaux, CNRS/IN2P3, LP2I Bordeaux, UMR 5797, F-33170, Gradignan, France

^jExtreme Light Infrastructure - Nuclear Physics, Horia Hulubei National Institute for R&D in Physics and Nuclear Engineering (IFIN-HH), 077125, Bucharest-Măgurele, Romania

^kNatural Resources Institute Finland, Yliopistokatu 6B, FI-80100, Joensuu, Finland

^lFinnish Institute for Educational Research, University of Jyväskylä, P.O. Box 35, FI-40014, Jyväskylä, Finland

^mCenter for Theoretical Physics, Sloane Physics Laboratory, Yale University, New Haven, Connecticut 06520-8120, Connecticut, USA



*Thank you
for
your attention*



Zhuang Ge---High precision decay energy measurements of low Q-value beta decays with JYFLTRAP at IGISOL

*Thank you for your
attention*