




JYVÄSKYLÄN YLIOPISTO
UNIVERSITY OF JYVÄSKYLÄ

Deutsche Physikalische Gesellschaft

Φ DPG



Search for low Q value beta decays for neutrino mass determination

Zhuang Ge

31-03-2022

GSI Helmholtzzentrum für Schwerionenforschung GmbH,
University of Jyväskylä


Determination of neutrino mass from single β^\pm /EC decay

Φ DPG

Current direct neutrino mass probes: Ground-state to ground-state (gs-to-gs) decays
 (β^- : Tritium, ^{187}Re ; EC: ^{163}Ho)


- Lower Q-value, higher sensitivity to neutrino mass
- Model independent method

Tritium (β^- -decay)




$$E_0 = Q_0 - E_{\text{rec}} \quad (\text{recoil corrections: } 1.72 \text{ eV})$$

Endpoint energy $E_0 \sim 18.57 \text{ keV}$



Our Purpose: Search for low Q-value decays
 $Q \rightarrow 0, \text{ and } Q < 1 \text{ keV (ultra-low)}$


^{163}Ho (Electron Capture)



Low Q-value decays for neutrino mass determination

We search for low Q-value ground state to nuclear excited state decays.

- Low Q-value (Q^*): < 1 keV



1. β -decay of $^{115}\text{In}(9/2^+) \rightarrow ^{115}\text{Sn}^*(9/2^+)$: Q^* -value = 0.147(10) keV
 E^* improvement: V.A. Zheltonozhsky et al. 2018 EPL 121 12001

J. Suhonen, Phys. Scr. 89, 054032 (2014)
N. D. Gamage et al., Hyp. Int. 240, 43 (2019)

$$Q^* = Q_{\text{gs}} - E^*$$

E^* From gamma spectroscopy

- Typical uncertainty ~100 eV
- Potentially ~10 eV

Our work: Q_{gs} measurements

- Penning trap mass spectrometry (JYFLTRAP)
- Q_{gs} through $E = mc^2$

Nuclear theory:

- Partial half-life based on Q^*

Summary of measured Q-values of potential candidates



- List of measured promising low Q-value decay candidates for neutrino mass determination

Parent	T1/2	Daughter	E* (keV)	decay type	Q* (keV)	Decay	Q ₀ (keV)	dQ ₀ (keV)
146Pm(3-)	5.53(5) y	146Nd(2+)	1470.63(6)	1st FNU	1.3(4.2)	EC	1472.000	4.000
149Gd(7/2-)	9.28(10) dy	149Eu(5/2+)	1312(4)	1st FNU	2(6.4)	EC	1314.100	4.000
155Tb(3/2+)	5.32(6) dy	155Gd{3/2+}	815.731(3)	Allowed{?}	4.2(10.1)	EC	820.000	10.000
159Dy(3/2-)	144.4(2) dy	159Tb(5/2-)	363.5449(14)	Allowed	1.7(1.2)	EC	365.200	1.200
<i>Z. Ge, T. Eronen et al., Phys. Rev. Lett. 127, 272301</i>		159Tb(11/2+)	362.050(40)	3rd FU	3.2(1.2)	EC	365.200	1.200
161Ho(5/2-)	18.479(4) hr	161Dy{7/2+}	858.502(7)	1st FNU	1.0(2.2)	EC	858.500	2.200
		161Dy{3/2-}	858.7919(18)	Allowed	-0.3(2.2)	EC	858.500	2.200
72As(2-)	26.0(1)h	72Ge{1}	4358.7(3)	Allowed{?}	-2.8(4.0)	EC	4356.000	4.000
<i>Z. Ge, T. Eronen et al., PHYSICAL REVIEW C 103, 065502 (2021)</i>		72Ge(3-)	3325.01(3)	Allowed	8.9(4.0)	β^+	4356.000	4.000
		72Ge(2+)	3327(3)	1st FNU	6.9(5.0)	β^+	4356.000	4.000
		72Ge{1+}	3338.0(3)	1st FNU{?}	-4.1(4.0)	β^+	4356.000	4.000
		72Ge{2-}	3341.76(4)	Allowed{?}	-7.9(4.0)	β^+	4356.000	4.000
159Gd(3/2-)	26.24(9) h	159Tb{1/2+}	971	1st FNU{?}	0.0(1.8)	β^-	970.900	0.800
77As(3/2-)	38.79(5) h	77Se(5/2+)	680.1035(17)	1st FNU	3.1(1.7)	β^-	683.200	1.700
76As(2-)	26.24(9) h	76Se{2-}	2968.4(7)	Allowed{?}	-7.8(1.1)	β^-	2960.600	0.900
153Tb(5/2+)	2.34(1)dy	153Gd(5/2-)	548.7645(18)	1st FNU	-1.2(4.0)	β^+	1569.000	4.000
		153Gd{5/2}	551.092(19)	Allowed{?}	-3.5(4.0)	β^+	1569.000	4.000
111In(9/2+)	3dy	111Cd(3/2+)	864.8(3)	2nd FU	-6.6(3.0)	EC	860.2	3.4
<i>Submitted</i>		111Cd(3/2+)	864.8(3)	2nd FU	-4.6(3.0)	EC	860.2	3.4
		111Cd(3/2+)	855.6(1.0)	2nd FU	4.6(3.2)	EC	860.2	3.4
		111Cd(7/2+)	853.94(7)	Allowed	6.3(3.0)	EC	860.2	3.4
131I(7/2+)	8dy	131Xe{9/2+}	971.22(13)	Allowed{?}	-0.42(0.61)	β^-	970.80	0.60
		131Xe(7/2+)	973.11(14)	Allowed	-2.31(0.62)	β^-	970.80	0.60
155Eu(5/2+)	5yr	155Gd(9/2-)	251.7056(10)	1st FU	0.1(1.8)	β^-	252.00	2.40


Q_0 from: M. Wang et al. , Chinese Physics C 45, 030003 (2021)

E^* from: National nuclear data center, Available at <https://www.nndc.bnl.gov>

The Ion Guide Isotope Separator On-Line facility (**IGISOL**)



J. Ärje, J. Äystö et al., PRL 54 (1985) 99




T. Eronen et al., Eur. Phys. J. A 48 (2012) 46

JYFLTRAP double Penning trap

Φ DPG

JYFLTRAP double Penning trap



Eronen et al., EPJA 48 (2012) 46

Cyclotron frequency

$$\nu_c = \frac{1}{2\pi} \frac{q}{m} B$$


Penning trap eigenfrequencies:

$$\nu_z = \frac{1}{2\pi} \sqrt{\frac{U_0}{d^2} \frac{q}{m}}$$

$$\nu_{\pm} = \frac{1}{2} \left(\nu_c \pm \sqrt{\nu_c^2 - 2\nu_z^2} \right)$$

Invariance theorem

$$\nu_c^2 = \nu_-^2 + \nu_+^2 + \nu_z^2$$



Q-value and mass measurements

- Cyclotron frequency:

$$\nu_c = \nu_+ + \nu_- = \frac{qB}{2\pi m}$$

- Frequency ratio r :

$$r = \frac{\nu_1}{\nu_2}$$

Determine

- Q -value:
$$Q = M_2 - M_1 = (r - 1)(M_1 - m_e) + m_e$$

- Mass:
$$M_2 = r(M_1 - m_e) + m_e$$

Eronen et al., EPJA 48 (2012) 46

1. TOF-ICR

Time-of-Flight Ion-Cyclotron-Resonance (TOF-ICR) technique
G. Gräff et al., Zeitschrift für Physik A: Atoms and Nuclei 297, 35 (1980).

- a. Normal TOF-ICR
- b. Ramsey TOF-ICR

2. PI-ICR


Phase-imaging Ion-Cyclotron-Resonance (PI-ICR) technique
S. Eliseev et al., Phys. Rev. Lett. 110, 082501 (2013).



Phase-imaging Ion-Cyclotron-Resonance (PI-ICR)


Φ DPG

Delay-Line Microchannel Channel Plate (MCP) Detector from Roentdek GmbH



S. Eliseev et al., Phys. Rev. Lett. 110, 082501 (2013).

D. A. Nesterenko, et al., European Physical Journal A 54, 0 (2018).




Angle between cyclotron and magnetron motion phases with respect to the center spot:

$$\alpha_c = \alpha_- + \alpha_+$$

cyclotron frequency:


$$\nu_c = \nu_+ + \nu_- \\ = \frac{\alpha_c + 2\pi n}{2\pi t}$$

Schematic of PI-ICR for $^{159}\text{Dy}-^{159}\text{Tb}$ Q-value measurement.



Imaging

Q-value measurement of ^{159}Dy



Gs-to-GS Q value ($Q_{\text{EC}}^{\text{gs}}$)



Obtained frequency ratio r with a precision of 1.3×10^{-9}



Q-value precision: 190 eV
now 6.3 times more precise
and 0.47 keV smaller than
literature value

Level scheme of ^{159}Dy with refined Q-value

Φ DPG



With the refined $Q_{\text{EC}}^{\text{gs}}$:

- Captures to 5/2⁻**
 - only from N1 or higher orbitals
 - M2 and M1 captures forbidden at $> 4\sigma$ level
- Captures to 11/2⁺**
 - from M1 and higher orbitals

Z. Ge, T. Eronen, K. S. Tyrin et al., Phys. Rev. Lett. 127, 272301(2021)

M. Wang et al. , Chinese Physics C 45, 030003 (2021) -> AME2020
National nuclear data center; Available at <https://www.nndc.bnl.gov>



GSI

JYU. Since 1863.

2022/3/11


13

Results and conclusion of ^{159}Dy



EC spectrum of ^{159}Dy ($3/2^- \rightarrow 5/2^-$) compared to ^{163}Ho
(Dirac-Hartree-Fock atomic many-body calculations)


Z. Ge, T. Eronen, K. S. Tyrin et al., Phys. Rev. Lett. 127, 272301(2021)



$Q_{EC}^* = 1.18(19) \text{ keV} \rightarrow$ the lowest EC Q value for the transition:
 $^{159}\text{Dy}(3/2^-) \rightarrow ^{159}\text{Tb}^*(5/2^-)$,
lower than the GS-to-GS Q_{EC} of ^{163}Ho ,
utilized in presently running or planned direct neutrino mass experiments

- allowed transition
- known branching ratio $1.9(5) \times 10^{-6}$
- Smallest EC Q value
→ the most promising gs-to-excited state transition
for future calorimetric experiment

More cases have been measured and in analysis/preparation



Collaboration list



T. Eronen,¹ Z. Ge,¹ A. de Roubin,² D. A. Nesterenko,¹ M. Hukkanen,^{1, 2} O. Beliuskina,¹ R. de Groot,¹ C. Delafosse,¹ S. Geldhof,¹, W. Gins,¹ A. Kankainen,¹ Á. Koszorús,⁷ I. D. Moore,¹ H. Penttilä,¹ A. Raggio,¹ S. Rinta-Antila,¹ M. Stryjczyk,¹ V. Virtanen,¹ A. P. Weaver,⁸ A. Zadvornaya,¹ A. Jokinen¹, T. Dickel^{9, 10}, W. R. Plaß^{9, 10} and the IGISOL collaboration

K. S. Tyrin,³ M. I. Krivoruchenko,^{3, 4} J. Kostensalo,¹ J. Kotila,^{5, 6} J. Suhonen,¹

¹ Department of Physics, University of Jyväskylä, P.O. Box 35, FI-40014, Jyväskylä, Finland

² Centre d'Etudes Nucléaires de Bordeaux Gradignan, UMR 5797 CNRS/IN2P3 - Université de Bordeaux,
19 Chemin du Solarium, CS 10120, F-33175 Gradignan Cedex, France

³ National Research Centre "Kurchatov Institute", Ploschad' Akademika Kurchatova 1, 123182 Moscow, Russia

⁴ Institute for Theoretical and Experimental Physics, NRC "Kurchatov Institute", B. Cheremushkinskaya 25, 117218 Moscow, Russia

⁵ Finnish Institute for Educational Research, University of Jyväskylä, P.O. Box 35, FI-40014, Jyväskylä, Finland

⁶ Center for Theoretical Physics, Sloane Physics Laboratory Yale University, New Haven, Connecticut 06520-8120, USA

⁷ Department of Physics, University of Liverpool, Liverpool, L69 7ZE, United Kingdom

⁸ School of Computing, Engineering and Mathematics, University of Brighton, Brighton BN2 4JG, United Kingdom

⁹ GSI Helmholtzzentrum für Schwerionenforschung GmbH, Darmstadt, Germany

¹⁰ Justus-Liebig University of Giessen, Germany

Financial Support

Academy of Finland projects No. 306980, 312544, 275389, 284516, 295207, 314733 and 320062.
EU Horizon 2020 research and innovation program under grant No. 771036 (ERC CoG MAIDEN)

Acknowledgements

the staff of the accelerator laboratory of University of Jyväskylä (JYFL-ACCLAB)
J. Jaatinen and R. Seppälä, for preparing the production target



Thank you for your attention