

## (PTIA0301) Elementary Linear Algebra

#### Dr. Gabor FACSKO, PhD

Senior Research Fellow facskog@gamma.ttk.pte.hu

University of Pecs, Faculty of Scienceüs, Institute of Mathematics and Informatics, 7624 Pecs, Ifjusag utja 6.
Wigner Research Centre for Physics, Department of Space Physics and Space Technology, 1121 Budapest, Konkoly-Thege Miklos ut 29-33.
https://facs.ko.ttk.pte.i/

September 6, 2024

### Multiply vector with a scalar and sum of vectors

**Definition:** Sum of vectors. If  $\mathbf{a}(a_1, a_2, a_3)$  and  $\mathbf{b}(b_1, b_2, b_3)$ , then

$$\mathbf{a} + \mathbf{b} = (a_1 + b_1, a_2 + b_2, a_3 + b_3),$$

where  $a_1$ ,  $a_2$ ,  $a_3$ ,,  $b_1$ ,  $b_2$ ,  $b_3 \in \mathbb{R}$ .

**Definition:** Subtracktion of vectors. If  $\mathbf{a}(a_1, a_2, a_3)$  and  $\mathbf{b}(b_1, b_2, b_3)$ , then

$$\mathbf{a} - \mathbf{b} = (a_1 - b_1, a_2 - b_2, a_3 - b_3),$$

where  $a_1$ ,  $a_2$ ,  $a_3$ ,  $b_1$ ,  $b_2$ ,  $b_3 \in \mathbb{R}$ .

#### Multiply vector with a scalar, difference of vectors

▶ Definition: Multiplication of vectors with a scalar. If  $\lambda \in \mathbb{R}$  and  $\mathbf{a}(a_1, a_2, a_3)$ , where  $a_1$ ,  $a_2$ ,  $a_3 \in \mathbb{R}$ , then

$$\lambda \mathbf{a} = (\lambda a_1, \lambda a_2, \lambda a_3).$$

Mind if  $\lambda$  is 0. 1. -1. <1. or >1.

- The sum of vectors is
  - ightharpoonup Commutative:  $\mathbf{a} + \mathbf{b} = \mathbf{b} + \mathbf{a}$ , where  $\mathbf{a}, \mathbf{b} \in \mathbb{R}^3$ .

  - Associative: (a + b) + c = a + (b + c), where a, b, c ∈ R³.
    The null vector exists: ∃0 ∈ R³, where a + 0 = a, where a ∈ R³.
    All vectors have an inverse vector: ∀a ∈ R³ ∃ (-a) ∈ R³, where a + (-a) = 0.
- ► The vector multiplication with scalar is

  - Associative:  $\lambda(\mu \mathbf{a}) = (\lambda \mu) \mathbf{a}$ , where  $\mathbf{a} \in \mathbb{R}^3, \lambda, \mu \in \mathbb{R}$ .
    Distributive:  $\lambda(\mathbf{a} + \mathbf{b}) = \lambda \mathbf{a} + \lambda \mathbf{b}$ , where  $\mathbf{a}, \mathbf{b} \in \mathbb{R}^3, \lambda \in \mathbb{R}$ .
  - Distributive:  $(\lambda + \mu) \mathbf{a} = \lambda \mathbf{a} + \mu \mathbf{a}$ , where  $\mathbf{a} \in \mathbb{R}^3$ ,  $\lambda, \mu \in \mathbb{R}$ .  $\forall \in \mathbf{a} \cdot \mathbf{1} = \mathbf{a}$ , where  $\mathbf{a} \in \mathbb{R}^3$ .

# The End

Thank you for your attention!