



(ENKEMNA0302) Applied Linear Algebra

Dr. Gabor FACSKO, PhD

Senior Research Fellow

facskog@gamma.ttk.pte.hu

University of Pecs, Faculty of Sciences, Institute of Mathematics and Informatics, 7624 Pecs, Ifjusag utja 6.
Wigner Research Centre for Physics, Department of Space Physics and Space Technology, 1121 Budapest, Konkoly-Thege Miklos ut 29-33.
<https://facsko.ttk.pte.hu>

February 27, 2025

Airy disc I

- ▶ During most of the practice, I finished the material from the previous day's lecture.
- ▶ In the last few minutes, we dealt with the Airy disc. The light of stars is formed on the CCD camera according to the formula $\frac{\sin(r)}{r}$, where r is the distance from the centre of the object. Rings appeared around the central peak. The goal is to clean the image from these rings.
- ▶ We determined that the height of the first ring is 0.13 if the central peak is 1.0. Thus, this needs to be removed.
- ▶ The matrix $\mathbf{A}_{m \times n}$ represents a $m \times n$ greyscale image.
- ▶ Each matrix element represents a color in the interval $\{0, \dots, 1\}$, where 0 is black and 1 is white.
- ▶ The matrix $\mathbf{B}_{m \times n}$ represents the background to be removed, with a value of 0.13.
- ▶ We are looking for the operation $\mathbf{A} \odot \mathbf{B} = [a_{ij} \odot b_{ij}]$, which removes the parts of the first image that are below 0.13.

Airy disc II

- ▶ In formula form: $[a_{ij} \odot b_{ij}] = \begin{cases} a_{ij}, & \text{if } a_{ij} > b_{ij}. \\ 0, & \text{otherwise.} \end{cases}$
- ▶ We use the function $x \mapsto \lceil x \rceil$, which rounds up.
- ▶ If $a < b$, then $\lceil a - b \rceil$ is 0; otherwise, it is 1.
- ▶ Alternatively, $\lceil a - b \rceil$ is 1 if and only if $a > b$, meaning the pixel is above the threshold; otherwise, it is 0.
- ▶ Therefore, $a \odot b = a \lceil a - b \rceil$ is the desired operation.

The End

Thank you for your attention!