



(ENKEMNA0302) Applied Linear Algebra

Dr. Gabor FACSKO, PhD

Senior Research Fellow

facskog@gamma.ttk.pte.hu

University of Pecs, Faculty of Sciences, Institute of Mathematics and Informatics, 7624 Pecs, Ifjusag utja 6.
Wigner Research Centre for Physics, Department of Space Physics and Space Technology, 1121 Budapest, Konkoly-Thege Miklos ut 29-33.
<https://facsko.ttk.pte.hu>

February 12, 2025

Actualities

- ▶ I booked Room No. F07 in the Building F from 10:00 to 14:00 on May 15, 2025 for retake test.
- ▶ Are you interested in building a CubeSat?
- ▶ Recruit lecture will be held at UP Faculty of Engineering and Information Technology in Room B224 from 14:00 to 16:00 on Tuesday February 18, 2025.

Requirements

- ▶ You will write tests based on the exercises of the practical courses. You can use everything during the test
- ▶ The minimum requirement is 41 % of both tests.
- ▶ Failed tests must be corrected
- ▶ You must take an oral written exam. You cannot use anything
- ▶ Grades: Insufficient/Fail (1): 0-40 %, Sufficient/Pass (2): 41-55 %, Average (3): 56-70 %, Good (4): 71-85 %, Excellent (5): 86-100 %.
- ▶ Mid-term test 1: March 13, mid-term test 2: May 8, retake tests: May 15, 2025.

Bibliography


Bernard Kolman and David Hill: Elementary Linear Algebra with Applications, 9th ed., Person, 2007

Philip N. Klein: Coding the Matrix: Linear Algebra through Applications to Computer Science, Newtonian Press 2013

K. F. Riley, M. P. Hobson, S. J. Bence: Mathematical Methods for Physics and Engineering: A Comprehensive Guide, Cambridge University Press, 3rd. ed. (2006)


Operators I

- ▶ Definition: The set is the sum of things. It is a fundamental term. You need a statement that collects the element. It means that you can decide whether an element is part of the set or not.
- ▶ Definition: The pair are sets consisting of two elements.
- ▶ Definition: Elements e_1 and e_2 consist of ordered pair if $\{e_1, \{e_2\}\}$. Its sign is (e_1, e_2) .
- ▶ Definition: Relation is the set of ordered pairs.




- ▶ Definition: The injection orders different elements (X) to different elements (Y).

Operators II



- Definition: Surjections are those relations, that the values of the relation agree to the values of the set to order.



- Definition: Bijection is an injection and a surjection. All elements are related to all elements of the other set.

Operators III

- ▶ Definition: The functions are such a set of ordered pairs in that one element shows up only once:

$$(\forall x) (\forall y_1) (\forall y_2) [(x, y_1) \in f \wedge (x, y_2) \in f \Rightarrow y_1 = y_2]$$

- ▶ Definition: V and U vector spaces above \mathbb{T} body. The $f : V \rightarrow U$ relation is linear if it is
 1. Additive, for all $v_1, v_2 \in V$ vectors $f(v_1 + v_2) = f(v_1) + f(v_2)$.
 2. Homogen, for all $v \in V$ vectors and $\lambda \in \mathbb{T}$ elements $f(\lambda v) = \lambda f(v)$.
- ▶ Definition: Operators are the linear vector-vector functions.
- ▶ Például:
 - ▶ Identical operator: $\mathbf{A} \cdot \mathbf{1} = \mathbf{A}$, for all \mathbf{A} operators.
 - ▶ Null operator: $\mathbf{A} \cdot \mathbf{0} = \mathbf{0}$, for all \mathbf{A} operators.

Operators IV

- ▶ Mirror operators: $(\mathbf{A} \cdot \mathbf{M}) \cdot \mathbf{M} = \mathbf{A}$, for all \mathbf{A} operators.
- ▶ Projection operator: $\mathbf{A} \cdot \mathbf{P} = \mathbf{P}$, for all \mathbf{A} operator.
- ▶ Rotational operator: later.
- ▶ Operators could be multiplied on both sides.
- ▶ The representation of operators is the matrixes. See $\alpha_{ij} \in \mathbb{R}$ for all $i \in \{1, 2, \dots, m\}$ and $j \in \{1, 2, \dots, n\}$, where $m, n \in \mathbb{N}^+$. The

$$A = \begin{pmatrix} \alpha_{11} & \alpha_{12} & \cdots & \alpha_{1n} \\ \alpha_{21} & \alpha_{22} & \cdots & \alpha_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ \alpha_{m1} & \alpha_{m2} & \cdots & \alpha_{mn} \end{pmatrix}$$

table is called $m \times n$ type matrix. The set of the $m \times n$ type matrixes is $M_{m \times n}$.

Operators V

- ▶ The spur of the matrix is the set of $\{\alpha_{11}, \alpha_{22}, \dots, \alpha_{nn}\}$.
- ▶ The first index of the elements α_{ij} is the rowindex (i), the 2nd index is the column index (j).
- ▶ The Row i of the Matrix is A_i , and the Column j of the matrix is A_j .
- ▶ Determinant!!!

Transpose I

- Definition: The transpose of the $A = (\alpha_{ij})_{m \times n}$ matrix is the $A^T = (\alpha_{ji})_{n \times m}$. This means the change of the rows and columns. The transpose is the mirror of a square matrix.

$$A_{m \times n} = \begin{pmatrix} \alpha_{11} & \alpha_{12} & \cdots & \alpha_{1n} \\ \alpha_{21} & \alpha_{22} & \cdots & \alpha_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ \alpha_{m1} & \alpha_{m2} & \cdots & \alpha_{mn} \end{pmatrix} \quad A_{n \times m}^T = \begin{pmatrix} \alpha_{11} & \alpha_{21} & \cdots & \alpha_{1m} \\ \alpha_{12} & \alpha_{22} & \cdots & \alpha_{2m} \\ \vdots & \vdots & \ddots & \vdots \\ \alpha_{1n} & \alpha_{2n} & \cdots & \alpha_{nm} \end{pmatrix}$$

- Examples for transpose.

Matrix Operations I

- Definition: $A = (\alpha_{ij})_{m \times n}$ and $B = (\beta_{ij})_{m \times n}$ are two matrixes with same type, $\lambda \in \mathbb{R}$ a scalar. The sum of Matrixes A and B is Matrix $A + B = (\alpha_{ij} + \beta_{ij})_{m \times n}$, the λ times Matrix A is Matrix $\lambda A = (\lambda \alpha_{ij})_{m \times n}$.

$$A_{m \times n} = \begin{pmatrix} \alpha_{11} & \alpha_{12} & \cdots & \alpha_{1n} \\ \alpha_{21} & \alpha_{22} & \cdots & \alpha_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ \alpha_{m1} & \alpha_{m2} & \cdots & \alpha_{mn} \end{pmatrix} \quad B_{m \times n} = \begin{pmatrix} \beta_{11} & \beta_{12} & \cdots & \beta_{1n} \\ \beta_{21} & \beta_{22} & \cdots & \beta_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ \beta_{m1} & \beta_{m2} & \cdots & \beta_{mn} \end{pmatrix}$$

$$A_{m \times n} + B_{m \times n} = \begin{pmatrix} \alpha_{11} + \beta_{11} & \alpha_{12} + \beta_{12} & \cdots & \alpha_{1n} + \beta_{1n} \\ \alpha_{21} + \beta_{21} & \alpha_{22} + \beta_{22} & \cdots & \alpha_{2n} + \beta_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ \alpha_{m1} + \beta_{m1} & \alpha_{m2} + \beta_{m2} & \cdots & \alpha_{mn} + \beta_{mn} \end{pmatrix}$$

Matrix Operations II

$$\lambda A_{m \times n} = \begin{pmatrix} \lambda\alpha_{11} & \lambda\alpha_{12} & \cdots & \lambda\alpha_{1n} \\ \lambda\alpha_{21} & \lambda\alpha_{22} & \cdots & \lambda\alpha_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ \lambda\alpha_{m1} & \lambda\alpha_{m2} & \cdots & \lambda\alpha_{mn} \end{pmatrix}$$

The elements of the matrixes are added and multiplying by a scalar means to multiply all elements of the matrix by the scalar.

- Examples for matrix operations.

Matrix Operations III

- Definition: $A = (\alpha_{ij})_{m \times n}$ and $B = (\beta_{ij})_{n \times k}$ are two matrixes. The product of Matrixes A and B is Matrix $A \cdot B = (\gamma_{ij})_{m \times k}$, where

$$\gamma_{ij} = \sum_{l=1}^n \alpha_{il} \beta_{lj}.$$

Or:

$$A_{m \times n} = \begin{pmatrix} \alpha_{11} & \alpha_{12} & \cdots & \alpha_{1n} \\ \alpha_{21} & \alpha_{22} & \cdots & \alpha_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ \alpha_{i1} & \alpha_{i2} & \cdots & \alpha_{in} \\ \vdots & \vdots & \ddots & \vdots \\ \alpha_{m1} & \alpha_{m2} & \cdots & \alpha_{mn} \end{pmatrix} \quad B_{n \times k} = \begin{pmatrix} \beta_{11} & \beta_{12} & \cdots & \beta_{1j} & \cdots & \beta_{1k} \\ \beta_{21} & \beta_{22} & \cdots & \beta_{2j} & \cdots & \beta_{2k} \\ \vdots & \vdots & \ddots & \vdots & \ddots & \vdots \\ \beta_{n1} & \beta_{n2} & \cdots & \beta_{nj} & \cdots & \beta_{nk} \end{pmatrix}$$

$$A \cdot B_{m \times k} = \begin{pmatrix} \alpha_{11}\beta_{11} + \alpha_{12}\beta_{21} + \cdots + \alpha_{1n}\beta_{n1} & \alpha_{11}\beta_{12} + \alpha_{12}\beta_{22} + \cdots + \alpha_{1n}\beta_{n2} & \cdots & \alpha_{11}\beta_{1k} + \alpha_{12}\beta_{2k} + \cdots + \alpha_{1n}\beta_{nk} \\ \alpha_{21}\beta_{11} + \alpha_{22}\beta_{21} + \cdots + \alpha_{2n}\beta_{n1} & \alpha_{21}\beta_{12} + \alpha_{22}\beta_{22} + \cdots + \alpha_{2n}\beta_{n2} & \cdots & \alpha_{21}\beta_{1k} + \alpha_{22}\beta_{2k} + \cdots + \alpha_{2n}\beta_{nk} \\ \vdots & \vdots & \ddots & \vdots \\ \alpha_{m1}\beta_{11} + \alpha_{m2}\beta_{21} + \cdots + \alpha_{mn}\beta_{n1} & \alpha_{m1}\beta_{12} + \alpha_{m2}\beta_{22} + \cdots + \alpha_{mn}\beta_{n2} & \cdots & \alpha_{m1}\beta_{1k} + \alpha_{m2}\beta_{2k} + \cdots + \alpha_{mn}\beta_{nk} \end{pmatrix}$$

- Examples for matrix multiplications.

Matrix Operations IV

- ▶ For similar square matrixes the condition of matrix production is fulfilled and the product will be the same type. Therefore, there is an exponentiation of matrixes:

$$A^1 = A \quad \text{és} \quad A^m = AA^{m-1}$$

where $(m \geq 2)$ és $A \in \mathcal{M}_{n \times n}$. Let us consider $A^0 = E_m$.

- ▶ Thesis: Equations of matrix exponentation:

$$\begin{aligned} A^m A^k &= A^{m+k} \\ (A^m)^k &= A^{mk}, \end{aligned}$$

ahol $m, k \in \mathbb{N}$.

Deduction: It is trivial based on the definition of matrix product.

- ▶ Examples for matrix exponentation.

Identity Matrix

- Definition: The n^{th} order identity matrix is:

$$E_n = \begin{pmatrix} 1 & 0 & \cdots & 0 \\ 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 1 \end{pmatrix}$$

- Thesis: For all $A \in \mathcal{M}_{n \times n}$: $A \cdot E_n = E_n \cdot A = A$, or matrix E_n is identity element of the $n \times n$ square matrixes for matrix production.

Deduction: $A = (\alpha_{ij})_{n \times n}$ and $E_n = (\beta_{ij})_{n \times n}$ are two matrixes, where $\beta_{ij} = 1$, if $i = j$, otherwise it is zero. The product of Matrixes A and E_n is Matrix $A \cdot E_n = (\sum_{l=1}^n \alpha_{il} \beta_{lj})_{n \times n}$. It is Matrix $A = (\alpha_{ij})_{n \times n}$, because the definition of β_{ij} erases all other elements than α_{ij} .

Matrix Rank I

- ▶ Definition: $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_s \in V$ are vectors. The rank of the $\{\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_s\}$ vector system is the dimension of the $\mathcal{L}(\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_s)$ subspace. Its sign is $\rho(\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_s)$.
- ▶ Thesis: The following transformation do not change the order of the $\{\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_s\}$ vector system:
 1. Multiplying a vectors by a $\lambda \neq 0$ scalar.
 2. Adding the vector multiplied by λ to another vector.
 3. Eliminating a vector that is a linear combination of the remaining vectors.
 4. Changing the order of vectors.
- ▶ Definition: The rank of Matrix $A \in \mathcal{M}_{m \times n}$ is the rank of its row vector system.
- ▶ The rank of a matrix is equal to the common rank of the maximal ranked non-disappearing subdeterminants.

Matrix Rank II

- ▶ The rank of a matrix is determined by transforming the matrix to trapezoid form by rank invariant transformations. You can change the columns. (A matrix has trapezoid shape if $\alpha_{ij} = 0$, $i > j$, and $\alpha_{ii} \neq 0$, where $(1 \leq i \leq \min\{m, n\})$.) Rows and columns containing 0 could be deleted. The rank of the trapezoid matrix is the number of its rows.

- ▶ Examples of determination of the rank of a matrix.

Image processing I

[illegible]

- ▶ Matrix $\mathbf{A}_{m \times n}$ is a representation of an $m \times n$ greyscale image.
- ▶ All matrix elements gives color from the $\{0, 1, \dots, k\}$ range, where 0 is black, $k - 1$ is white and k means transparency.
- ▶ The $\mathbf{B}_{m \times n}$ background is transparent.
- ▶ Let us construct the $\mathbf{A} \odot \mathbf{B} = [a_{ij} \odot b_{ij}]$ operations that copies the 2nd image to the background of the first image.
- ▶ Formula: $[a_{ij} \odot b_{ij}] = \begin{cases} b_{ij}, & \text{if } a_{ij} = k. \\ a_{ij}, & \text{otherwise.} \end{cases}$
- ▶ Let us use the $x \mapsto \lfloor x \rfloor$ function, that means to rounding to the lower integer value.
- ▶ If $a \in [0, k]$, then $0 \leq a/k \leq 1$, therefore $\lfloor a/k \rfloor$ is 0, or 1.

Moment of inertia

- ▶ Mass times square of distance from the axis.
- ▶ For a rigid object of N point masses m_k , the moment of inertia tensor is given by

$$\mathbf{I} = \begin{pmatrix} I_{11} & I_{12} & I_{13} \\ I_{21} & I_{22} & I_{23} \\ I_{31} & I_{32} & I_{33} \end{pmatrix},$$

where

$$I_{ij} = \sum_{k=1}^N m_k \left(\|\mathbf{r}_k\|^2 \delta_{ij} - x_i^{(k)} x_j^{(k)} \right)$$

and $i, j \in \{1, 2, 3\}$, $\mathbf{r}_k = (x_1^{(k)}, x_2^{(k)}, x_3^{(k)})$ is the vector to the point mass m_k from the point about which the tensor is calculated, and δ_{ij} is the Kronecker delta.

Transformation matrixes I

- ▶ Rotational matrix in 2D:

$$\begin{pmatrix} \cos \alpha & -\sin \alpha \\ \sin \alpha & \cos \alpha \end{pmatrix}.$$

- ▶ Rotational matrixes in 3D around z, x, y axes:

$$\begin{pmatrix} \cos \alpha & -\sin \alpha & 0 \\ \sin \alpha & \cos \alpha & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} 1 & 0 & 1 \\ 0 & \cos \alpha & -\sin \alpha \\ 0 & \sin \alpha & \cos \alpha \end{pmatrix} \begin{pmatrix} \cos \alpha & 0 & \sin \alpha \\ 0 & 1 & 0 \\ -\sin \alpha & 0 & \cos \alpha \end{pmatrix}.$$

- ▶ Mirror of the vectors of the plan for the $\alpha/2$ angular line:

$$\begin{pmatrix} \cos \alpha & \sin \alpha \\ \sin \alpha & -\cos \alpha \end{pmatrix}.$$

Transformation matrixes II

- ▶ Mirror of the vectors of the 3D space for the \mathbf{n} normal vector planes:

$$\mathbf{M} = \mathbf{I} - 2\mathbf{n} \otimes \mathbf{n}^T.$$

- ▶ Perpendicular projection to a line with \mathbf{b} direction vector:

$$\mathbf{P} = \frac{1}{\mathbf{b}\mathbf{b}^T} \mathbf{b} \otimes \mathbf{b}^T.$$

- ▶ Perpendicular projection to the plan with \mathbf{n} normal vector:

$$\mathbf{P} = \mathbf{I} - \mathbf{n} \otimes \mathbf{n}^T.$$

Transformation matrixes III

- ▶ Shift by (a, b) vector in 2D:

$$\begin{pmatrix} 1 & 0 & a \\ 0 & 1 & b \\ 0 & 0 & 1 \end{pmatrix}.$$

- ▶ Shift by (a, b, c) vector in 3D:

$$\begin{pmatrix} 1 & 0 & 0 & a \\ 0 & 1 & 0 & b \\ 0 & 0 & 1 & c \\ 0 & 0 & 0 & 1 \end{pmatrix}.$$

Comming soon...

- ▶ Diagonal matrixes
- ▶ Permutation matrixes and snakes
- ▶ Triangular matrixes
- ▶ Symmetric and skew-symmetric matrixes

The End

Thank you for your attention!