



(PTIA0301) Elementary Linear Algebra

Dr. Gabor FACSKO, PhD

Senior Research Fellow

facskog@gamma.ttk.pte.hu

University of Pécs, Faculty of Sciences, Institute of Mathematics and Informatics, 7624 Pécs, Ifjuság útja 6.
Wigner Research Centre for Physics, Department of Space Physics and Space Technology, 1121 Budapest, Konkoly-Thege Miklós ut 29-33.
<https://facsko.ttk.pte.hu>

November 14, 2024

Announcements

- ▶ The tests can be reviewed during consultin hours.
- ▶ The 3rd test will be held in December. Students can take them to pass or improve their grades.
- ▶ I should grade your tests during the first week.
- ▶ I am sorry that some of you had difficulties with vector operations and the triple product.
- ▶ The Sarrus rule applies to 2×2 and 3×3 matrices. It is faster than the expansion theorem. It is not applicable to 4×4 matrices.
- ▶ The Cramer rule makes it harder to make mistakes when solving systems of linear equations.

Linear Transformations I

- ▶ Definition: Let V_1 and V_2 be vector spaces. A function $\varphi : V_1 \rightarrow V_2$ is called a linear mapping if

$$\text{additive : } \varphi(\mathbf{a} + \mathbf{b}) = \varphi(\mathbf{a}) + \varphi(\mathbf{b})$$

$$\text{and homogeneous : } \varphi(\lambda \mathbf{a}) = \lambda \varphi(\mathbf{a}),$$

where $\mathbf{a}, \mathbf{b} \in V_1$ and $\lambda \in \mathbb{R}$.

- ▶ Thesis: (Matrix Representation) A mapping $\varphi : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is linear if and only if $\exists A \in \mathcal{M}_{m \times n}$ such that $\varphi(\mathbf{x}) = A \cdot \mathbf{x}$, where $\mathbf{x} \in \mathbb{R}^n$.
- ▶ Definition: Let V be a vector space. A mapping $\varphi : V \rightarrow V$ is called a linear transformation. The set of all linear transformations on V is denoted by \mathcal{T}_V .
- ▶ Definition: A mapping $f : V \rightarrow \mathbb{R}$ is called a linear form.

Linear Transformations II

- Definition: A mapping $L : V \times V \rightarrow \mathbb{R}$ is called a bilinear form if it is linear in both arguments, that is,

$$L(\mathbf{x} + \mathbf{y}, \mathbf{z}) = L(\mathbf{x}, \mathbf{z}) + L(\mathbf{y}, \mathbf{z})$$

$$L(\lambda \mathbf{x}, \mathbf{y}) = \lambda L(\mathbf{x}, \mathbf{y})$$

$$L(\mathbf{x}, \mathbf{y} + \mathbf{z}) = L(\mathbf{x}, \mathbf{y}) + L(\mathbf{x}, \mathbf{z})$$

$$L(\mathbf{x}, \lambda \mathbf{y}) = \lambda L(\mathbf{x}, \mathbf{y}),$$

where $\mathbf{x}, \mathbf{y}, \mathbf{z} \in V$ and $\lambda \in \mathbb{R}$.

- Thesis: A mapping $L : V \times V \rightarrow \mathbb{R}$ is a bilinear form if and only if (for a given basis) there exist unique coefficients $\alpha_{ik} \in \mathbb{R}$ such that
$$L(x, y) = \sum_{i=1}^n \sum_{k=1}^n \alpha_{ik} x_i y_k.$$

Linear Transformations III

- ▶ Consider the canonical basis of \mathbb{R}^n , denoted as $\{\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n\}$. In this case, $\alpha_{ik} = L(\mathbf{e}_i, \mathbf{e}_k)$. The matrix $A = (\alpha_{ik})_{n \times n}$ is called the matrix of the bilinear form L (with respect to the canonical basis).
- ▶ Definition: A bilinear form L is symmetric if $L(\mathbf{x}, \mathbf{y}) = L(\mathbf{y}, \mathbf{x})$ for all $\mathbf{x}, \mathbf{y} \in V$.
- ▶ Definition: Let L be a symmetric bilinear form on the vector space V . The function $Q(x) = L(x, x)$ is called a quadratic form.
- ▶ Definition: A quadratic form Q is positive definite if $\forall \mathbf{x} \neq \mathbf{0}$ we have $Q(\mathbf{x}) > 0$.
Note: Q is positive semidefinite if $\forall \mathbf{x} \neq \mathbf{0}$ we have $Q(\mathbf{x}) \geq 0$ and $\exists \mathbf{y} \neq \mathbf{0}$ such that $Q(\mathbf{y}) = 0$. The concepts of negative definite and negative semidefinite are defined similarly.
- ▶ Definition: A symmetric bilinear form whose associated quadratic form is positive definite is called an inner product.
Example: In the space \mathbb{R}^3 , the scalar product is an inner product.

Linear Transformations IV

- ▶ A mapping $\varphi : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is called an isomorphism if it is bijective.
- ▶ It can be shown that two vector spaces are isomorphic if and only if their dimensions are equal:

$$V_1 \cong V_2 \Leftrightarrow \dim V_1 = \dim V_2.$$

The End

Thank you for your attention!