



(ENKEMNA0302) Applied Linear Algebra

Dr. Gabor FACSKO, PhD

Senior Research Fellow

facskog@gamma.ttk.pte.hu

University of Pécs, Faculty of Sciences, Institute of Mathematics and Informatics, 7624 Pécs, Ifjuság útja 6.
Wigner Research Centre for Physics, Department of Space Physics and Space Technology, 1121 Budapest, Konkoly-Thege Miklós ut 29-33.
<https://facsko.ttk.pte.hu>

March 20, 2025

Eigenvalue, Eigenvector, Eigenspace I

- Definition: (Determination of Eigenvalues) The characteristic polynomial of a matrix $A \in \mathcal{M}_{n \times n}$ is defined as the n^{th} -degree polynomial

$$f(x) = |A - xE_n| = \begin{vmatrix} a_{11} - x & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} - x & \dots & a_{2n} \\ \vdots & \vdots & \dots & \vdots \\ a_{n1} & a_{n2} & \dots & a_{nn} - x \end{vmatrix}.$$

- Statement: (Eigenvalues of Triangular Matrices). The eigenvalues of triangular matrices, and thus also of diagonal matrices, are equal to the elements of their main diagonal.

Eigenvalue, Eigenvector, Eigenspace II

- Statement: (Determinant, Trace, and Eigenvalues). If the eigenvalues of an $n \times n$ matrix \mathbf{A} are $\lambda_1, \dots, \lambda_n$, then

$$\det(\mathbf{A}) = \lambda_1 \lambda_2 \dots \lambda_n$$

$$\text{trace}(\mathbf{A}) = \lambda_1 + \lambda_2 + \dots + \lambda_n$$

These values appear in the characteristic polynomial: the determinant corresponds to the constant term, while the trace is the coefficient of $(-\lambda)^{n-1}$.

Eigenvalue, Eigenvector, Eigenspace III

- ▶ Theorem: (Eigenspaces of 2×2 Symmetric Matrices). Let $\mathbf{A} \in \mathbb{R}^{2 \times 2}$ be a symmetric matrix. Then:
 1. Every eigenvalue of \mathbf{A} is real.
 2. \mathbf{A} has two identical eigenvalues if and only if it is of the form $a\mathbf{I}$, in which case every vector in the plane is an eigenvector.
 3. If \mathbf{A} has two distinct eigenvalues, then its eigenspaces are orthogonal to each other.
- ▶ Theorem: (Matrix Invertibility and the Eigenvalue 0). A matrix \mathbf{A} is invertible if and only if 0 is not an eigenvalue.
- ▶ Theorem: (Eigenvalues of Special Matrices). Let \mathbf{A} be an $n \times n$ real matrix. Then:
 1. If \mathbf{A} is symmetric, all of its eigenvalues are real.
 2. If \mathbf{A} is skew-symmetric, all of its eigenvalues are imaginary.

The End

Thank you for your attention!