



(PTIA0301) Elementary Linear Algebra

Dr. Gabor FACSKO, PhD

Senior Research Fellow

facskog@gamma.ttk.pte.hu

University of Pécs, Faculty of Sciences, Institute of Mathematics and Informatics, 7624 Pécs, Ifjuság útja 6.
Wigner Research Centre for Physics, Department of Space Physics and Space Technology, 1121 Budapest, Konkoly-Thege Miklós ut 29-33.
<https://facsko.ttk.pte.hu>

October 11, 2024

Vector Space I

- Definition: The set $V \neq \emptyset$ a vector space above \mathbb{R} set, if there is an "+" operation with the following features:

$$\mathbf{a} + \mathbf{b} = \mathbf{b} + \mathbf{a}, \text{ where } (\mathbf{a}, \mathbf{b} \in V)$$

$$(\mathbf{a} + \mathbf{b}) + \mathbf{c} = \mathbf{a} + (\mathbf{b} + \mathbf{c}), \text{ where } (\mathbf{a}, \mathbf{b}, \mathbf{c} \in V)$$

$$\exists \mathbf{0} \in V \text{ that } \mathbf{a} + \mathbf{0} = \mathbf{a} \quad \forall \mathbf{a} \in V$$

$$\forall \mathbf{a} \in V \exists (-\mathbf{a}) \in V : \mathbf{a} + (-\mathbf{a}) = \mathbf{0},$$

Vector Space II

furthermore, for all $\lambda \in \mathbb{R}$ and for all $\mathbf{a} \in V$ the term $\lambda \mathbf{a} \in V$ exists, and the following features are fulfilled:

$$\lambda(\mu \mathbf{a}) = (\lambda\mu) \mathbf{a}, \text{ where } (\mathbf{a} \in V, \lambda, \mu \in \mathbb{R})$$

$$\lambda(\mathbf{a} + \mathbf{b}) = \lambda \mathbf{a} + \lambda \mathbf{b}, \text{ where } (\mathbf{a}, \mathbf{b} \in V, \lambda \in \mathbb{R})$$

$$(\lambda + \mu) \mathbf{a} = \lambda \mathbf{a} + \mu \mathbf{a}, \text{ where } (\mathbf{a} \in V \text{ és } \lambda, \mu \in \mathbb{R})$$

$$\forall \mathbf{a} \in V \text{ --re } 1 \cdot \mathbf{a} = \mathbf{a}.$$

These features are true for the V^2 and V^3 set, however the $\mathbb{R}^n = \mathbb{R} \times \mathbb{R} \times \dots \times \mathbb{R}$ set are vector space too. Furthermore, the set of the maximum n^{th} ordered polynomials ($R_n[x]$) are vector spaces too.

Vector Space III

- ▶ Definition: The V vector space's non empty subset L is a linear subspace if L also is a vector space with the operations in V .
- ▶ Thesis: The V vector space's L non-empty set is a linear subspace if and only if the following conditions are fulfilled:

$$\begin{aligned}\forall \mathbf{a}, \mathbf{b} \in L \quad \text{és} \quad \mathbf{a} + \mathbf{b} \in L \\ \forall \lambda \in \mathbb{R}, \forall \mathbf{a} \in L \quad \text{és} \quad \lambda \mathbf{a} \in L.\end{aligned}$$

- ▶ Definition: $H \neq \emptyset$ is subset of the V vector space. The subspace generated by H is the minimal size of such a subset of V that contains H . (Or it is a subset of all subspace containing H .) Sign: $\mathcal{L}(H)$.
- ▶ Always exists a minimal subspace containing H that is the intersection of all subspaces containing H .

Vector Space IV

- ▶ Definition: Set H is a generator system of V vector space if $\mathcal{L}(H) = V$.
- ▶ Definition: V vector space is generated finitely if it has a generator system containing a finite number of elements.
- ▶ Comment: The $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$ vector system is a generator system of a finitely generated V vector system if all elements of the V set could be written as the linear combination of $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$ vectors.
- ▶ Definition: The linear independent generator system of the V vector space is a basis of the V vector space.
- ▶ Tétel: In a finitely generated vector space all bases have the same cardinality.
- ▶ Definition: The common element number of the basis of a vector space is called dimension. Sign: $\dim(V)$.

Vector Space V

- ▶ Definition: $\{\mathbf{b}_1, \mathbf{b}_2, \dots, \mathbf{b}_n\}$ is a basis in V and $\mathbf{a} \in V$. Those $\lambda_1, \lambda_2, \dots, \lambda_n$ numbers for which $\mathbf{a} = \lambda_1 \mathbf{b}_1 + \lambda_2 \mathbf{b}_2 + \dots + \lambda_n \mathbf{b}_n$ we call coordinates the \mathbf{a} vector for the $\{\mathbf{b}_1, \mathbf{b}_2, \dots, \mathbf{b}_n\}$ basis.
- ▶ Definition: If $\mathbf{v} \in V^2 \setminus \{\mathbf{0}\}$, or $\mathbf{v} \in V^3 \setminus \{\mathbf{0}\}$. Then the $l = \{\alpha \mathbf{v} : \alpha \in \mathbb{R}\}$ set is a line (crossing the origo).
- ▶ Definition: If $\mathbf{u}, \mathbf{v} \in V^3 \setminus \{\mathbf{0}\}$ és $\nexists \lambda \in \mathbb{R}$, so $\mathbf{u} = \lambda \mathbf{v}$. Then the $L = \{\alpha \mathbf{u} + \beta \mathbf{v} : \alpha, \beta \in \mathbb{R}\}$ set a plane (crossing the origo).
- ▶ The set $L = \{\alpha_1 \mathbf{u}_1 + \alpha_2 \mathbf{u}_2 + \dots + \alpha_m \mathbf{u}_m : \alpha_1, \dots, \alpha_m \in \mathbb{R}\}$ is just the linear subspace generated by $\{\mathbf{u}_1, \dots, \mathbf{u}_m\}$. If $\mathbf{u}_1, \dots, \mathbf{u}_m$ are linear independent, then the set L is an mD subspace- Then the set $K = \{\alpha_1 \mathbf{u}_1 + \alpha_2 \mathbf{u}_2 + \dots + \alpha_m \mathbf{u}_m + \mathbf{v} : \alpha_1, \dots, \alpha_m \in \mathbb{R}\}$ is an mD affine subspace. All affine subspace can be constructed in the $K = L + \mathbf{v}$ form, where L is a linear subspace and $\mathbf{v} \in V$.

Vector Space VI

- ▶ The equation of a plane which has the $\mathbf{n}(n_1, n_2, n_3)$ normal vector and crossed the point $P(p_1, p_2, p_3)$ is $n_1x + n_2y + n_3z = n_1p_1 + n_2p_2 + n_3p_3$.
- ▶ Definition: The (Minkowski-)sum of set is $A + B = \{a + b : a \in A, b \in B\}$.
- ▶ Állítás: The sum and intersection of subspaces are $L_1 + L_2$ and $L_1 \cap L_2$ is altér.
- ▶ Definition: The $L_1 + L_2$ is a direct sum if $L_1 \cap L_2 = \{\mathbf{0}\}$.

The End

Thank you for your attention!