



(ENKEMNA0302) Applied Linear Algebra

Dr. Gabor FACSKO, PhD

Senior Research Fellow

facskog@gamma.ttk.pte.hu

University of Pécs, Faculty of Sciences, Institute of Mathematics and Informatics, 7624 Pécs, Ifjuság útja 6.
Wigner Research Centre for Physics, Department of Space Physics and Space Technology, 1121 Budapest, Konkoly-Thege Miklós ut 29-33.
<https://facsko.ttk.pte.hu>

March 5, 2025

Operations with Block Matrices I

- ▶ Operations on extremely large matrices can be parallelized if the matrices are divided into blocks and the operations are performed on these smaller submatrices.
- ▶ When a matrix is divided into submatrices by horizontal and vertical lines, we say that this matrix is a block matrix composed of submatrices—also known as blocks.
- ▶ The rows and columns of a block matrix are called the block rows and block columns of the matrix.
- ▶ Block matrices are also referred to as hypermatrices, but this term is also used for multidimensional arrays. Therefore, we prefer to call them block matrices.

Operations with Block Matrices II

- ▶ The augmented matrix $[\mathbf{A}|\mathbf{b}]$ of a system of equations is a block matrix consisting of two blocks:

$$\left(\begin{array}{ccc|cc|c} 1 & 0 & 0 & 1 & 2 & 4 \\ 0 & 1 & 0 & 2 & 0 & 3 \\ 0 & 0 & 1 & 1 & 0 & 3 \\ \hline 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{array} \right)$$

This is the augmented matrix of a system of 5 equations with 5 unknowns. The first block column corresponds to the bound variables, the second to the free variables, and the third to the right-hand side of the system of equations. The second block row contains the zero rows.

Operations with Block Matrices III

- Statement: (Operations with Block Matrices). Multiplication of block matrices by a scalar and the addition of two block matrices partitioned in the same way can be performed block-wise, that is,

$$c[\mathbf{A}_{ij}] = [c\mathbf{A}_{ij}], \quad [\mathbf{A}_{ij}] + [\mathbf{B}_{ij}] = [\mathbf{A}_{ij} + \mathbf{B}_{ij}].$$

If $\mathbf{A} = [\mathbf{A}_{ik}]_{m \times t}$ and $\mathbf{B} = [\mathbf{B}_{kj}]_{t \times n}$ are two block matrices, and for every k , the number of columns of \mathbf{A}_{ik} matches the number of rows of \mathbf{B}_{kj} , then the product $\mathbf{C} = \mathbf{AB}$ can be computed by applying the multiplication rule to blocks, that is, \mathbf{C} is a block matrix where the block in the i -th block row and j -th block column is given by:

$$\mathbf{C}_{ij} = \sum_{k=1}^t \mathbf{A}_{ik} \mathbf{B}_{kj}.$$

Operations with Block Matrices IV

- Example of block matrix multiplication:

$$\left(\begin{array}{cc|c} 1 & 0 & 1 \\ 2 & 1 & 1 \\ 0 & 3 & 1 \end{array} \right) \left(\begin{array}{c|c} 1 & 1 \\ 1 & 2 \\ 0 & 1 \end{array} \right) = \left(\begin{array}{cc|c} (1 \ 0) \begin{pmatrix} 1 \\ 1 \end{pmatrix} + (1) (0) & (1 \ 0) \begin{pmatrix} 1 \\ 2 \end{pmatrix} + (1) (1) \\ \hline \begin{pmatrix} 2 & 1 \\ 0 & 3 \end{pmatrix} \begin{pmatrix} 1 \\ 1 \end{pmatrix} + \begin{pmatrix} 1 \\ 1 \end{pmatrix} (0) & \begin{pmatrix} 2 & 1 \\ 0 & 3 \end{pmatrix} \begin{pmatrix} 1 \\ 2 \end{pmatrix} + \begin{pmatrix} 1 \\ 1 \end{pmatrix} (1) \end{array} \right) =$$
$$\left(\begin{array}{c|c} \begin{pmatrix} 1 \\ 3 \\ 3 \end{pmatrix} & \begin{pmatrix} 2 \\ 5 \\ 7 \end{pmatrix} \end{array} \right) = \left(\begin{array}{c|c} 1 & 2 \\ 3 & 5 \\ 3 & 7 \end{array} \right).$$

Kronecker Product and the Vec Function I

- ▶ Certain block matrix operations cannot be derived from simple matrix operations.
- ▶ The vec function transforms an arbitrary matrix into a vector by stacking its column vectors on top of each other. If $\mathbf{A} = [\mathbf{a}_1 | \mathbf{a}_2 | \dots | \mathbf{a}_n]$, then

$$\text{vec}(\mathbf{A}) = \begin{pmatrix} \mathbf{a}_1 \\ \vdots \\ \mathbf{a}_n \end{pmatrix}$$

For example, if $\mathbf{A} = \begin{pmatrix} 1 & 2 \\ 3 & 4 \end{pmatrix}$, then $\text{vec}(\mathbf{A}) = \begin{pmatrix} 1 \\ 2 \\ 3 \\ 4 \end{pmatrix}$.

Kronecker Product and the Vec Function II

- ▶ Let \mathbf{A} be an $m \times n$ matrix and \mathbf{B} a $p \times q$ matrix. Their Kronecker product (also called the tensor product) is the $mp \times nq$ matrix denoted by $\mathbf{A} \otimes \mathbf{B}$, which has the block matrix form:

$$\mathbf{A} \otimes \mathbf{B} = \begin{pmatrix} a_{11}\mathbf{B} & a_{12}\mathbf{B} & \dots & a_{1n}\mathbf{B} \\ a_{21}\mathbf{B} & a_{22}\mathbf{B} & \dots & a_{2n}\mathbf{B} \\ \vdots & \vdots & \dots & \vdots \\ a_{m1}\mathbf{B} & a_{m2}\mathbf{B} & \dots & a_{mn}\mathbf{B} \end{pmatrix}.$$

- ▶ For example,

$$\begin{pmatrix} -1 & 2 \\ 0 & 1 \end{pmatrix} \otimes \begin{pmatrix} 0 & 1 & 2 \\ 3 & 3 & 3 \end{pmatrix} = \begin{pmatrix} 0 & -1 & -2 & 0 & 2 & 4 \\ -3 & -3 & -3 & 6 & 6 & 6 \\ 0 & 0 & 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 3 & 3 & 3 \end{pmatrix}.$$

Kronecker Product and the Vec Function III

► Theorem: (Properties of the Kronecker Product). Given the matrices $\mathbf{A}_{m \times n}$, $\mathbf{B}_{m \times n}$, $\mathbf{C}_{p \times q}$, and $\mathbf{D}_{r \times s}$, we have:

1. $(\mathbf{A} + \mathbf{B}) \otimes \mathbf{C} = \mathbf{A} \otimes \mathbf{C} + \mathbf{B} \otimes \mathbf{C}$, $\mathbf{C} \otimes (\mathbf{A} + \mathbf{B}) = \mathbf{C} \otimes \mathbf{A} + \mathbf{C} \otimes \mathbf{B}$,
2. $(\mathbf{A} \otimes \mathbf{C}) \otimes \mathbf{D} = \mathbf{A} \otimes (\mathbf{C} \otimes \mathbf{D})$,
3. $(\mathbf{A} \otimes \mathbf{C})^T = \mathbf{C}^T \otimes \mathbf{A}^T$.

Proof: By definition, e.g., for (2):

$$(\mathbf{AXB})_{*j} = \mathbf{AXB}_{*j} = \sum_{i=1}^n (b_{ij}\mathbf{A}) \mathbf{X}_{*i} = (b_{1j}\mathbf{A} | \dots | b_{nj}\mathbf{A}) \text{vec}(\mathbf{X}) = \left(\mathbf{B}^T \otimes \mathbf{A} \right)_{*j} \text{vec}(\mathbf{X}).$$

Hypermatrixes I

- ▶ Certain data can be well organized in arrays of dimension higher than 2.
- ▶ Definition: (Hypermatrix). Let $n_1, n_2, \dots, n_d \in \mathbb{N}^+$ and let S be an arbitrary set (e.g., $S = \mathbb{R}, \mathbb{Q}, \mathbb{N}, \mathbb{Z} \dots$). A d th-order (or d -dimensional) $n_1 \times n_2 \times \dots \times n_d$ -type hypermatrix is a mapping of the form

$$\mathbf{A} : \{1, \dots, n_1\} \times \{1, \dots, n_2\} \times \dots \times \{1, \dots, n_d\} \rightarrow S$$

The element $\mathbf{A}(i_1, i_2, \dots, i_d)$ is denoted by $a_{i_1 i_2 \dots i_d}$, which is an element of a d -dimensional table, and similarly to matrices, we can write:

$$\mathbf{A} = (a_{i_1 i_2 \dots i_d})_{i_1, i_2, \dots, i_d}^{n_1, n_2, \dots, n_d} = 1, \text{ or simply } \mathbf{A} = (a_{i_1 i_2 \dots i_d}).$$

If $n_1 = n_2 = \dots = n_d = n$, then we refer to a hyper-cube matrix.

Hypermatrices II

- ▶ The set of all $n_1 \times n_2 \times \cdots \times n_d$ -type hypermatrices formed from the elements of S is denoted by $S^{n_1 \times n_2 \times \cdots \times n_d}$.
- ▶ Second-order hypermatrices coincide with matrices.
- ▶ The elements of third-order hypermatrices can be described by slicing them according to the third index. Each slice is a matrix, and these matrices are written next to each other separated by vertical lines. For example, the general form of a $4 \times 2 \times 3$ -type hypermatrix is:

$$\begin{pmatrix} a_{111} & a_{121} & a_{112} & a_{122} & a_{113} & a_{123} \\ a_{211} & a_{221} & a_{212} & a_{222} & a_{213} & a_{223} \\ a_{311} & a_{321} & a_{312} & a_{322} & a_{313} & a_{323} \\ a_{411} & a_{421} & a_{412} & a_{422} & a_{413} & a_{423} \end{pmatrix}$$

Hypermatrixes III

- Addition of two hypermatrixes of the same type and scalar multiplication of a hypermatrix follow element-wise operations similar to matrices:

$$(a_{i_1 i_2 \dots i_d}) + (b_{i_1 i_2 \dots i_d}) = (a_{i_1 i_2 \dots i_d} + b_{i_1 i_2 \dots i_d}),$$
$$c(a_{i_1 i_2 \dots i_d}) = (ca_{i_1 i_2 \dots i_d}).$$

- Definition: (Transpose of a hypermatrix). Let π be a permutation of the set $\{1, 2, \dots, d\}$. The π -transpose of the d th-order hypermatrix $\mathbf{A} = (a_{i_1 i_2 \dots i_d}) \in S^{n_1 \times n_2 \times \dots \times n_d}$ is defined as:

$$\mathbf{A}^\pi = \left(a_{i_{\pi(1)} i_{\pi(2)} \dots i_{\pi(d)}} \right) \in S^{n_{\pi(1)} \times n_{\pi(2)} \times \dots \times n_{\pi(d)}}$$

A hyper-cube matrix $\mathbf{A} \in S^{n \times n \times \dots \times n}$ is symmetric if for all permutations π , we have $\mathbf{A}^\pi = \mathbf{A}$, and it is skew-symmetric if $\mathbf{A}^\pi = \text{sgn}(\pi) \mathbf{A}$, where $\text{sgn}(\pi) = -1$ for odd permutations and 1 for even permutations.

Hypermatrices IV

- Accordingly, the general form of $2 \times 2 \times 2$ hypermatrices and symmetric hypermatrices is:

$$\left(\begin{array}{cc|cc} a_{111} & a_{121} & a_{112} & a_{122} \\ a_{211} & a_{221} & a_{212} & a_{222} \end{array} \right), \quad \left(\begin{array}{cc|cc} a & b & b & c \\ b & c & c & d \end{array} \right)$$

The general form of $3 \times 3 \times 3$ hypermatrices, symmetric, and skew-symmetric hypermatrices is:

$$\left(\begin{array}{ccc|ccc|ccc} a_{111} & a_{121} & a_{131} & a_{112} & a_{122} & a_{132} & a_{113} & a_{123} & a_{133} \\ a_{211} & a_{221} & a_{231} & a_{212} & a_{222} & a_{232} & a_{213} & a_{223} & a_{233} \\ a_{311} & a_{321} & a_{331} & a_{312} & a_{322} & a_{332} & a_{313} & a_{323} & a_{333} \end{array} \right),$$
$$\left(\begin{array}{ccc|ccc|ccc} a & b & c & b & d & e & c & e & f \\ b & d & e & d & g & h & e & h & i \\ c & e & f & e & h & i & f & i & j \end{array} \right), \quad \left(\begin{array}{ccc|ccc|ccc} 0 & 0 & 0 & 0 & 0 & -a & 0 & a & 0 \\ 0 & 0 & a & 0 & 0 & 0 & -a & 0 & 0 \\ 0 & -a & 0 & a & 0 & 0 & 0 & 0 & 0 \end{array} \right),$$

where $a, b, c, d, e, f, g, h, i, j \in S$ are not necessarily distinct elements.

The End

Thank you for your attention!