



# (PTIA0301) Elementary Linear Algebra

Dr. Gabor FACSKO, PhD

Senior Research Fellow

*facskog@gamma.ttk.pte.hu*

University of Pécs, Faculty of Sciences, Institute of Mathematics and Informatics, 7624 Pécs, Ifjuság útja 6.  
Wigner Research Centre for Physics, Department of Space Physics and Space Technology, 1121 Budapest, Konkoly-Thege Miklós ut 29-33.  
<https://facsko.>

September 5, 2024

# Course Objectives

- ▶ Terms and basic methods of Linear Algebra methodology
- ▶ Using elementary methods of Linear Algebra to fix simple problems
- ▶ Recognition when Linear Algebra could be used to solve problems

# Requirements

- ▶ You will write tests based on the exercises of the practical courses.
- ▶ The minimum requirement is 40 % of both tests.
- ▶ I will offer a grade based on the test results for the colloquium if you get at least an Average (3) grade.
- ▶ If you do not like the offered grade, you can take a (written) exam.
- ▶ Grades: Insufficient/Fail (1): 0-40 %, Sufficient/Pass (2): 41-55 %, Average (3): 56-70 %, Good (4): 71-85 %, Excellent (5): 86-100 %.

## Bibliography

*Bernard Kolman and David Hill: Elementary Linear Algebra with Applications, 9th ed., Person, 2007*


*Henry Ricardo: A Modern Introduction to Linear Algebra, Taylor & Francis Group, LLC, 2010*

# Scalars and vectors

- ▶ Scalars are quantities without directions. For example mass ( $m$ ), speed ( $|\mathbf{v}|$ , *not* velocity), temperature ( $T$ ), length, volume ( $V$ ).
- ▶ Vectors are quantities with direction. For example: weight ( $\mathbf{F}$ ), velocity ( $\mathbf{v}$ ), position ( $\mathbf{r}$ ), acceleration/deceleration ( $\mathbf{a}$ ), rotation, rotational speed ( $\omega$ ).
- ▶ Two types of vectors:
  - ▶ Weight ( $\mathbf{F}$ ), velocity ( $\mathbf{v}$ ), position ( $\mathbf{r}$ ), acceleration/deceleration ( $\mathbf{a}$ ).
  - ▶ Rotation, rotational speed ( $\omega$ ).
- ▶ You can write vectors in three different ways: boldface ( $\mathbf{v}$ ), underline ( $\underline{v}$ ), and arrow ( $\vec{v}$ )
- ▶ Weird association by **François Villon (1431– after 1463)**: *"Francis I am, which weighs me down, / born in Paris near Pontoise town, / and with a stretch of rope my pate / will learn for once my arse's weight."*

## Other definitions for vectors I

- ▶ Limited length line segments from Point A to Point B:  $\vec{AB}$ . The start point is Point A, and the end is Point B.



- ▶ Two vectors are equal if you can transform the first vector to the second vector using parallel shift/translation displacement.
- ▶ Or, if the lengths and directions of the vectors are the same.
- ▶ The vectors you can transform to each other form the group of additive vectors.

## Other definitions for vectors II


- ▶ Definition: Planar ( $V^2$ ) or spatial vectors ( $V^3$ ) are the group of those that you can transform to each other using translation (parallel displacement).

# Coordinates of vectors I

- The pairs/triplets of numbers ( $\mathbb{R}^2$ , or  $\mathbb{R}^3$ ) can identify as vectors:

$$\mathbf{v} = (v_1, v_2) = \begin{pmatrix} v_1 \\ v_2 \end{pmatrix},$$


where  $v_1 \in \mathbb{R}$ ,  $v_2 \in \mathbb{R}$  are the components of the vector in 2D.



## Coordinates of vectors II

$$\mathbf{v} = (v_1, v_2, v_3) = \begin{pmatrix} v_1 \\ v_2 \\ v_3 \end{pmatrix},$$

where  $v_1 \in \mathbb{R}$ ,  $v_2 \in \mathbb{R}$ ,  $v_3 \in \mathbb{R}$  are the components of the vector in 3D.






## Length and equality of vectors I

- ▶ Definition: Two vectors are equal, and only equal if their origo centred representations are the same.
- ▶ In coordinates it means that  $\mathbf{a}(a_1, a_2, a_3)$  and  $\mathbf{b}(b_1, b_2, b_3)$  are equal, and only equal if  $a_1 = b_1$ ,  $a_2 = b_2$ , and  $a_3 = b_3$ , where  $a_1, a_2, a_3 \in \mathbb{R}$ .

## Length and equality of vectors II

- Thesis: The magnitude of the  $\mathbf{a} = (a_1, a_2, a_3)$  vector is the following non-zero number:

$$|\mathbf{a}| = \|\mathbf{a}\| = \sqrt{a_1^2 + a_2^2 + a_3^2}.$$



Deduction: Points AOM form a right triangle, where the right angle is located at  $\angle OMA$ . Applying the Pythagoras-thesis you got  $|\mathbf{a}|^2 = OM^2 + a_3^2$ . Points O,  $(a_1, 0, 0)$ ,  $(0, a_2, 0)$  form a right triangle too, where the right angle is located at  $\angle [O, (a_1, 0, 0), (0, a_2, 0)]$  according too the Pythagoras theorem:  $OM^2 = a_1^2 + a_2^2$ . After substituting the later equation into the first equation we got  $|\mathbf{a}|^2 = OM^2 + a_3^2 = a_1^2 + a_2^2 + a_3^2$ , therefore  $|\mathbf{a}| = \sqrt{a_1^2 + a_2^2 + a_3^2}$ .  
q. e. d.

## Length and equality of vectors III


- ▶ The null vector has no length:  $|\mathbf{0}(0, 0, 0)|=0$ .

# Multiply vector with a scalar, sum of vectors, and difference of vectors I

- Definition: Sum of vectors. If  $\mathbf{a} (a_1, a_2, a_3)$  and  $\mathbf{b} (b_1, b_2, b_3)$ , then

$$\mathbf{a} + \mathbf{b} = (a_1 + b_1, a_2 + b_2, a_3 + b_3),$$

where  $a_1, a_2, a_3, b_1, b_2, b_3 \in \mathbb{R}$ .




*To add vectors  $\mathbf{a}$  and  $\mathbf{b}$  you should transform vector  $\mathbf{b}$  to the end point of the vector  $\mathbf{a}$ . Their sum  $(\mathbf{a} + \mathbf{b})$  is the vector from the start point of  $\mathbf{a}$  to the endpoint of the vector  $\mathbf{b}$ .*

## Multiply vector with a scalar, sum of vectors, and difference of vectors II

- Definition: Multiplication of vectors with a scalar. If  $\lambda \in \mathbb{R}$  and  $\mathbf{a} (a_1, a_2, a_3)$ , where  $a_1, a_2, a_3 \in \mathbb{R}$ , then

$$\lambda \mathbf{a} = (\lambda a_1, \lambda a_2, \lambda a_3).$$

Mind if  $\lambda$  is 0, 1, -1,  $<1$ , or  $>1$ .




*To multiply an  $\mathbf{a}$  vector by a  $\lambda$  number, you should draw a vector from the starting point of  $\mathbf{a}$  vector in the direction of  $\mathbf{a}$  that has  $\lambda$  length of the  $\mathbf{a}$  vector.*

## Multiply vector with a scalar, sum of vectors, and difference of vectors III

- Definition: Subtraction of vectors. If  $\mathbf{a} (a_1, a_2, a_3)$  and  $\mathbf{b} (b_1, b_2, b_3)$ , then

$$\mathbf{a} - \mathbf{b} = (a_1 - b_1, a_2 - b_2, a_3 - b_3),$$

where  $a_1, a_2, a_3, b_1, b_2, b_3 \in \mathbb{R}$ .



*To get the difference of vectors **a** and **b**, you should transfer to the vectors in a common start point. The difference of the vectors (**a** - **b**) starts from the end point of the vector **b** to the endpoint of vector **a**.*

# Multiply vector with a scalar, sum of vectors, and difference of vectors IV

► The sum of vectors is

1. Thesis: Commutative  $\mathbf{a} + \mathbf{b} = \mathbf{b} + \mathbf{a}$ , where  $\mathbf{a}, \mathbf{b} \in \mathbb{R}^3$ .

Deduction:  $\mathbf{a} + \mathbf{b} = (a_1, a_2, a_3) + (b_1, b_2, b_3) = (a_1 + b_1, a_2 + b_2, a_3 + b_3) =$   
 $(b_1 + a_1, b_2 + a_2, b_3 + a_3) = (b_1, b_2, b_3) + (a_1, a_2, a_3) = \underline{\underline{\mathbf{b} + \mathbf{a}}}$  *q. e. d.*

2. Thesis: Associative  $(\mathbf{a} + \mathbf{b}) + \mathbf{c} = \mathbf{a} + (\mathbf{b} + \mathbf{c})$ , where  $\mathbf{a}, \mathbf{b}, \mathbf{c} \in \mathbb{R}^3$ .

Deduction:  $(\mathbf{a} + \mathbf{b}) + \mathbf{c} = [(a_1, a_2, a_3) + (b_1, b_2, b_3)] + (c_1, c_2, c_3) =$   
 $[(a_1 + b_1) + c_1, (a_2 + b_2) + c_2, (a_3 + b_3) + c_3] =$   
 $[a_1 + (b_1 + c_1), a_2 + (b_2 + c_2), a_3 + (b_3 + c_3)] =$   
 $(a_1, a_2, a_3) + [(b_1, b_2, b_3) + (c_1, c_2, c_3)] = \underline{\underline{\mathbf{a} + (\mathbf{b} + \mathbf{c})}}$  *q. e. d.*

3. The null vector exists:  $\exists \mathbf{0} \in \mathbb{R}^3$ , where  $\mathbf{a} + \mathbf{0} = \mathbf{a}$ , and  $\mathbf{a} \in \mathbb{R}^3$ .

4. All vectors have an inverse vector:  $\forall \mathbf{a} \in \mathbb{R}^3 \exists (-\mathbf{a}) \in \mathbb{R}^3$ , where  $\mathbf{a} + (-\mathbf{a}) = \mathbf{0}$ .

# Multiply vector with a scalar, sum of vectors, and difference of vectors V

► The vector multiplication with scalar is

- Thesis: Multiplication of vectors by a scalar is associative,  $\lambda(\mu \mathbf{a}) = (\lambda\mu) \mathbf{a}$ , where  $\mathbf{a} \in \mathbb{R}^3$ ,  $\lambda, \mu \in \mathbb{R}$ .

Deduction:  $\lambda(\mu \mathbf{a}) = \lambda[\mu(a_1, a_2, a_3)] = \lambda(\mu a_1, \mu a_2, \mu a_3) = (\lambda\mu a_1, \lambda\mu a_2, \lambda\mu a_3) = (\lambda\mu)(a_1, a_2, a_3) = \underline{(\lambda\mu) \mathbf{a}}$  *q.e.d.*

- Thesis: Addition of vectors is distributive for the multiplication by a scalar,  $\lambda(\mathbf{a} + \mathbf{b}) = \lambda \mathbf{a} + \lambda \mathbf{b}$ , where  $\mathbf{a}, \mathbf{b} \in \mathbb{R}^3$ ,  $\lambda \in \mathbb{R}$ .

Deduction:  $\lambda(\mathbf{a} + \mathbf{b}) = \lambda[(a_1, a_2, a_3) + (b_1, b_2, b_3)] = \lambda(a_1 + b_1, a_2 + b_2, a_3 + b_3) = [\lambda(a_1 + b_1), \lambda(a_2 + b_2), \lambda(a_3 + b_3)] = (\lambda a_1 + \lambda b_1, \lambda a_2 + \lambda b_2, \lambda a_3 + \lambda b_3) = (\lambda a_1, \lambda a_2, \lambda a_3) + (\lambda b_1, \lambda b_2, \lambda b_3) = \lambda(a_1, a_2, a_3) + \lambda(b_1, b_2, b_3) = \underline{\underline{\lambda \mathbf{a} + \lambda \mathbf{b}}}$  *q. e. d*



# Multiply vector with a scalar, sum of vectors, and difference of vectors VI

- Thesis: The addition of scalars is distributive for multiplication by a vector,

$$(\lambda + \mu) \mathbf{a} = \lambda \mathbf{a} + \mu \mathbf{a}, \text{ where } \mathbf{a} \in \mathbb{R}^3, \lambda, \mu \in \mathbb{R}.$$

$$\text{Deduction: } (\lambda + \mu) \mathbf{a} = (\lambda + \mu) (a_1, a_2, a_3) = [(\lambda + \mu) a_1, (\lambda + \mu) a_2, (\lambda + \mu) a_3] =$$

$$(\lambda a_1 + \mu a_1, \lambda a_2 + \mu a_2, \lambda a_3 + \mu a_3) = (\lambda a_1, \lambda a_2, \lambda a_3) + (\mu a_1, \mu a_2, \mu a_3) =$$

$$\lambda (a_1, a_2, a_3) + \mu (a_1, a_2, a_3) = \underline{\underline{\lambda \mathbf{a} + \mu \mathbf{a}}} \quad \text{q. e. d.}$$

- $\forall \mathbf{a} \cdot 1 = \mathbf{a}$ , where  $\mathbf{a} \in \mathbb{R}^3$ .

# The End

Thank you for your attention!