



# (ENKEMNA0302) Applied Linear Algebra

Dr. Gabor FACSKO, PhD

Senior Research Fellow

*facskog@gamma.ttk.pte.hu*

University of Pecs, Faculty of Sciences, Institute of Mathematics and Informatics, 7624 Pecs, Ifjusag utja 6.  
Wigner Research Centre for Physics, Department of Space Physics and Space Technology, 1121 Budapest, Konkoly-Thege Miklos ut 29-33.  
<https://facsko.ttk.pte.hu>

April 10, 2025

## Orthonormal basis - orthogonal matrix I

- ▶ Orthogonal basis: the basis vectors are pairwise orthogonal, since in that case the scalar product of different basis vectors is 0. I
- ▶ Orthonormal basis: if the basis vectors are unit vectors.
- ▶ A set of pairwise orthogonal vectors is called an orthogonal basis (OB). An orthogonal system may contain zero vectors.
- ▶ A set of pairwise orthogonal unit vectors is called an orthonormal basis (ONB). Orthonormal bases do not contain zero vectors.
- ▶ From an orthogonal basis, an orthonormal basis can always be obtained by normalizing each basis vector (i.e., dividing each by its length).

## Orthonormal basis - orthogonal matrix II

- ▶ Independence of orthogonal vectors: Wettl notes
- ▶ Best approximation in the case of an ONB: Wettl notes
- ▶ Definition: (Orthogonal and semi-orthogonal matrix). A real square matrix is called orthogonal if its column vectors or row vectors form an orthonormal system. If we do not require the matrix to be square, we speak of a semi-orthogonal matrix.
- ▶ Theorem: (Equivalent definitions of semi-orthogonal matrices) Let  $m \geq n$  and  $\mathbf{Q} \in \mathbb{R}^{m \times n}$ . The following statements are equivalent:

1.  $\mathbf{Q}$  is semi-orthogonal,
2.  $\mathbf{Q}^T \mathbf{Q} = \mathbf{E}_n$ .

Similarly, for  $m \leq n$ ,  $\mathbf{Q}$  is semi-orthogonal if and only if  $\mathbf{Q} \mathbf{Q}^T = \mathbf{E}_m$ . Statement 2 expresses in algebraic terms that for  $m \leq n$ ,  $\mathbf{Q}$  is semi-orthogonal if and only if its transpose is its left inverse.

Proof: Wettl notes.

## Orthonormal basis - orthogonal matrix III

► Theorem: (Equivalent definitions of orthogonal matrices). Let  $\mathbf{Q} \in \mathbb{R}^{n \times n}$ . The following statements are equivalent:

1. The column vectors of  $\mathbf{Q}$  form an orthonormal system.
2.  $\mathbf{Q}^T \mathbf{Q} = \mathbf{E}_n$ .
3.  $\mathbf{Q}^{-1} = \mathbf{Q}^T$ .
4.  $\mathbf{Q} \mathbf{Q}^T = \mathbf{E}_n$ .
5. The row vectors of  $\mathbf{Q}$  form an orthonormal system.

Proof: Wette notes.

► Theorem: (Matrix transformation associated with an orthogonal matrix) Let  $\mathbf{Q} \in \mathbb{R}^{n \times n}$ . The following statements are equivalent:

1.  $\mathbf{Q}$  is orthogonal.
2.  $|\mathbf{Q}\mathbf{x}| = |\mathbf{x}|$  for all  $\mathbf{x} \in \mathbb{R}^n$ .
3.  $\mathbf{Q}\mathbf{x} \cdot \mathbf{Q}\mathbf{y} = \mathbf{x} \cdot \mathbf{y}$  for all  $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$ .

Proof: Wette notes.

## Orthonormal basis - orthogonal matrix IV

► Theorem: (Properties of orthogonal matrices)

1. If  $\mathbf{Q}$  is a real orthogonal matrix, then  $|\det(\mathbf{Q})| = 1$ .
2. The set of real  $n \times n$  orthogonal matrices, denoted by  $O(n)$ , is closed under matrix multiplication and inversion.

Proof: Wettl notes.

- Theorem: Every orthogonal matrix in  $O(2)$  is either a rotation or a reflection across a line.
- The set of  $n \times n$  real orthogonal matrices with determinant 1 is also closed under matrix multiplication and inversion. This set is denoted by  $SO(n)$ .

## Orthonormal basis - orthogonal matrix $V$

- ▶ Orthogonal transformations in 2D and 3D spaces. Orthogonal matrices can be described using rotations and reflections.
- ▶ Theorem: Every orthogonal matrix in  $O(2)$  is either a rotation or a reflection across a line.  
Proof: Wettel notes.
- ▶ ~~Givens rotation, Householder reflection~~
- ▶ ~~Reflection of a vector into another~~
- ▶ Gram–Schmidt orthogonalization: studied last semester.
- ▶ QR decomposition: Just like LU decomposition compactly encodes the row operations needed to bring a matrix to triangular form, QR decomposition encodes the results of the orthogonalization process. This decomposition plays a crucial role in both the least squares method and in solving eigenvalue problems.

## Orthonormal basis - orthogonal matrix VI

- ▶ Definition: (QR decomposition). Let  $\mathbf{A}$  be a full column-rank real matrix. The decomposition  $\mathbf{A} = \mathbf{QR}$  is called a QR decomposition (or reduced QR decomposition) if  $\mathbf{Q}$  is a semi-orthogonal matrix of the same size as  $\mathbf{A}$ , and  $\mathbf{R}$  is a square upper triangular matrix with positive elements on the diagonal.
- ▶ Theorem: (Existence and uniqueness of QR decomposition). Every real full column-rank matrix  $\mathbf{A}$  has a QR decomposition, i.e., there exist a semi-orthogonal matrix  $\mathbf{Q}$  and an upper triangular matrix  $\mathbf{R}$  with positive diagonal elements such that  $\mathbf{A} = \mathbf{QR}$ . This decomposition is unique.

Proof: Wettel notes.

- ▶ QR decomposition with primitive orthogonal transformations: Wettel notes.
- ▶ Optimal solution of a system of equations using QR decomposition: Wettel notes.
- ▶ Least squares using QR decomposition: Wettel notes.

# Spaces over the Complex Field and Finite Fields I

- ▶ Complex vectors and spaces, scalar product of complex vectors.
- ▶ Definition: (Adjugate of a complex matrix). The adjugate (or Hermitian transpose) of a complex matrix  $\mathbf{A}$  is the transpose of its elementwise conjugate. The adjugate of  $\mathbf{A}$  is denoted by  $\mathbf{A}^*$  or, after Hermite, by  $\mathbf{A}^H$ , thus  $\mathbf{A}^H = \overline{\mathbf{A}}^T$ .
- ▶ Definition: (Scalar product of complex vectors). For vectors  $\mathbf{z} = (z_1, z_2, \dots, z_n)$  and  $\mathbf{w} = (w_1, w_2, \dots, w_n)$  in  $\mathbb{C}^n$ , the scalar product is defined as

$$\mathbf{z} \cdot \mathbf{w} = \overline{z_1} w_1 + \overline{z_2} w_2 + \dots + \overline{z_n} w_n$$

which is a complex scalar. Its matrix product form is  $\mathbf{z} \cdot \mathbf{w} = \mathbf{z}^H \mathbf{w}$ .



## Spaces over the Complex Field and Finite Fields II

► Theorem: (Properties of the adjugate). Let **A** and **B** be complex matrices and let  $c$  be a complex number. Then:

1.  $(\mathbf{A}^H)^H = \mathbf{A}$ ,
2.  $(\mathbf{A} + \mathbf{B})^H = \mathbf{A}^H + \mathbf{B}^H$ ,
3.  $(c\mathbf{A})^H = \bar{c}\mathbf{A}^H$ ,
4.  $(\mathbf{AB})^H = \mathbf{B}^H\mathbf{A}^H$ .

► The properties of the adjugate immediately imply:

► Theorem: (Properties of complex scalar product). Let  $\mathbf{u}, \mathbf{v}, \mathbf{w} \in \mathbb{C}^n$  and let  $c \in \mathbb{C}$ . Then:

1.  $\mathbf{u} \cdot \mathbf{v} = \overline{\mathbf{v} \cdot \mathbf{u}}$ ,
2.  $\mathbf{u} \cdot (\mathbf{v} + \mathbf{w}) = \mathbf{u} \cdot \mathbf{v} + \mathbf{u} \cdot \mathbf{w}$ ,
3.  $(c\mathbf{u}) \cdot \mathbf{v} = c(\mathbf{u} \cdot \mathbf{v})$  and  $\mathbf{u} \cdot (c\mathbf{v}) = \bar{c}(\mathbf{u} \cdot \mathbf{v})$ ,
4.  $\mathbf{u} \cdot \mathbf{u} > 0$  if  $\mathbf{u} \neq \mathbf{0}$ , and  $\mathbf{u} \cdot \mathbf{u} = 0$  if  $\mathbf{u} = \mathbf{0}$ .

Proof: See Wettl notes.

## Spaces over the Complex Field and Finite Fields III

- ▶ Distinguished subspaces of complex matrices: see Wettl notes.
- ▶ Self-adjoint matrices: Just as the adjoint is the extension of the transpose (considering complex scalar product), the self-adjungate matrix is the extension of the symmetric matrix. A symmetric matrix is equal to its own transpose, while a self-adjungate matrix is equal to its own adjungate.
- ▶ A complex matrix  $\mathbf{A}$  is self-adjungate if  $\mathbf{A}^H = \mathbf{A}$ .
- ▶ Self-adjungate matrices are also called Hermitian matrices.
- ▶ The diagonal of a self-adjungate matrix contains only real numbers, as only real numbers are equal to their own conjugate.
- ▶ Every real symmetric matrix is self-adjungate, since real numbers are equal to their own conjugate. Since non-real complex numbers are not equal to their conjugates, a complex symmetric matrix is self-adjungate if and only if all its elements are real.

## Spaces over the Complex Field and Finite Fields IV

- ▶ Distance and orthogonal projection in complex spaces: The scalar product allows the definition of distance and orthogonality between complex vectors.
- ▶ The length or absolute value of a complex vector  $\mathbf{u} \in \mathbb{C}^n$  is  $|\mathbf{u}| = \sqrt{\mathbf{u} \cdot \mathbf{u}}$ ; the distance between two vectors equals the length of their difference: for vectors  $\mathbf{u}, \mathbf{v} \in \mathbb{C}^n$ ,  $d(\mathbf{u}, \mathbf{v}) = |\mathbf{u} - \mathbf{v}|$ . Two vectors are considered orthogonal if their scalar product is 0.
- ▶ Theorem: (Cauchy–Bunyakovsky–Schwarz inequality). For all  $\mathbf{u}, \mathbf{v} \in \mathbb{C}^n$ :



$$|\mathbf{u} \cdot \mathbf{v}| \leq |\mathbf{u}| |\mathbf{v}|.$$

Equality holds if and only if  $\mathbf{u}$  and  $\mathbf{v}$  are linearly dependent, i.e., one is a scalar multiple of the other.

## Spaces over the Complex Field and Finite Fields V

- ▶ The complex analogue of orthogonal matrices is the unitary matrix.
- ▶ Definition: (Unitary matrix). A square complex matrix  $\mathbf{U}$  is unitary if  $\mathbf{U}^H \mathbf{U} = \mathbf{E}$ .
- ▶ Just as in the case of orthogonal matrices, one can prove that a matrix  $\mathbf{U} \in \mathbb{C}^{n \times n}$  is unitary if any of the following equivalent conditions hold:
  1.  $\mathbf{U} \mathbf{U}^H = \mathbf{E}$ ,
  2.  $\mathbf{U}^{-1} = \mathbf{U}^H$ ,
  3. The column vectors of  $\mathbf{U}$  form an orthonormal basis with respect to the complex scalar product,
  4. The row vectors of  $\mathbf{U}$  form an orthonormal basis with respect to the complex scalar product,
  5.  $|\mathbf{U}\mathbf{x}| = |\mathbf{x}|$  for every vector  $\mathbf{x} \in \mathbb{C}^n$ ,
  6.  $\mathbf{U}\mathbf{x} \cdot \mathbf{U}\mathbf{y} = \mathbf{x} \cdot \mathbf{y}$ .

# Complex numbers are cool beautiful



# The End

Thank you for your attention!