



(KVXFI2BBNE, KVXFI21ONC, KVXFI6BBNE) Physics II.

Dr. Gábor FACSKÓ, PhD
Assistant Professor / Senior Research Fellow
facsko.gabor@uni-obuda.hu

Óbuda University, Faculty of Electrical Engineering, 1084 Budapest, Tavaszmező u. 17.
Wigner Research Centre for Physics, Department of Space Physics and Space Technology, 1121 Budapest, Konkoly-Thege Miklós út 29–33.
<https://wigner.hu/~facsko.gabor>

October 6, 2025

Planck Radiation Law I

- ▶ The Planck Radiation Law can be expressed mathematically as follows:

$$E(f, T) = \frac{8\pi hf^3}{c^3} \frac{1}{e^{\frac{hf}{kT}} - 1},$$

and

$$E(\lambda, T) = \frac{8\pi ch}{\lambda^5} \frac{1}{e^{\frac{hc}{\lambda kT}} - 1},$$

where c [m/s] is the speed of light in vacuum, f [1/s] is the frequency of the radiation, λ [m] is the wavelength, k is the Boltzmann constant, T [K] is the absolute temperature, and h is the Planck constant.

The Photoelectric Effect - Nobel Prize, Albert EINSTEIN, 1905

Foton $E = h \cdot f$

$E_0 = \frac{3}{2} kT$

$E' = E_0 + hf$

Az elektron elnyeli a fotont

Az elektron mozog a felület felé. Ez a mozgás ΔE energiát felemészthet.




Az elektron kilép a felületen. Ez $W_{ki} = e \cdot U$ energiába kerül.

Ha $\Delta E = 0$ és $E_0 \rightarrow 0$, akkor:

$$\frac{1}{2} m v_{max}^2 = h \cdot f - W_{ki}$$

Compton Scattering I

COMPTON-SZÓRÁS



Compton-formula:

Compton-hullámhossz:

$$\Delta\lambda = \frac{h}{m_0 c} \cdot (1 - \cos\theta)$$

$$\lambda_c = \frac{h}{m_0 c}$$

Exercises I

- ▶ See other documents
- ▶ Photoelectric Effect
- ▶ Compton Scattering
- ▶ etc.

The End

Thank you for your attention!