


ANSWER SCHEME
SESSION 2017/2018

NO	ANSWER SCHEME	MARKS
1	$3^{2x+1} - (16)3^x + 5 = 0$ $3(3^x)^2 - 16(3^x) + 5 = 0$ Let $m = 3^x$ $3m^2 - 16m + 5 = 0$ $(3m - 1)(m - 5) = 0$ $m = \frac{1}{3} \quad m = 5$ $3^x = 3^{-1} \quad 3^x = 5$ $x = -1 \quad x = 1.465$	K1 – 3^x J1 J1 K1 J1 J1
TOTAL		6 marks
2	$3 \log_9 x = (\log_3 x)^2$ $\frac{3 \log_3 x}{\log_3 9} = (\log_3 x)^2$ $\frac{3 \log_3 x}{\log_3 3^2} = (\log_3 x)^2$ $\frac{3 \log_3 x}{2} = (\log_3 x)^2$ let $u = \log_3 x$ $\frac{3u}{2} = (u)^2$ $2u^2 - 3u = 0$ $u(2u - 3) = 0$ $u = 0 \quad u = \frac{3}{2}$ $\log_3 x = 0 \quad \log_3 x = \frac{3}{2}$ $x = 1 \quad x = \sqrt{27}$ $x = 5.196$	K1 – same base K1 – change $9 = 3^2$ K1 – log rule J1 K1 - quadratic equation and equate with zero J1 – both K1 – either one to find x J1 J1
TOTAL		9 marks

NO	ANSWER SCHEME	MARKS
3(a)	$z = 2 + i$ $\frac{-}{z} - \frac{1}{z} = 2 - i - \frac{1}{2 - i}$ $= 2 - i - \left(\frac{1}{2 - i} \times \frac{2 + i}{2 + i} \right)$ $= 2 - i - \left(\frac{2 + i}{5} \right)$ $= \frac{5(2 - i) - 2 - i}{5}$ $= \frac{10 - 5i - 2 - i}{5}$ $= \frac{8 - 6i}{5}$ $= \frac{8}{5} - \frac{6}{5}i$	B1 K1 – multiply conjugate K1 – equalize denominator J1
3(b)	$\left \frac{-}{z} - \frac{1}{z} \right = \left \frac{8}{5} - \frac{6}{5}i \right $ $= \sqrt{\left(\frac{8}{5}\right)^2 + \left(-\frac{6}{5}\right)^2}$ $= 2$ $\left(\frac{-}{z} - \frac{1}{z} \right)^2 = \left(\frac{8}{5} - \frac{6}{5}i \right) \left(\frac{8}{5} - \frac{6}{5}i \right)$ $= \frac{64}{25} - 2 \left(\frac{48}{25} \right) i - \frac{36}{25}$ $= \frac{28}{25} - \frac{96}{25}i$ $\alpha + \beta i = \left \frac{-}{z} - \frac{1}{z} \right \left(\frac{-}{z} - \frac{1}{z} \right)^2$ $= 2 \left(\frac{28}{25} - \frac{96}{25}i \right)$ $= \frac{56}{25} - \frac{192}{25}i$ compare $\alpha = \frac{56}{25} \quad \beta = -\frac{192}{25}$	K1 - modulus J1 K1 - expansion J1 K1 – multiply 2 K1 – compare J1 J1
TOTAL		12 marks

NO	ANSWER SCHEME	MARKS																								
4(a)	$\frac{5}{x+3} - 1 \leq 0$ $\frac{5-x-3}{x+3} \leq 0$ $\frac{2-x}{x+3} \leq 0$ C.N: $x = -3, x = 2$ <table><tr><td></td><td>-3</td><td>2</td><td></td></tr><tr><td>Interval</td><td>$(-\infty, -3)$</td><td>$(-3, 2]$</td><td>$[2, \infty)$</td></tr><tr><td>Value x</td><td>-5</td><td>0</td><td>3</td></tr><tr><td>$2-x$</td><td>-</td><td>-</td><td>+</td></tr><tr><td>$x+3$</td><td>-</td><td>+</td><td>+</td></tr><tr><td>Conc</td><td> + </td><td>-</td><td> + </td></tr></table> $(-\infty, -3) \cup [2, \infty)$		-3	2		Interval	$(-\infty, -3)$	$(-3, 2]$	$[2, \infty)$	Value x	-5	0	3	$2-x$	-	-	+	$x+3$	-	+	+	Conc	+	-	+	K1 – equate denominator K1 – simplify K1 J1
	-3	2																								
Interval	$(-\infty, -3)$	$(-3, 2]$	$[2, \infty)$																							
Value x	-5	0	3																							
$2-x$	-	-	+																							
$x+3$	-	+	+																							
Conc	+	-	+																							
4(b)	$\left \frac{3x-2}{2x+3} \right > 2$ $\frac{3x-2}{2x+3} > 2 \qquad \text{or} \qquad \frac{3x-2}{2x+3} < -2$ $\frac{3x-2}{2x+3} - 2 > 0 \qquad \frac{3x-2}{2x+3} + 2 < 0$ $\frac{3x-2-2(2x+3)}{2x+3} > 0 \qquad \frac{3x-2+2(2x+3)}{2x+3} < 0$ $\frac{-x-8}{2x+3} > 0 \qquad \frac{7x+4}{2x+3} < 0$ C.N: $x = -8, x = -\frac{3}{2}$ <table><tr><td></td><td>-8</td><td>$-\frac{3}{2}$</td><td></td></tr><tr><td>Interval</td><td>$(-\infty, -8)$</td><td>$\left(-8, -\frac{3}{2}\right)$</td><td>$\left(-\frac{3}{2}, \infty\right)$</td></tr><tr><td>Value x</td><td>-9</td><td>-2</td><td>0</td></tr><tr><td>$-x-8$</td><td>+</td><td>-</td><td>-</td></tr><tr><td>$2x+3$</td><td>-</td><td>-</td><td>+</td></tr><tr><td>Conc</td><td>-</td><td> + </td><td>-</td></tr></table> $\left(-8, -\frac{3}{2}\right)$		-8	$-\frac{3}{2}$		Interval	$(-\infty, -8)$	$\left(-8, -\frac{3}{2}\right)$	$\left(-\frac{3}{2}, \infty\right)$	Value x	-9	-2	0	$-x-8$	+	-	-	$2x+3$	-	-	+	Conc	-	+	-	B1 - definition K1 – collect one side J1 - both K1 - method J1
	-8	$-\frac{3}{2}$																								
Interval	$(-\infty, -8)$	$\left(-8, -\frac{3}{2}\right)$	$\left(-\frac{3}{2}, \infty\right)$																							
Value x	-9	-2	0																							
$-x-8$	+	-	-																							
$2x+3$	-	-	+																							
Conc	-	+	-																							

NO	ANSWER SCHEME	MARKS
5(b)	$(g \circ f)(x) = g[(x-2)^2 + 1]$ $= \ln\left[\left((x-2)^2 + 1\right) + 1\right]$ $= \ln\left[(x-2)^2 + 2\right]$ $(g \circ f)(2) = \ln\left[(2-2)^2 + 2\right]$ $= \ln 2$ $= 0.693$	K1 – subs $f(x)$ J1 K1 – subs $x = 2$ J1
TOTAL		13 marks
6(a)	$g(x) = \frac{1}{2x-5}$ $2x-5 \neq 0$ $x \neq \frac{5}{2}$ $D_g : \left(-\infty, \frac{5}{2}\right) \cup \left(\frac{5}{2}, \infty\right)$ $R_g : (-\infty, 0) \cup (0, \infty)$	K1 J1 J1
6(b)	$g(x_1) = g(x_2)$ $\frac{1}{2x_1-5} = \frac{1}{2x_2-5}$ $2x_2-5 = 2x_1-5$ $2x_1 = 2x_2$ $x_1 = x_2$ $\therefore g(x) \text{ is one-to-one function. } \}$ $g[g^{-1}(x)] = x$ $\frac{1}{2[g^{-1}(x)]-5} = x$ $2[g^{-1}(x)]-5 = \frac{1}{x}$ $2[g^{-1}(x)] = \frac{1}{x} + 5$ $2[g^{-1}(x)] = \frac{1+5x}{x}$ $g^{-1}(x) = \frac{1+5x}{2x}$	K1 - state J1 K1 – using properties K1 – attempt to find inverse J1

NO	ANSWER SCHEME	MARKS
6(c)	 $g^{-1}(x) = \frac{1+5x}{2x}$ $g(x) = \frac{1}{2x-5}$	R1 – shape and label R1 – with pair R1 – all correct (asymptote, intercept)
6(d)	$g \circ g^{-1}(x) = \frac{1}{2\left(\frac{1+5x}{2x}\right) - 5}$ $= \frac{1}{\frac{1+5x}{x} - 5}$ $= \frac{x}{1+5x-5x}$ $= x$	K1 – to get $g \circ g^{-1}$ J1
TOTAL		13 marks
7(a)	$\lim_{x \rightarrow 2} \frac{x^3 - 8}{x^2 - 2x} = \lim_{x \rightarrow 2} \frac{(x-2)(x^2 + 2x + 4)}{x(x-2)}$ $= \lim_{x \rightarrow 2} \frac{x^2 + 2x + 4}{x}$ $= \frac{2^2 + 2(2) + 4}{4}$ $= 6$	K1 – factorize K1 - simplify J1 - answer
7(b)	$\lim_{x \rightarrow \infty} \sqrt{\frac{5x+7}{6x-5}} = \sqrt{\lim_{x \rightarrow \infty} \frac{5x+7}{6x-5}}$ $= \sqrt{\lim_{x \rightarrow \infty} \frac{\frac{5x}{x} + \frac{7}{x}}{\frac{6x}{x} - \frac{5}{x}}}$ $= \sqrt{\lim_{x \rightarrow \infty} \frac{5 + \frac{7}{x}}{6 - \frac{5}{x}}}$ $= \sqrt{\frac{5}{6}}$	K1 K1 J1
TOTAL		6 marks

NO	ANSWER SCHEME	MARKS
8	$y = e^{-2x} \sin 3x$ $u = e^{-2x} \quad v = \sin 3x$ $u' = -2e^{-2x} \quad v' = 3 \cos 3x$ $\frac{dy}{dx} = -2e^{-2x} \sin 3x + 3e^{-2x} \cos 3x$ $= e^{-2x} [3 \cos 3x - 2 \sin 3x]$ $u = e^{-2x} \quad v = -2 \sin 3x + 3 \cos 3x$ $u' = -2e^{-2x} \quad v' = -6 \cos 3x - 9 \sin 3x$ $\frac{d^2y}{dx^2} = (-2 \sin 3x + 3 \cos 3x)(-2e^{-2x}) + (-6 \cos 3x - 9 \sin 3x)(e^{-2x})$ $= e^{-2x} [4 \sin 3x - 6 \cos 3x + (-6 \cos 3x - 9 \sin 3x)]$ $= e^{-2x} [-5 \sin 3x - 12 \cos 3x]$ $\frac{d^2y}{dx^2} + 4 \frac{dy}{dx} + 13y$ $= e^{-2x} [-5 \sin 3x - 12 \cos 3x] + 4(e^{-2x} [-2 \sin 3x + 3 \cos 3x]) + 13(e^{-2x} \sin 3x)$ $= e^{-2x} [-5 \sin 3x - 12 \cos 3x - 8 \sin 3x + 12 \cos 3x + 13 \sin 3x]$ $= e^{-2x} (0)$ $= 0 \quad (\text{shown})$	K1 – differentiate J1 - $\frac{dy}{dx}$ K1 – differentiate J1 - $\frac{d^2y}{dx^2}$ K1 – substitute J1 - answer
TOTAL		6 marks
9	The condition for continuity of $f(x)$ at $x = a$ (a) $f(a)$ is defined. (b) $\lim_{x \rightarrow a} f(x)$ is exist. (c) $f(a) = \lim_{x \rightarrow a} f(x)$	K1J1
9(a)	At $x = 0$ $f(0) = \lim_{x \rightarrow 0^-} (n - 2 \cos x) = \lim_{x \rightarrow 0^+} (2 + mx^2)$ $\lim_{x \rightarrow 0^-} (n - 2 \cos x) = \lim_{x \rightarrow 0^+} (2 + mx^2)$ $n - 2 = 2$ $n = 4$ At $x = 2$ $f(2) = \lim_{x \rightarrow 2^-} (2 + mx^2) = \lim_{x \rightarrow 2^+} (m - x)$ $\lim_{x \rightarrow 2^-} (2 + mx^2) = \lim_{x \rightarrow 2^+} (m - x)$ $2 + m(2)^2 = m - 2$ $m = \frac{-4}{3}$	B1 K1 J1 B1 K1 J1

NO	ANSWER SCHEME	MARKS
9(b)	$m = -2, n = 4$ $f(x) = \begin{cases} 4 - 2\cos x, & x < 0 \\ 2 - 2x^2, & 0 \leq x < 2 \\ -2 - x, & x \geq 2 \end{cases}$ $f'(2^-) = \lim_{x \rightarrow 2^-} \frac{(2 - 2x^2) - (-4)}{x - 2}$ $= \lim_{x \rightarrow 2^-} \frac{-2x^2 + 6}{x - 2}$ $= +\infty \text{ (undefined)}$ $f'(2^+) = \lim_{x \rightarrow 2^+} \frac{(-2 - x) - (-4)}{x - 2}$ $= \lim_{x \rightarrow 2^+} \frac{-(x - 2)}{x - 2}$ $= -1$ Since $f'(2^-) \neq f'(2^+)$, therefore $f(x)$ is not differentiable at $x = 2$	K1 K1 J1 – both J1
TOTAL		12 marks
10(a)	$x^2 - 3y^2 = ae^{y-2x} + by - 6$ At the point (1,2) $(1)^2 - 3(2)^2 = ae^{2-2(1)} + b(2) - 6$ $a + 2b = -5 \quad \dots \quad (i)$ $2x - 6y \frac{dy}{dx} = ae^{y-2x} \frac{d}{dx}(y - 2x) + b \frac{dy}{dx}$ $2x - 6y \frac{dy}{dx} = ae^{y-2x} \left[\frac{dy}{dx} - 2 \right] + b \frac{dy}{dx}$ At the point (1,2), $\frac{dy}{dx} = 1$ $2(1) - [6(2)(1)] = ae^{2-2(1)} [1 - 2] + b(1)$ $-a + b = -10$ $a - b = 10 \quad \dots \quad (ii)$ (i) – (ii) $3b = -15$ $b = -5$ $a + 2(-5) = -5$ $a = 5$ $\therefore a = 5, \quad b = -5$	K1 -Substitute K1 – 1st equation K1 - differentiate K1 - substitute K1 – 2nd equation K1 – any method to find a and b J1, J1

NO	ANSWER SCHEME	MARKS
10(b)	$2x - 6y \frac{dy}{dx} = 5e^{y-2x} \left[\frac{dy}{dx} - 2 \right] - 5 \frac{dy}{dx}$ $2 - \left[6y \frac{d^2y}{dx^2} + 6 \left(\frac{dy}{dx} \right)^2 \right] = \left[\left(\frac{dy}{dx} - 2 \right) \left(5e^{y-2x} \left(\frac{dy}{dx} - 2 \right) \right) + \left(5e^{y-2x} \right) \frac{d^2y}{dx^2} \right] - 5 \frac{d^2y}{dx^2}$ At the point (1,2), $\frac{dy}{dx} = 1$ $2 - \left[6(2) \frac{d^2y}{dx^2} + 6(1)^2 \right] = \left[(1-2) \left(5e^{2-2(1)} (1-2) \right) + \left(5e^{2-2(1)} \right) \frac{d^2y}{dx^2} \right] - 5 \frac{d^2y}{dx^2}$ $2 - 12 \frac{d^2y}{dx^2} - 6 = 5 + 5 \frac{d^2y}{dx^2} - 5 \frac{d^2y}{dx^2}$ $-12 \frac{d^2y}{dx^2} = 9$ $\frac{d^2y}{dx^2} = -\frac{3}{4}$	K1 - differentiate K1 - substitute K1 - simplify J1
TOTAL		12 marks
11(a)	$f(x) = \frac{\ln(x-1)}{x-1}$ $u = \ln(x-1) \quad v = x-1$ $u' = \frac{1}{x-1} \quad v' = 1$ $f'(x) = \frac{(x-1) \left(\frac{1}{x-1} \right) - \ln(x-1)(1)}{(x-1)^2}$ $= \frac{1 - \ln(x-1)}{(x-1)^2}$ $f'(x) = 0 \quad f'(x) \text{ undefined}$ $\frac{1 - \ln(x-1)}{(x-1)^2} = 0$ $\ln(x-1) = 1 \quad x-1 = 0$ $x-1 = e \quad x = 1 \text{ (rejected because } x > 1)$ $x = e + 1$	K1 - differentiate K1 - 1st derivative K1 - equate with zero J1 - value of x

NO	ANSWER SCHEME	MARKS
11(a)	$u = 1 - \ln(x - 1) \quad v = (x - 1)^2$ $u' = -\frac{1}{x - 1} \quad v' = 2(x - 1)$ $f''(x) = \frac{(x - 1)^2 \left(-\frac{1}{x - 1} \right) - (1 - \ln(x - 1))(2(x - 1))}{(x - 1)^4}$ $= \frac{-(x - 1) - (1 - \ln(x - 1))(2(x - 1))}{(x - 1)^4}$ When $x = e + 1$ $f''(e + 1) = \frac{-(e + 1 - 1) - (1 - \ln(e + 1 - 1))(2(e + 1 - 1))}{(e + 1 - 1)^4}$ $= \frac{-e - (1 + \ln 1)(2e)}{e^4}$ $= \frac{-3e}{e^4}$ $= \frac{-3}{e^3} < 0 \quad (\text{maximum})$	K1 - differentiate K1 - 2nd derivative K1 - substitute J1
11(b)	When $x = e + 1$ $f(e + 1) = \frac{\ln(e + 1 - 1)}{e + 1 - 1}$ $= \frac{\ln e}{e}$ $= \frac{1}{e}$ So the exact coordinates of the maximum point is $\left(e + 1, \frac{1}{e} \right)$	K1 - substitute into $f(x)$ J1
11(c)	$\frac{d^2y}{dx^2} = 0$ $\frac{-(x - 1) - (1 - \ln(x - 1))(2(x - 1))}{(x - 1)^4} = 0$ $-(x - 1) - (1 - \ln(x - 1))(2(x - 1)) = 0$ $-(x - 1)[1 + 2(1 - \ln(x - 1))] = 0$ $-(x - 1)[3 - 2\ln(x - 1)] = 0$	K1

NO	ANSWER SCHEME	MARKS
11(c)	$3 - 2\ln(x - 1) = 0$ $2\ln(x - 1) = 3$ $-(x - 1) = 0$ $x - 1 = 0$ $x = 1$ $\ln(x - 1) = \frac{3}{2}$ $x - 1 = e^{\frac{3}{2}}$ $x = e^{\frac{3}{2}} + 1$ Since $x \neq 1$, therefore $x = e^{\frac{3}{2}} + 1$.	J1
TOTAL		12 marks
12(a)	$x = 3t - \frac{1}{t} \qquad y = t + \frac{3}{t}$ $\frac{dx}{dt} = 3 + \frac{1}{t^2} \qquad \frac{dy}{dt} = 1 - \frac{3}{t^2}$ $= \frac{3t^2 + 1}{t^2} \qquad = \frac{t^2 - 3}{t^2}$ $\frac{dy}{dx} = \frac{t^2 - 3}{t^2} \times \frac{t^2}{3t^2 + 1}$ $= \frac{t^2 - 3}{3t^2 + 1} \quad (\text{shown})$ $u = t^2 - 3 \qquad v = 3t^2 + 1$ $u' = 2t \qquad v' = 6t$ $\frac{d}{dt} \left(\frac{t^2 - 3}{3t^2 + 1} \right) = \frac{(3t^2 + 1)(2t) - (t^2 - 3)(6t)}{(3t^2 + 1)^2}$ $= \frac{6t^3 + 2t - 6t^3 + 18t}{(3t^2 + 1)^2}$ $= \frac{20t}{(3t^2 + 1)^2}$	K1 – differentiate K1 – chain rule J1 K1 K1 – differentiate $\frac{dy}{dx}$ J1

NO	ANSWER SCHEME	MARKS
12(a)	$\frac{d^2y}{dx^2} = \frac{d}{dt} \left(\frac{t^2 - 3}{3t^2 + 1} \right) \times \frac{t^2}{3t^2 + 1}$ $= \frac{20t}{(3t^2 + 1)^2} \times \frac{t^2}{3t^2 + 1}$ $= \frac{20t^3}{(3t^2 + 1)^3}$	K1 – 2nd derivative J1
12(b)	$\frac{dy}{dx} = \frac{t^2 - 3}{3t^2 + 1}$ $= \frac{1}{3} + \frac{-\frac{10}{3}}{3t^2 + 1}$ $= \frac{1}{3} + \frac{-10}{3(3t^2 + 1)}$ $t^2 > 0$ $3t^2 > 0$ $3t^2 + 1 > 0 + 1$ $3t^2 + 1 > 1$ $3(3t^2 + 1) > 3$ $3 < 3(3t^2 + 1) < \infty$ $0 < \frac{1}{3(3t^2 + 1)} < \frac{1}{3}$ $0 < \frac{10}{3(3t^2 + 1)} < \frac{10}{3}$ $-\frac{10}{3} < -\frac{10}{3(3t^2 + 1)} < 0$ $\frac{1}{3} - \frac{10}{3} < \frac{1}{3} - \frac{10}{3(3t^2 + 1)} < \frac{1}{3} - 0$ $-3 < \frac{1}{3} - \frac{10}{3(3t^2 + 1)} < \frac{1}{3}$ $-3 < \frac{dy}{dx} < \frac{1}{3}$	K1 – using long division J1 K1 K1 K1 K1 J1
TOTAL		15 marks