

Algebras for a Functor

Gašper Golob in Nejc Zajc

Project presentation, 26.5.2022

Motivation

Modelling inductive types.

F-Algebras

category C , endofunctor $F: C \rightarrow C$

F-Algebras

category \mathcal{C} , endofunctor $F: \mathcal{C} \rightarrow \mathcal{C}$

$$FA \xrightarrow{\alpha} A$$

F-Algebras

category C , endofunctor $F: C \rightarrow C$

$$\begin{array}{ccc} FA & \xrightarrow{\alpha} & A \\ \downarrow Ff & & \downarrow f \\ FB & \xrightarrow{\beta} & B \end{array}$$

Initial Objects

Such an object I , that for every object X , there exist a **unique** morphism $I \rightarrow X$.

Lambek Lemma

Lemma (Lambek)

If $I = (A, \alpha)$ is an initial algebra, then A is isomorphic to FA via α .

Lambek Lemma

Lemma (Lambek)

If $I = (A, \alpha)$ is an initial algebra, then A is isomorphic to FA via α .

$$\begin{array}{ccc} FA & \xrightarrow{\alpha} & A \\ \downarrow Fi & & \downarrow i \\ F(FA) & \xrightarrow{F\alpha} & FA \end{array}$$

Polynomial Functor

- project only defines them on **Sets**
- $PX = \sum_{i \in I} X^{A_i}$, for $A : I \rightarrow \mathbf{Set}$
- natural numbers from $PX = 1 + X$

Initial algebra for polynomial functors

- Tree has a constructor $\text{Node } \sum_{i \in I} \text{Tree}^{A_i}$
- initial object is the F-algebra of the Tree

$$\begin{array}{ccc} \sum_{i \in I} \text{Tree}^{A_i} & \xrightarrow{\alpha} & \text{Tree} \\ Pf \downarrow & & \downarrow f \\ \sum_{i \in I} B^{A_i} & \xrightarrow{\beta} & B \end{array}$$

Future work

- implement presentation of natural numbers with F-algebras
- implement presentation of lists with F-algebras
- generalize polynomial functors
- generalize existence of initial algebras

Problems in Implementation

Problems in Implementation

```
; _°_ = λ f g → record {  
  f = F-Algebra-Morphism.f f ° F-Algebra-Morphism.f g ;  
  commutes = glue {! 0!} (F-Algebra-Morphism.commutes f)  
  (F-Algebra-Morphism.commutes g)}
```

Problems in Implementation

```
; _°_ = λ f g → record {
  f = F-Algebra-Morphism.f f ° F-Algebra-Morphism.f g ;
  commutes = glue {! 0!} (F-Algebra-Morphism.commutes f)
  (F-Algebra-Morphism.commutes g)}
```

Initial $/ \Rightarrow \text{record } \{ \perp, \perp\text{-is-initial} \} \Rightarrow \text{record } \{ A, \alpha \} .$

Sources

Awodey, Steve (2010). *Category Theory*. 2nd. USA: Oxford University Press, Inc. ISBN: 0199237182.

nLab authors (May 2022). *initial algebra of an endofunctor*. URL: <http://ncatlab.org/nlab/show/initial%20algebra%20of%20an%20endofunctor>.