

Triangulation (computing depth)

Given the intrinsic parameters, the projections of scene point on the two image sensors are:

$$\begin{bmatrix} u_l \\ v_l \\ 1 \end{bmatrix} = \begin{bmatrix} f_x^{(l)} & 0 & o_x^{(l)} & 0 \\ 0 & f_y^{(l)} & o_y^{(l)} & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix} \begin{bmatrix} x_l \\ y_l \\ z_l \\ 1 \end{bmatrix}$$

left camera imaging equation

$$\begin{bmatrix} u_r \\ v_r \\ 1 \end{bmatrix} = \begin{bmatrix} f_x^{(r)} & 0 & o_x^{(r)} & 0 \\ 0 & f_y^{(r)} & o_y^{(r)} & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix} \begin{bmatrix} x_r \\ y_r \\ z_r \\ 1 \end{bmatrix}$$

right camera imaging equation

We also know the relative position and orientation between the two cameras:

$$\begin{bmatrix} x_l \\ y_l \\ z_l \\ 1 \end{bmatrix} = \begin{bmatrix} r_{11} & r_{12} & r_{13} & t_x \\ r_{21} & r_{22} & r_{23} & t_y \\ r_{31} & r_{32} & r_{33} & t_z \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x_r \\ y_r \\ z_r \\ 1 \end{bmatrix}$$

$$\begin{bmatrix} u_l \\ v_l \\ 1 \end{bmatrix} = \begin{bmatrix} f_x^{(l)} & 0 & o_x^{(l)} & 0 \\ 0 & f_y^{(l)} & o_y^{(l)} & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix} \begin{bmatrix} r_{11} & r_{12} & r_{13} & t_x \\ r_{21} & r_{22} & r_{23} & t_y \\ r_{31} & r_{32} & r_{33} & t_z \\ 0 & 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x_r \\ y_r \\ z_r \\ 1 \end{bmatrix}$$

$\tilde{u}_l = P_l \tilde{x}_r$

Substitute in the left imaging equation

$$\begin{bmatrix} u_r \\ v_r \\ 1 \end{bmatrix} = \begin{bmatrix} f_x^{(r)} & 0 & o_x^{(r)} & 0 \\ 0 & f_y^{(r)} & o_y^{(r)} & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix} \begin{bmatrix} x_r \\ y_r \\ z_r \\ 1 \end{bmatrix}$$

$\tilde{u}_r = r_{int_r} \tilde{x}_r$

Keep the right imaging equation as it is

$$\begin{bmatrix} u_l \\ v_l \\ 1 \end{bmatrix} = \begin{bmatrix} p_{11} & p_{12} & p_{13} & p_{14} \\ p_{21} & p_{22} & p_{23} & p_{24} \\ p_{31} & p_{32} & p_{33} & p_{34} \end{bmatrix} \begin{bmatrix} x_r \\ y_r \\ z_r \\ 1 \end{bmatrix}$$

$\tilde{u}_l = P_l \tilde{x}_r$

Known Unknown

$$\begin{bmatrix} u_r \\ v_r \\ 1 \end{bmatrix} = \begin{bmatrix} m_{11} & m_{12} & m_{13} & m_{14} \\ m_{21} & m_{22} & m_{23} & m_{24} \\ m_{31} & m_{32} & m_{33} & m_{34} \end{bmatrix} \begin{bmatrix} x_r \\ y_r \\ z_r \\ 1 \end{bmatrix}$$

$\tilde{u}_r = M_r \tilde{x}_r$

Known Unknown

Rearranging the terms :

$$\begin{array}{|c|} \hline \begin{array}{ccc} u_r m_{31} - m_{11} & u_r m_{32} - m_{12} & u_r m_{33} - m_{13} \\ v_r m_{31} - m_{21} & v_r m_{32} - m_{22} & v_r m_{33} - m_{23} \\ u_l p_{31} - p_{11} & u_l p_{32} - p_{12} & u_l p_{33} - p_{13} \\ u_l p_{31} - p_{21} & v_l p_{32} - p_{22} & v_l p_{33} - p_{23} \end{array} \\ \hline \end{array} \quad \begin{array}{|c|} \hline \begin{array}{c} x_r \\ y_r \\ z_r \end{array} \\ \hline \end{array} = \begin{array}{|c|} \hline \begin{array}{c} m_{14} - m_{34} \\ m_{24} - m_{34} \\ p_{14} - p_{34} \\ p_{24} - p_{34} \end{array} \\ \hline \end{array}$$

$A_{4 \times 3}$ (known) x_r (Unknown) $b_{4 \times 1}$ (known)

Find least squares solution using pseudo-inverse :

$$A x_r = b$$

$$A^T A x_r = A^T b$$

$$x_r = (A^T A)^{-1} A^T b$$

we repeat this for every pair of corresponding points in the left and right image. This gives us the complete 3-dimensional depth map of the scene.