

Explicit Scale Simulation for analysis of RNA-sequencing ~~count data~~ with ALDEx2

Gregory B. Gloor¹

Michelle Pistner Nixon²

Justin D. Silverman^{~~3,4,5~~2,3,4,5}

April 14, 2025

¹Department of Biochemistry, University of Western Ontario ²Department of Population Health Sciences, Geisinger, Danville, PA ³Department of Medicine, Pennsylvania State University ⁴Institute for Computational and Data Science, Pennsylvania State University ⁵Department of Statistics, Pennsylvania State University

Abstract

In high-throughput sequencing (HTS) studies, sample-to-sample variation in sequencing depth is driven by technical factors, and not by variation in the scale (size) of the biological system. Typically a statistical normalization removes unwanted technical variation in the data or the parameters of the model to enable differential abundance analyses. We recently showed that all normalizations make implicit assumptions about the unmeasured system scale and that errors in these assumptions can dramatically increase false positive and false negative rates. We demonstrated that these errors can be mitigated by accounting for uncertainty using a *scale model*, which we integrated into the ALDEx2 R package. This article provides new insights focusing on the application to transcriptomic analysis. We provide transcriptomic case studies demonstrating how scale models, rather than traditional normalizations, can reduce false positive and false negative rates in practice while enhancing the transparency and reproducibility of analyses. These scale models replace the need for dual cutoff approaches often used to address the disconnect between practical and statistical significance. We demonstrate the utility of scale models built based on known housekeeping genes in complex metatranscriptomic datasets. Thus this work provides guidance on how to incorporate scale into transcriptomic data sets.

Introduction

High-throughput sequencing (HTS) is a ubiquitous tool used to explore many biological phenomenon such as gene expression (single-cell sequencing, RNA-sequencing, meta-transcriptomics), microbial community composition (16S rRNA gene sequencing, shotgun metagenomics) and differential enzyme activity (selex, CRISPR killing). HTS proceeds by taking a sample from the environment, making a library, multiplexing (merging) multiple libraries together, and then applying a sample of the multiplexed library to a flow cell. Each of these steps is a compositional sampling step as only a fixed-size subsample of nucleic acid is carried over to subsequent steps. Thus, with each sampling step the connection between the actual ~~number of molecules in size of~~ the sampled DNA pool and the ~~environmental~~-scale (e.g., ~~total number of molecules~~~~size~~, microbial load, or total gene expression) of the measured biological system is degraded or lost. In the end, the information contained in the data relates only to relative abundances and has an arbitrary scale imposed by the sequencing process (1–3). Researchers can use modified experimental protocols (4–7) or machine learning methods (8) to uncover the biological variation in scale. However, wet-lab protocols only provide information on the size of the data downstream of the step in the sample preparation protocol where the intervention was made and introduce an additional source of variation that must be accounted for (5). The computational methods developed for microbiome analysis have low correlation with the actual scale but are useful (8). In short, the disconnect between sample-to-sample variation in sequencing depth and biological variation in scale remains an open challenge.

The analysis of HTS data suffers from several known problems that can be traced, in whole or in part, to misspecification of scale in the output data. The first issue is poor control of the false discovery rate (FDR) (4–8, 9–13), exhibited as dataset-dependent FDR control which is observed as a disconnect between statistical and biological significance (8, 9–14). In current practice, this issue is addressed by a dual-filtering method, whereby both a low p-value (or equivalently a low q-value following FDR correction (10–15)) and a large difference between groups is used to identify interesting transcripts or genes of interest for follow-up analysis (9, 11–14, 16). This double-filtering approach is graphically exemplified by the volcano plot (11–16), but is known to not appropriately control the FDR (12, 13). Since there is no standard way of determining what fold-change cutoff should be used researchers have unlimited degrees of freedom which is known to lead to unreliable inference (14). In particular Li et al. (8) used patient or clinical-derived transcriptome datasets and found that many methods suffer from an extremely high false positive rate (17, 18). The second issue is poor performance when analyzing data where the mean change between groups is non-zero (3). Such asymmetric data can arise when a gene set is expressed in one group but not the other, or when one group contains different gene content from the other. This type of data frequently arises in in-vitro selection experiments (SELEX), transcriptome analysis, and microbiome analysis (15–19). The third issue is that the actual scale of the environment is often a major confounding variable during analysis (3, 16). This has long been known in transcriptome analysis and was a major driver for the development of normalization factors (17) and the use of molecular spike-ins to provide a reference set of known counts (18–21) to estimate scale. While often useful, spike-in methods only provide information downstream of the step in the sample preparation protocol where the intervention was made and introduce an additional source of variation that must be accounted for (19). In the microbiome field, a recent landmark paper showed that biological scale was a major unacknowledged confounder in many human analyses (16). These authors built a machine learning model to uncover the biological variation in scale and including this information was useful despite exhibiting only a modest correlation (estimated to be 0.6) with the scale of data external to the training set (16). The final issue is that all the above these problems become more pronounced as more samples are collected; that is, more information results in optimizing for a precise but inaccurate a worsening of the accuracy of the analysis (3, 8, 22–23, 20).

The four problems were recently shown by Nixon et al. (3) to be a result of a mismatch between the underlying size or scale of the system and the assumptions of the normalizations used for the analysis of HTS. Biological variation in scale often represents an important unmeasured confounder in HTS analyses (15, 16, 19, 23–21). For example, cells transformed by the cMyc oncogene have about 3 times the amount of mRNA and about twice the rRNA content than non-transformed cells (24–22), and this dramatically skews transcriptome analysis unless spike-in approaches are used (19–5). In addition, wild-type and mutant strains of cell lines, yeast or bacteria have different growth rates and RNA contents under different conditions, which affect our ability to identify truly differentially abundant genes (25–27–23–25). As another example, the total bacterial load of the vaginal microbiome differs by 1–2 orders of magnitude in absolute abundance between the healthy and bacterial vaginosis states (28–26), and the cell and RNA composition between these states is dramatically different (29, 30–27, 28). Thus, a full description of any of these systems includes both relative change (composition) and absolute abundance (scale). Current methods access only the compositional information yet make implicit assumptions about the scale (22–20).

Recently, Nixon et al. (3) showed that the challenge of non-biological variation in sequencing depth could be explained be viewed as a problem of partially-identified models. They showed that all normalizations make some assumption about scale but these implicit assumptions are often inappropriate and difficult to interpret. As a result, different normalizations This causes different normalizations to provide different outputs when applied to the same dataset (4, 6, 17, 31, 32–9, 11, 29–31). Intuitively, normalizations in widespread use assume that either all samples have the same scale, e.g. proportions, rarefaction (33–32), RPKM (34, 35–33, 34), etc; or that a subset of features in one sample can be chosen as a reference to which the others are scaled e.g. the TMM (36), the LVHA (15) qPCR (37, 35), or LVHA (19) or the additive log-ratio (38–36); or that different sub-parts of each sample maintain a constant scale across samples e.g. the RLE (39–37); or that the geometric mean of the parts is appropriate e.g. the CLR (40–38) and its derivatives. Notably when the assumption of similar scale across samples is not violated, or violated only weakly, any method will provide a reasonable analysis. The problem arises when this assumption is violated, in which case all tools fail without warning.

Mismatches between estimated scale and true scale lead to poor estimation of high throughput sequencing data. All normalizations used for differential abundance analysis make some strict assumption about the scale of the environment as shown in line 1. In this example all normalizations produce a biased estimate of the environmental scale, but estimate C is the closest to the truth. Adding uncertainty to normalization C as represented by the distribution in the line 2 leads to less bias and now includes the actual environmental scale in the assumption. As shown in line 3 in some cases it may be useful to adjust the centre of the scale uncertainty estimate if the initial normalizations give very poor estimates of the underlying environment.

The original scale-naive ~~The original naive~~ ALDEx2 (41,39) model unwittingly made a strict assumption about scale through the CLR normalization and we found that the CLR was very sensitive to violations of the assumption of scale identity between groups (15). Moreover, when the assumption of identity was not true the CLR used by ALDEx2 (3). This assumption was often close enough to the true value to be useful, but was not always the a good estimate and could be outperformed by other normalizations in simulation studies (42). As illustrated graphically in Figure 1, (40). Nixon et al. (3) showed through simulation that introducing uncertainty in the scale assumptions, and in extreme cases altering the location of the scale assumption, that better scale assumptions resulted in more reproducible data analysis including better control of both false positive and false negative results. We recently modified ALDEx2 to explicitly model uncertainty in the scale over a range of reasonable normalization parameters, and showed significant improvements in performance in microbiome and in-vitro selection experiments (22) and in a vaginal metatranscriptome analysis (43,20). Here, we briefly review these modifications and show how adding scale uncertainty can greatly improve modeling in transcriptome and meta-transcriptome datasets to provide substantially more robust and reproducible results.

Implementation

Formal and expanded descriptions of the concepts that follow are given in (3, 22,20). To be concrete, we let \mathbf{Y} denote the *measured* $D \times N$ matrix of sequence counts with elements \mathbf{Y}_{dn} indicating the number of measured DNA molecules mapping to feature d (e.g., a taxon, transcript or gene) in sample n . Likewise, we can denote \mathbf{W}_{dn} as the *true* amount of class d in the biological system from which sample n was obtained. We can think of \mathbf{W} as consisting of two parts, the scale $\mathbf{W}^{Tot} \mathbf{W}^\perp$ (e.g., totals) and the composition $\mathbf{W}^{Comp} \mathbf{W}^\parallel$ (i.e., proportions). That is, $\mathbf{W}^{Tot} \mathbf{W}^\perp$ is a N -vector with elements $\mathbf{W}_n^{Tot} = \sum_d \mathbf{W}_{dn}$ while $\mathbf{W}^{Comp} \mathbf{W}^\perp = \sum_d \mathbf{W}_{dn}$ while \mathbf{W}^\parallel is a $D \times N$ matrix with elements $\mathbf{W}_{dn}^{Comp} = \mathbf{W}_{dn} / \mathbf{W}_n^{Tot}$ $\mathbf{W}_{dn}^\parallel = \mathbf{W}_{dn} / \mathbf{W}_n^\perp$. Note that with these definitions \mathbf{W} can be written as the element-wise combination of scale and composition: $\mathbf{W}_{dn} = \mathbf{W}_{dn}^{Comp} \mathbf{W}_n^{Tot} \mathbf{W}_{dn}^\perp = \mathbf{W}_{dn}^\parallel \mathbf{W}_n^\perp$, or as the logarithm $\log \mathbf{W}_{dn} = \log \mathbf{W}_{dn}^{Comp} + \log \mathbf{W}_n^{Tot} \log \mathbf{W}_{dn}^\perp = \log \mathbf{W}_{dn}^\parallel + \log \mathbf{W}_n^\perp$.

Many of the normalizations used in widespread use in tools such as DESeq2 (44,41), edgeR (36,35), metagenomeSeq (45,42) ALDEx2 (46,43) can be stated as ratios of the form $\hat{\mathbf{W}}_{dn} \approx \mathbf{Y}_{dn} / f(\mathbf{Y})$, where the denominator is determined by some function of the observation. We use the ~~hat notation~~ $\hat{\mathbf{W}}$ notation to indicate that the output is an estimate of the true value. The technical variation in sequencing depth, which is often called “library size” has no relationship with the actual number of molecules in the sampled environment (1). In other words $(\mathbf{Y}_n^{Tot} = \sum_d \mathbf{Y}_{dn})$ the $(\mathbf{Y}_n^\perp = \sum_d \mathbf{Y}_{dn})$ implies that observed data \mathbf{Y} provides us with information about the system composition $\mathbf{W}^{Comp} \mathbf{W}^\parallel$ but little to no information in the system scale $\mathbf{W}^{Tot} \mathbf{W}^\perp$ (Lovell et al. 2011).

Adding Scale Uncertainty in ALDEx2

The ALDEx2 R package (41, 46,39, 43) is a general purpose toolbox to model the uncertainty of HTS data and to use that model to estimate the significance of the underlying LFC (log-fold change) significance. At a high-level, ALDEx2 has three connected components to estimate the uncertainty inherent in HTS datasets. First, the tool accounts for the uncertainty of the sequencing counts using Dirichlet multinomial sampling to build a probabilistic model of the data; i.e., $\hat{\mathbf{W}}^{Comp} \approx \text{Dir}(\mathbf{Y}) \hat{\mathbf{W}}^\parallel \approx \text{Dir}(\mathbf{Y})$. Secondly, ALDEx2 uses the centred log-ratio transformation to scale the data (41). It was this step that was modified (39). However,

this step was modified recently to account for scale uncertainty and misspecification ~~(22)–(20)~~ via a scale model, explained with more details in (3, 2220) and summarized in the next paragraph. Finally, a standard null-hypothesis test and a non-parametric estimate of mean standardized difference are used to report on the finite sample variation. These sources of uncertainty and variation are combined via reporting the expected values from a Monte-Carlo simulation framework. For simplicity, we use the term ‘difference’ to refer to the absolute difference between groups, and ‘dispersion’ to refer to the within-condition difference or pooled variance as defined in (4139). These are calculated on a \log_2 scale. For more details on ALDEx2 see (3, 22, 41, 4620, 39, 43).

Scale models ~~were~~ can be incorporated into ALDEx2, turning the ALDEx2 model into a specialized type of statistical model which Nixon et al. (3) term a *Scale Simulation Random Variable* (SSRV). To do this, Nixon et al. (3) generalized the concept of normalizations by introducing the concept of a *scale model* to account for potential error in the centred log-ratio normalization step. They did this by including a model for $\hat{\mathbf{W}}_n^{Tot} \hat{\mathbf{W}}_n^\perp$. The CLR normalization used by ALDEx2 makes the assumption $\hat{\mathbf{W}}_n^{Tot} = 1/G_n \hat{\mathbf{W}}_n^\perp = 1/G_n$, where G_n is the geometric mean of ~~the counts (or the corresponding proportions) of each part in~~ sample n , which while being a random variable, is essentially constant across each Monte-Carlo replicate, but that differs between samples. With this modification, ALDEx2 can be generalized by considering probability models for the scale $\hat{\mathbf{W}}_n^{Tot} \hat{\mathbf{W}}_n^\perp$ that have mean $1/G_n$. For example, the following scale model generalizes the CLR:

$$\log \hat{\mathbf{W}}_n^{Tot \perp} = -\log G_n + \Lambda x_n \quad \Lambda \sim N(\mu, \gamma^2)$$

~

This formulation is quite flexible (3, 2220). In the simple or ‘default’ configuration, $\mu = 0$ and γ is a tunable parameter drawn from a log-Normal distribution(3). Adding scale uncertainty with the γ parameter ~~(as shown in Figure 1:1)~~ controls only the degree of uncertainty of the CLR assumption for the x_n binary condition indicator (e.g., $x_n = 1$ denotes case and $x_n = 0$ denotes control). In the advanced or ‘informed’ configuration, μ takes different values for each group and controls the location of the LFC assumption ~~(as shown in Figure 1:2)~~; combining μ with a γ estimate allows for uncertainty in both the location and the scale. ~~In the manuscript where the idea of scale was original derived, Nixon et al. (3) conducted extensive simulations showing that~~ An example of both the default and informed approaches ~~exhibit~~ is given for a microbiome dataset in (20) showing increased sensitivity and specificity ~~(22). In this report. Here we show that these approaches also work well to control the FDR in transcriptome and metatranscriptome datasets. We also provide some additional insights into how this is achieved.~~ These modifications are instantiated in ALDEx2 which is the first software package designed for SSRV-based inference.

Results

Adding scale uncertainty replaces the need for dual significance cutoffs.

Gierliński et al. (4744) generated a highly replicated yeast transcriptome dataset to compare gene expression between a wild-type strain and a *snf2* gene knockout, $\Delta snf2$. This dataset ~~of 86 samples (44 and 42 per group) is an example of technical growth replicate experiments common in the literature. The dataset~~ was used to test several RNA-seq tools for their power to detect the set of differentially abundant transcripts identified in the full dataset when the data was subset (914). In this ~~original~~ study each tool had its own ‘gold standard’ set of transcripts with different tools identifying between between 65% to >80% of all transcripts as being significantly different. Since the majority of transcripts were significantly different, the authors suggested that it was more appropriate to apply a dual cutoff composed of both a Benjamini-Hochberg (4845) corrected p-value (q-value) plus a difference cutoff to limit the number of identified transcripts to a much smaller fraction of the total. ~~In other datasets,~~ Nixon et al, (2220) showed that adding even a small amount of scale uncertainty with ALDEx2 dramatically reduced the number of significant transcripts identified, removing the need for the dual cutoff approach in this dataset and others. Below we include an intuitive explanation of why and how incorporating scale uncertainty achieved this outcome using a setting of $\gamma = 0.5$. The approach is in line with the recommendations of (20) and gives results comparable to those proposed in (14).

Volcano and effect plots for unscaled and scaled transcriptome analysis. ALDEx2 was used to conduct a differential expression (DE) analysis on the yeast transcriptome dataset. The results were plotted to show the relationship between difference and dispersion using effect plots or difference and the q-values using volcano plots. Panels A,C are for the naive analyses, and Panels B,D are for the default analyses that include scale uncertainty. Each point represents the values for one transcript, with the color indicating if that transcript was significant in the both analyses (red) or in the naive analysis only (orange). Points in grey are not statistically significantly different under any condition. The horizontal dashed lines represent a \log_2 difference of ± 1.4 .

We start with the assumption that not all statistically significant differences are biologically relevant (4946), and that ~~a result where the majority of transcripts are significant such a large number of significantly different transcripts~~ breaks the necessary assumption for DA/DE expression that most parts be invariant (31). ~~Transcriptomic analysis 30). As noted, transcriptomics commonly uses a dual cutoff approach that is~~ graphically exemplified by volcano plots (9, 1114, 16). Using either DESeq2 or ALDEx2, a majority of transcripts are statistically significantly different between groups with a q-value cutoff of ≤ 0.05 ; i.e. 4636 (79%, DESeq2) or 4172 (71%, ALDEx2) of the 5891 transcripts. These values are in line with those observed by (914). Such large numbers of statistically significant transcripts seems biologically unrealistic. That 118 transcripts are identified by ALDEx2 and not DESeq2, while DESeq2 identifies 582 transcripts that ALDEx2 does not, suggests that the choice of normalization plays a role in which results are returned as significant and that some, if not the majority, are driven by technical differences in the analysis (8, 31) ~~or are false positives~~ (13, 30).




Figure 1: ~~Volcano and effect plots for unscaled and scaled transcriptome analysis. ALDEx2 was used to conduct a differential expression (DE) analysis on the yeast transcriptome dataset. The results were plotted to show the relationship between difference and dispersion using effect plots or difference and the q-values using volcano plots. Panels A,C are for the naive analyses, and Panels B,D are for the default analyses that include scale uncertainty. Each point represents the values for one transcript, with the color indicating if that transcript was significant in the both analyses (red) or in the naive analysis only (orange). Points in grey are not statistically significantly different under any condition. The horizontal dashed lines represent a \log_2 difference of ± 1.4 .~~

The Volcano plots in Figure 2-1 A and B show that adding scale uncertainty increases the minimum q-value and increases the concordance between the q-value and the difference between groups (compare panels A and B). The effect plots (5047) in Figure 2C-1C shows that the majority of significant transcripts (red, orange) have negligible differences between groups and very low dispersion. We suggest that this low dispersion is driven by the experimental design which is actually a technical wet lab replication rather than a true biological replication design (4744). Scale uncertainty can be incorporated using the `gamma` parameter that controls the amount of noise added to the CLR mean assumption when we call either `aldex()`, or `aldex.clr()`. Figure 2-1 B,D shows that setting $\gamma = 0.5$ ~~now~~ results in 205 which is far fewer ~~statistically~~ significant transcripts than in the naive analysis and we observe that the minimum dispersion increases from 0.12 ($\gamma = 0$) to 0.67 ($\gamma = 0.5$).

It is common practice to use a dual-cutoff by choosing transcripts based on a thresholds for both q-values and fold-changes (9), ~~and these were first proposed for microarray experiments through volcano plots (1114, 16). Note that there is considerable variation in recommended cutoff values(9), and that this controversy has persisted ever since fold-change was suggested (51). Unfortunately, universal cutoff fold-change values cannot be identified in part because different tools have intrinsically different variance in their \log_2 fold change ranges (52). This has led to the widespread practice of applying a post-hoc fold-change cutoff to~~




Figure 2: Adding scale uncertainty changes the dispersion distribution. Panel A shows a plot of the expected value for relative abundance vs the expected value for the pooled dispersion as output by `aldex.effect`. The dashed horizontal lines show the median value for the features with a `rAbundance` between -0.5 and 0.5, and the light colored lines are lowess lines of fit through the center of mass of the data. Panel B plots the dispersion difference between $\gamma = 1$ and $\gamma = 0$; note the non-linear relationship that highlights the rotation that is evident in Panel A. The colored lines indicate the lowess line of fit through the centre of mass of the plot for the various populations of points. The grey line is the total population and shows the difference Δ , the red line is the population of significant transcripts (*) with $\gamma = 0$, the orange line is the population of significant transcripts with a difference threshold (T) of about $\pm 2^{1.4}$ -fold change, the blue line is the population of significant transcripts with $\gamma = 0.5$, and the cyan line is the significant population with $\gamma = 1$. Δ : Difference, *: significant, T: thresholded.

reduce the number of positive identifications to a manageable proportion of the whole dataset. Here, we applied the 14). Here, applying a dual-cutoff method using a fold-change using a heuristic of at least a $2^{1.4}$ fold change that reduces the number of significant outputs to 193 for DESeq2 and to 186 for ALDEx2. This cutoff was chosen for convenience and is mid-way between the high and low fold-change in-line with the recommendations of (9). These limits are 14) with the fold change limits shown by the dashed grey lines in Figure 2 and we can see that a 1. The $2^{1.4}$ fold change (~ 2.6 fold) cutoff identifies a similar number of transcripts as does ALDEx2 using $\gamma = 0.5$ which identifies 205 transcripts.

205. Supplementary Figures 1 shows an example of and 2 shows how to use the `aldex.senAnalysis()` function to identify those transcripts that are very sensitive to scale uncertainty in this dataset. Here, In this supplementary figure we see that even adding a very small amount of scale $\gamma = 0.1$ reduces the number of significant transcripts by more than half in the yeast dataset. This allows the analyst us to ignore those low-dispersion transcripts that were significant only because of an absence of scale uncertainty.

We next examined how adding scale would alter the analysis in a real dataset to which synthetically generated true positive counts had been added. We show results from the anti-PD-1 therapy RNA-seq dataset (53) which examined changes in gene expression when cells were exposed or not to a cell-cycle checkpoint inhibitor. This dataset was used by Li et al. (8) as an example of the dangers of relying on tools with high false positive error rates when analyzing clinical or clinically-related transcriptome samples. Indeed, in the benchmarking analysis done by this group, they found that parameter based methods such as DESeq2 and edgeR that on reported p-value. In practice, we suggest that a gamma parameter between 0.5 and fold-change cutoffs often led to conclusions in the original dataset that were indistinguishable from permutations of the dataset. In other words, that the analysis of transcriptome datasets from patient-derived samples often exhibited many false-positive identifications. 1 is realistic for most experimental designs (20).

The effect on dispersion with increasing amounts of scale uncertainty are shown in Figure 2A, where we can see that the dispersion increases as uncertainty is added. Note that the dispersion in the unscaled analysis in Figure 2A reaches a minimum near the mid-point of the distribution, and also does so when the analysis is conducted with DESeq2 (Supplementary Figure 3). This shows more clearly that dispersion of many transcripts is almost negligible in the absence of scale uncertainty. This plot makes the counter-intuitive suggestion that the variance in expression of the majority of genes with moderate expression is more predictable than highly-expressed genes or of housekeeping genes (48). This is at odds with the known biology of cells where single cell counting of highly-expressed transcripts shows that they have little intrinsic variation (23, 49).

Results of modeling true positive (TP) differences in the PD-1 dataset. For this the data were shuffled and

5% of the transcripts were modeled to have TP differences between groups where the differences were drawn from a Normal distribution with a mean difference of 0 and a sd of 2. Panel A shows the mean FDR of 10 instances for scale-naive ALDEx2 (A 0), and ALDEx2 with $\gamma = 0.2$ (A 0.2) or $\gamma = 0.5$ (A 0.5), and where significant features were identified by DESeq2 without (D 0), or with (D 0.5) a 0.5-fold change difference. The x-axis shows how the FDR changes for each tool as a function of the estimated modeled difference between groups. Panel B shows the mean sensitivity (TP found /all TP). Panels C and D show volcano and effect plots for the ALDEx2 output with the transcripts identified as significant at each scale setting as colored points, and non-significant transcripts in black.

Figure 3 shows the results of ten permutations of this dataset while simulating 5% of the transcripts to be true positives where the difference from no change was derived from a Normal distribution(54). This simulation was conducted with Adding scale uncertainty by setting $\gamma = 0.5$ or $\gamma = 1.0$ increases the minimum dispersion as shown in Figure 2A by the red and blue data points, and by the colored lines of fit through the centre of mass of the `seqgendiff` R package (54) to both permute the dataset and to add true positive features. We did ten permutations and kept track of data. Less obvious is that the number of true and false positive identifications. Figure 3A shows the actual false discovery rate for ALDEx2 with and without the addition of scale uncertainty and for DESeq2 with or without a fold-change cutoff of $2^{0.5}$. When the modeled difference was greater than 0, ALDEx2 exhibited a FDR of $\gamma = 0$: 0.002, $\gamma = 0.2$: 0, and $\gamma = 0.5$: 0, while DESeq2 with no fold-change cutoff had an FDR of 0.32 and with a 0.5 fold-change cutoff an FDR of 0.31. Figure 3 also plots these results as a function of the minimum modelled fold change of the true positive transcripts. First, we can see that this analysis recapitulates the observations of Li et al (8) in that DESeq2 has very poor false positive control at a nominal FDR of 0.05. The FDR is not controlled any better when a fold-change cutoff is applied, and this agrees with previous work showing that fold-change cutoffs do not materially improve FDR control in high throughput datasets (12, 13). additional dispersion is not applied equally to all points. Figure 3B shows that DESeq2 has higher sensitivity than does ALDEx2, and not surprisingly this sensitivity increases as 2B shows a plot of the difference between the difference between groups increases. For the case where the mean difference is 0 or greater, ALDEx2 exhibited a sensitivity of $\gamma = 0$: 0.66, $\gamma = 0.2$: 0.54, and $\gamma = 0.5$: 0.35, while DESeq2 with no fold-change cutoff had a sensitivity of 0.76 and with a 0.5 fold-change cutoff a sensitivity of 0.73. In this example, scale-naive ALDEx2 has near perfect FDR control and reasonable sensitivity at low modeled difference between groups. However, when the modeled difference between groups becomes large, then the scale-naive version of ALDEx2 begins to exhibit unacceptable rates of false positives and the false positive rate for DESeq2 also increases. Adding in even small amounts of scale uncertainty drops the true FDR rate to 0, but at the expense of sensitivity. The major contributor to the increase in FDR with larger modeled differences is that the tools are identifying as positives those transcripts that are modeled to have differences just below the threshold. We need to recognize that there is no such thing as a statistical free lunch; the analyst can have high sensitivity but low confidence that any individual transcript is truly different, or have lower sensitivity but have very high confidence that the difference is real. In other words, the sensitivity of a method is directly tied to how much error the investigator is willing to tolerate.

Examination of the volcano plot (11) and effect plot (50) in Figure 3C and D provides some insight into why adding scale uncertainty provides better FDR control than does the approach of using a p-value and a fold-change cutoff. In the volcano plot, adding scale uncertainty differentially excludes transcripts with a combination of marginal p-values and low difference between groups, and $\gamma = 0$ and $\gamma = 1$ data and here we can see that scale uncertainty is preferentially increasing the dispersion of the mid-expressed transcripts that this becomes more pronounced with a larger scale value. This effect is also seen in Figure 2 but is more nuanced. In contrast, the fold-change cutoff does not incorporate the magnitude of the p-value and so transcripts with large differences, but marginal p-values are retained. The effect plot in Figure 3D shows that the transcripts with marginal p-values and large differences that are excluded when scale uncertainty is added are those that have a large dispersion.

As a concrete example consider the point that is circled in panels C and D with a marginal p-value, with a difference between of nearly 4 and a dispersion of greater than 6. This transcript is no longer significant when $\gamma = 0.2$, but would require a very large fold-change cutoff to be excluded by the standard approach. In addition, transcripts with very small dispersion and very small differences are also excluded when scale

uncertainty is added. Thus, the addition of scale uncertainty achieves the desired outcome of lowering the FDR for those formerly had negligible dispersion; examine the grey line of best fit (overlaid by the red line) for the trend. Panel B also shows the trend of the expression-dispersion relationship for transcripts that are either marginally differentially abundant, or where the underlying dispersion and hence the uncertainty in measurement is very high, or for transcripts that fit both criteria.

Supplementary Figure 2 shows a second permuted dataset with the addition of modeled differences between groups. Here we used another real dataset with over 200 biological replicates of BRCA1 tumor and control tissue samples from Li et al. (8). This Supplementary Figure shows that the FDR control of DESeq2 is somewhat better than in the PD-1 dataset, although still much greater than the anticipated 5%. Further, a fold-change cutoff reduces the FDR of DESeq2 from about 30% to just over 20% with a power of over 80%. However, we can see that scale-naive ALDEx2 performs substantially better with a negligible FDR and comparable power. Adding scale uncertainty again improves FDR even for those transcripts modeled to have larger differences and classified as statistically significant. The red line shows the power is substantially better than in the PD-1 dataset, reaching the same power as DESeq2 or scale-naive ALDEx2 when the modeled difference is large. As before, this is driven by removing from consideration those transcripts with either a small difference between or a marginal p-value, or both. trendline with no added scale uncertainty, and this trendline exactly overlays with the grey trendline for the bulk of transcripts. The orange trendline indicates those transcripts that are both statistically significant and that have a thresholded expression level of ± 1.4 , and the dark blue and cyan lines show the statistically significant trendline for $\gamma = 0.5$, or 1.0.

Supplementary Figures 3 shows that effect of applying $\gamma = 0.5$ to this dataset results in reducing the number of positive transcripts from being $\sim 70\%$ of the whole dataset to less than 10% of the dataset and that this is largely because of a reduction in significance of those transcripts with low dispersion. Supplementary Figures 4 shows a sensitivity analysis of the BRCA1 dataset showing that different scale uncertainty amounts alter the number of significant transcripts in a biological replicate experiment similarly to a technical replicate experiment. Supplementary Figures 5 and 6 delve into how Thus, taking Figures 1 and 2 together adding scale uncertainty affects the variance-abundance relationship in subtle ways, and may help readers to understand the observations seen in Figure 3 and supplementary Figure 2.

Together the results in this section show that adding scale uncertainty has the desirable effect of altering the changing the distribution of transcripts identified as significantly different between groups in a way that exhibits better control of FDR albeit with a corresponding reduction in sensitivity. Those parts that were statistically significantly different *only because of low dispersion or that had marginal p-values or both*, are now are preferentially excluded from statistical significance. In practice, we suggest that a gamma parameter of between 0.2 and 0.5 is realistic for most experimental designs (22) regardless if the replication is technical or biological while those parts that were significantly different because of a high difference between groups remain.

Housekeeping genes and functions to guide scale model choices.

Dos Santos et al. (55,50) used a vaginal metatranscriptome dataset to compare the gene expression in bacteria collected from healthy (H) and bacterial vaginosis (BV) affected women. This dataset is derived from two publicly available datasets composed of a set of 20 non-pregnant women from London, Ontario Canada (56) and a subset of 22 non-pregnant women collected from German women who underwent metronidazole treatment for BV (57). Batch effects for these two groups were removed with ComBat-seq (58) and the two datasets were merged into one, giving a total of 16 H and 26 BV samples. In the Dos Santos paper, all results from this initial analysis were replicated in a much larger dataset derived from the MOMS-PI study (59)

In this vaginal In this environment, both the relative abundance of species between groups and the gene expression level within a species is different (60,51). Additionally, prior research suggests that the total number of bacteria is about 10 times more in the BV than in the H condition (28,26). Thus, these are extremely challenging datasets in which to determine differential abundance as there are both compositional and scale changes between conditions. The usual method to analyze vaginal metatranscriptome data is to do so on an organism-by-organism basis (57, 59, 60,51-53) because the scale confounding of the environment is less

pronounced. One attempt at system-wide analysis returned several housekeeping functions as differentially expressed between groups (5752); a result likely due to a disconnect between the assumptions of the normalization used and the actual scale of the environment (4519).

~~Analysis of vaginal transcriptome data aggregated at the Kegg Orthology (KO) functional level. Panel A shows an effect plot for the default analysis where the functions that are elevated in the healthy individuals have positive values and functions that are elevated in BV have negative values. Highlighted in the box are KOs that are almost exclusively housekeeping functions; these are colored cyan. These housekeeping functions should be located on the midline of no difference. Panel B shows the same data scaled with $\gamma = 0.5$, which increase the minimum dispersion as before. Panel C shows the same data scaled with $\gamma = 0.5$ and a 0.14 fold difference in dispersion applied to the BV samples relative to the H samples. In these plots statistically significant (q-value < 0.01) functions in the informed model are in red, false positive functions are in blue, non-significant functions in black and false negative functions are in orange.~~

In this example, we show how to specify and interpret ~~a user-defined or informed~~ an informed scale model that can explicitly account for some of these modeling difficulties (2220) even in a difficult ~~to analyze~~ dataset. An informed scale model can control for both the mean difference of scale between groups (e.g., directly incorporate information on the differences in total number of bacteria between the BV and H conditions) as well as the uncertainty of that difference ~~as illustrated in Figure 1:3.~~ To specify a user-defined scale model, we can pass a matrix of scale values instead of an estimate of just the scale uncertainty to `aldex.clr()`. This matrix should have the same number of rows as the of Monte-Carlo Dirichlet samples, and the same number of columns as the number of samples. While this matrix can be computed from scratch by the analyst, there is an `aldex.makeScaleModel()` function that can be used to simplify this step in most cases. This encodes the scale model as $\Lambda \sim N(\log_2 \mu_n, \gamma^2)$, where μ_n represents the scale value for each sample or group and ~~gamma-gamma~~ is the uncertainty as before. The scale estimate can be a measured value (cell count, nucleic acid input, ~~spike-ins~~, etc) or an estimate. Nixon et al. (3, 2220) showed that only the ratio of the means are important when providing values for μ_n ; i.e., the ratio between the $\log_2 \mu_i$ and $\log_2 \mu_j$ values. See the supplement to Nixon et al. (2220) for more information.

Figure ~~4A-3A~~ shows an effect plot of the data where reads are grouped by homologous function regardless of the organism of origin. Each point represents one of 3728 KEGG functions (6154). There are many more functions represented in the BV group (bottom) than in the healthy group (top). This is because the *Lactobacilli* that dominate a healthy vaginal microbiome have reduced genome content relative to the anaerobic organisms that dominate in BV, because there is a greater diversity of organisms in BV than in H samples, and because the BV condition has about an order of magnitude more bacteria than does the H condition.

The naive scale model appears to be reflecting the bacterial load as observed by calculating the mean scale value for each group. Using a negligible scale value; i.e., $\gamma = 1e - 3$ exposes the naive scale estimate for samples in the `@scaleSamps` slot from the `aldex.clr` output. ~~The~~ the naive scale estimate for the

healthy group is 17.41 and for the BV group is 14.59 for a difference of 2.82. This is interpreted as the scale of the H group of samples being 7.06 greater than the BV group. This precise but incorrect estimate places the location of the housekeeping genes off the midline of no difference.

Applying the default scale model of $\gamma = 0.5$ increases the dispersion slightly but does not move the housekeeping functions toward the midline. This is as expected; the mean of the default scale model is based on the CLR normalization so no shift in location would be expected over the ~~scale-naive-original~~ ALDEx2 model. Nevertheless, about 30% of the housekeeping functions are no longer statistically significantly different. Note that this change is simple to conduct, has no additional computational complexity and requires only a slight modification for the analyst.

There are 101 functions with low dispersion that appear to be shared by both groups (boxed area in Figure 3A, and colored in cyan). Inspection shows that these largely correspond to core metabolic functions such as transcription, translation, ribosomal functions, glycolysis, replication, chaperones, etc (Supplementary file housekeeping.txt). The transcripts of many of these are commonly used as invariant reference sequences in wet lab experiments (6248) and so are not be expected to contribute to differences in ecosystem behaviour.

~~Because we expect housekeeping functions to be nearly invariant in their expression and to occur in all organisms, the~~ The average location of these should be centred on 0 difference to represent an internal reference set. However, ~~with the naive without an informed~~ scale model, the mean of these housekeeping functions is approximately located at ~~+2.3. Thus, we~~




Figure 3: ~~Analysis of vaginal transcriptome data aggregated at the Kegg Orthology (KO) functional level. Panel A shows an effect plot for the default analysis where the functions that are elevated in the healthy individuals have positive values and functions that are elevated in BV have negative values. Highlighted in the box are KOs that are almost exclusively housekeeping functions; these are colored cyan. These housekeeping functions should be located on the midline of no difference. Panel B shows the same data scaled with $\gamma = 0.5$, which increase the minimum dispersion as before. Panel C shows the same data scaled with $\gamma = 0.5$ and a 0.14 fold difference in dispersion applied to the BV samples relative to the H samples. In these plots statistically significant (q-value < 0.01) functions in the informed model are in red, false positive functions are in blue, non-significant functions in black and false negative functions are in orange.~~

~~We~~ desire a scale model that approximately centres the housekeeping functions~~and~~, ~~because we expect housekeeping functions to be nearly invariant; thus~~ an appropriate scale in this dataset for functional analysis ~~will place these functions is likely~~ closer to 0 than ~~does~~ the naive estimate. One way to choose an appropriate value for μ_n is to use the `aldex.clr` function on only the presumed invariant functions setting $\gamma > 0$, and then accessing the `@scaleSamps` slot as before. Doing so suggests that the difference in scale should be about 14%. A second approach would be to identify the functions used as the denominator with the `denom="lvha"` option (~~1519~~) for the `aldex.clr` function, and then to use these values as before. This approach suggests a 5% difference in scale, and is potentially less subject to user interpretation.

For the purposes of this example, if we assume a 14% difference in scale, we can set $\mu_i = 1$ and $\mu_j = 1.14$ using the `makeScaleMatrix` function. This function uses a logNormal distribution to build a scale matrix given a user-specified mean difference between groups and uncertainty level. Applying a per-group relative differential scale of 0.14 moves the housekeeping functions close to the midline of no difference (Figure 3C, assuming 14% mean difference = -0.24, assuming a 5% mean difference = -0.34), and applying a gamma of 0.5 provides the same dispersion as in panel B of Figure 3. Note that now a significant number of functions are differentially up in BV that were formerly classed as not different without the full scale model (orange), or when only a default scale was applied. Inspection of the functions shows that these are largely missing from the *Lactobacillus* species and so should actually be captured as differentially abundant in the BV group. Supplementary Figure ~~7-4~~ shows that the using either the 5% or the 14% scale difference give imperceptibly different results suggesting that an informed scale model does not have to precisely estimate the scale difference to be useful. Nixon et al, (~~2220~~) also found that multiple reasonable estimates for the μ_n part of the informed scale model were similarly useful in microbiome data.

Thus, applying an informed scale allows us to distinguish between both false positives (housekeeping functions in cyan, and others in blue) and false negatives (orange functions) even in a very difficult to analyze dataset. ~~We used this scale model to uncover hiter-to-now unknown differences in microbiome functional activity between the Healthy and BV cohorts that were missed in previos analyses and that explain important clinical differences between them (55).~~ The remarkable improvements in biological interpretation afforded by an informed scale model, and the transferrability of it between sample cohorts of the same condition is outlined ~~in dos Santos et al. (55elsewhere (50)).~~ We suggest that the default scale model is sufficient when the data are approximately centred~~but that~~. ~~However~~, an informed model is more appropriate with datasets are not well centred or when the investigator has prior information about the underlying biology.

Discussion

~~Scale estimates affect two parts of the analysis. Modeling uncertainty in scale prevents false certainty in the precision of estimation and controls false positive identification. Modeling between-group scale relaxes the assumption of identity between the sizes of the environments and allows better control of false negative identification. The scale estimates can be derived from the total number of molecules in the environment or from other estimates of input size (cell counts, initial concentrations, spike-ins, growth rates, etc).~~

Biological systems are both predictably variable and stochastic (6349) and systems biology experiments show that there are transcripts with approximately constant concentrations in the cell and those with large variability under different growth conditions (2523). Current measurement methods that rely on high throughput sequencing fail to capture all of the variation, particularly variation due to scale (3, 2220). In the absence of external information (19, 20, 645, 6, 55) sequencing depth normalisation methods cannot recapture the scale information (19, 235, 21), and can only normalize for the technical variation due to sequencing depth. Here we demonstrated that even approximate estimates of the true system scale and the uncertainty of measuring it can aid in the interpretation of RNA-sequencing experiments.

Nixon et al. (3) introduced the idea of explicitly modeling the scale of a HTS dataset, and showed how to incorporate these models in the analysis of microbiome and other datasets (2220). They demonstrated that many tools commonly used to analyze HTS datasets had substantial Type 1 and Type 2 error rates, in line with recent findings by others (5, 7, 810, 12, 13). A version of ALDEx2 with the ability to include scale uncertainty was shown to be able to correct for the high Type 1 error rate for that tool, albeit with some loss of sensitivity. Finally, they showed that incorporating an informed scale model incorporating both location and scale uncertainty estimates could both control for Type 1 and Type 2 error rates (22, 4320).

~~The process of choosing the parameters are experiment-specific and can be anchored in known information such as cell counts, spike-ins, information from the literature or similar (3, 22, 65). In the metatranscriptome example used in this report (43); the choice of parameter was driven by the assumption inherent in the biology that core housekeeping functions would serve as an appropriate standard (37). The choice of parameters must be guided by the experimental question and other approaches in the literature which suggest normalizing the transcript levels to the bacterial metagenomic levels (66) could be used to set the scale parameters, but in this case are more granular at the individual organism level rather than at the systemic functional level.~~

Building and using a scale model thus has substantial benefits relative to the dual cutoff approach that is advocated for many gene expression experiments (9, 1114, 16). In particular, the dual cutoff approach has long been known to not control for Type 1 errors (12, 1317, 18), and the frequent lack of concordance between tools when benchmarked on transcriptomes (5, 7-9, 17, 6710, 12-14, 29, 56) and microbiomes (4, 6, 32, 42, 68, 699, 11, 31, 40, 57, 58) suggests poor control of Type 2 errors as well (5, 810, 13). Thus, incorporating a scale model during the analysis of HTS data promises the best of both worlds. A default scale model can control for Type 1 errors with minimal prior knowledge of the environment and this can be done with essentially no additional computational overhead. Furthermore, this work and previous (22, 4320) show that even minimal information about the underlying environment can be used to build a relatively robust informed scale model that controls for both Type 1 and 2 error rates. ~~It is important to note that the approach advocated here is distinct from that suggested by Zhang et al. (66, 70) where the DNA amount for a gene is a covariate in the model for transcriptomic differential abundance. In our analysis we grouped all the transcript information to functional level regardless of organism, instead of modeling per-organism gene abundances. In the future we anticipate being able to build more complex models similar to those used by Zhang et al. with the additional information of uncertainty in the underlying gene count.~~

In the analysis of HTS data it is often observed that larger datasets converge on the majority of parts being significantly different (3, 8, 913, 14). Li et al. (813) conducted a permutation-based benchmarking study and found that widely used tools performed worse than simple Wilcoxon rank-sum tests ~~coupled with the TPM normalization~~ in controlling the FDR when sample sizes became large. Li et al. ~~They~~ suggested that the presence of outliers were one of the factors driving the extreme FDR in some tests. ~~We found that when the Wilcoxon test was used within the ALDEx2 framework that it had essentially the same outputs as did the t-test. For example, in the PD-1 dataset where $\gamma = 0$ the ALDEx2 t-test exhibited a mean FDR of 0.2% and mean sensitivity of 65.9% while the corresponding values from the ALDEx2 Wilcoxon test were 0.3%~~

and 68.9%. This result again supports that the assumptions of the normalizations are as important or more important than the statistical test. [this observation](#). Brooks et al. (7159) suggested that inappropriate choice of benchmarking methods are also a major contributing factor and that better objective standards of truth are needed. In this report we generated semi-synthetic test data used binomial thinning which produces data that more closely mimic the properties of real high throughput sequencing data, and so can more rigorously test different tools (54). [important](#). From the perspective of our work the disagreement between tools can be explained by the observation that different analytic approaches produce different parameter estimates for either location or scale, or for both, as suggested in Figure 1. Thus, more data produces worse estimates because the additional data simply increases the precision of a flawed estimate (3, 7260).

Scale simulation is now built into ALDEx2 (22) and in this report (20) and [here](#) we suggest that there are two main root causes to common HTS data pathologies. The first contributing factor is the observed very low dispersion estimate for many features that is a by-product of some experimental designs and [normalizations](#) (Figure of normalization). In the Schurch et al. (14 dataset), the data were from single colonies derived from a single culture. Thus, it is more accurate to describe the 96 samples as wet-lab technical replicates rather than independent samples. This type of replication approach is standard in the molecular literature, and would be expected to result in the very low dispersion that is observed. Applying the default scale model with $\gamma = 0.5$ a large number of transcripts have their dispersion increased (Figure 1D), with the effect being largest for those with the lowest initial dispersion (Figure 2). [Supplying additional uncertainty alleviates many FP but in a way that more appropriately controls the FDR as](#) Adding scale uncertainty results in modest number of transcripts, 205, being called significantly different as shown in the volcano plot in Figure 1B (red points). In addition, there is now a strong concordance between the difference and q-values. In hindsight, it is not obvious why the unscaled volcano plot shows such poor correspondence. We suggest that this is explained by random fluctuations of the many very low variance estimates and this is supported by the plots shown in Figure 3-2.

The second contributing factor is unacknowledged asymmetry in many datasets (4519); i.e., different gene content or a directional change in the majority of features. In the case of asymmetry, the use of a user-specified scale model can be very useful for otherwise difficult-to-analyze datasets such as meta-transcriptomes and in-vitro selection datasets where the majority of features can change as shown in Figure 4. We showed one such example for the metatranscriptome dataset in Figure 3. Here the dataset was highly asymmetrical. Incorporating differential scale on a per-group basis moves the mass of the housekeeping functions towards the midline of no difference and so affects both Type I and Type II error rates. We showed two ways of estimating the scale difference between groups and found that any reasonable estimate is an improvement over the naive approach and also over the default scale model. This is in line with the observations by Nixon et al (2220) in a 16S rRNA gene sequencing dataset. It is also of note that in the case of true biological replicates (different individuals) that adding a modest amount of scale $\gamma = 0.5$ had little effect on the the difference between groups and on the dispersion. Thus, in this dataset the scale mis-specification was affecting mainly the location of the difference between groups. While we acknowledge that some prior information is needed on which housekeeping transcripts should not be classed as differentially abundant is needed, we suggest that this information is widely available and is already used when performing the gold-standard quantitative PCR test of differential abundance (73, 7461, 62).

Beyond concerns of fidelity and rigor, scale models also enhance the reproducibility and transparency of HTS analyses. The development of HTS and the associated problems of very high dimensional data that was not always statistically well-behaved led to many different proposed solutions including multiple normalizations and moderated test statistics. That these perform poorly in real data is shown by the simulation data in this report and elsewhere where both DESeq2 (which we used) and edgeR (which uses moderated statistical tests) performed poorly in controlling the FDR (8). While these approaches often work in many datasets they fail to address the underlying problems of information that is missing in the data which is what is supplied by adding uncertainty around the information we have about that data. The addition of scale uncertainty directly addresses the missing information by testing the [essentially tests the](#) model over a range of normalizations (3). In doing so, the scale-based approach removes the need for moderated statistics and [and so](#) can replace the consensus approach that has been proposed by some groups (6, 7511, 63) with no additional computational overhead. Thus, an advantage of incorporating scale is that analyses can be made

much more robust such that actual or potential differences in scale can be tested and accounted for explicitly. While it is beyond the scope of the present article, we note that there are many ways of building scale models that enhance the interpretability of the parameters and assumptions and a detailed description of these points is describe elsewhere (3).

In summary, we supply a toolkit that makes incorporating scale uncertainty and location information simple to incorporate for transcriptomes or indeed any type of HTS dataset. While the underlying scale of the system is generally inaccessible, the effect of scale ~~uncertainty~~ on the analysis outcomes can be modelled and can help explain some of the underlying biology, [and help to expose known issues with the analysis of HTS data](#). Adding scale information to the analysis allows for more robust inference because the features that are sensitive to scale can be identified and their impact on conclusions weighted accordingly. ~~The~~ [Additionally, the](#) use of informed scale models permits difficult to analyze datasets to be examined in a robust and principled manner even when the majority of features are asymmetrically distributed or expressed (or both) in the groups (5550). Thus, using and reporting scale uncertainty should become a standard practice in the analysis of HTS datasets.

References

1. Lovell,D., Müller,W., Taylor,J., Zwart,A. and Helliwell,C. (2011) Proportions, percentages, ppm: Do the molecular biosciences treat compositional data right? In Pawlowsky-Glahn,V., Buccianti,A. (eds), *Compositional Data Analysis: Theory and Applications*. John Wiley; Sons New York, NY, London, pp. 193–207.
2. Quinn,T.P., Erb,I., Gloor,G., Notredame,C., Richardson,M.F. and Crowley,T.M. (2019) A field guide for the compositional analysis of any-omics data. *Gigascience*, **8**.
3. Nixon,M.P., McGovern,K.C., Letourneau,J., David,L.A., Lazar,N.A., Mukherjee,S. and Silverman,J.D. (2024) Scale reliant inference.
4. [Jiang,L., Schlesinger,F., Davis,C.A., Zhang,Y., Li,R., Salit,M., Gingeras,T.R. and Oliver.B. \(2011\) Synthetic spike-in standards for RNA-seq experiments. *Genome Res*, **21**, 1543–51.](#)
5. [Lovén,J., Orlando,D.A., Sigova,A.A., Lin,C.Y., Rahl,P.B., Burge,C.B., Levens,D.L., Lee,T.I. and Young,R.A. \(2012\) Revisiting global gene expression analysis. *Cell*, **151**, 476–82.](#)
6. [Vandeputte,D., Kathagen,G., D’hoë,K., Vieira-Silva,S., Valles-Colomer,M., Sabino,J., Wang,J., Tito,R.Y., De Commer,L., Darzi,Y., et al. \(2017\) Quantitative microbiome profiling links gut community variation to microbial load. *Nature*, **551**, 507–511.](#)
7. [Props,R., Kerckhof,F.-M., Rubbens,P., De Vrieze,J., Hernandez Sanabria,E., Waegeman,W., Monsieurs,P., Hammes,F. and Boon,N. \(2017\) Absolute quantification of microbial taxon abundances. *ISME J*, **11**, 584–587.](#)
8. [Nishijima,S., Stankevic,E., Aasmets,O., Schmidt,T.S.B., Nagata,N., Keller,M.I., Ferretti,P., Juel,H.B., Fullam,A., Robbani,S.M., et al. \(2024\) Fecal microbial load is a major determinant of gut microbiome variation and a confounder for disease associations. *Cell*, 10.1016/j.cell.2024.10.022.](#)
9. Thorsen,J., Brejnrod,A., Mortensen,M., Rasmussen,M.A., Stokholm,J., Al-Soud,W.A., Sørensen,S., Bisgaard,H. and Waage,J. (2016) Large-scale benchmarking reveals false discoveries and count transformation sensitivity in 16S rRNA gene amplicon data analysis methods used in microbiome studies. *Microbiome*, **4**, 62.

- ~~5–10.~~ Quinn,T.P., Crowley,T.M. and Richardson,M.F. (2018) Benchmarking differential expression analysis tools for RNA-seq: Normalization-based vs. Log-ratio transformation-based methods. *BMC Bioinformatics*, **19**, 274.
- ~~6–11.~~ Nearing,J.T., Douglas,G.M., Hayes,M.G., MacDonald,J., Desai,D.K., Allward,N., Jones,C.M.A., Wright,R.J., Dhanani,A.S., Comeau,A.M., *et al.* (2022) Microbiome differential abundance methods produce different results across 38 datasets. *Nat Commun*, **13**, 342.
- ~~7–12.~~ Ge,X., Chen,Y.E., Song,D., McDermott,M., Woyshner,K., Manousopoulou,A., Wang,N., Li,W., Wang,L.D. and Li,J.J. (2021) Clipper: P-value-free FDR control on high-throughput data from two conditions. *Genome Biol*, **22**, 288.
- ~~8–13.~~ Li,Y., Ge,X., Peng,F., Li,W. and Li,J.J. (2022) Exaggerated false positives by popular differential expression methods when analyzing human population samples. *Genome Biol*, **23**, 79.
- ~~9–14.~~ Schurch,N.J., Schofield,P., Gierliński,M., Cole,C., Sherstnev,A., Singh,V., Wrobel,N., Gharbi,K., Simpson,G.G., Owen-Hughes,T., *et al.* (2016) How many biological replicates are needed in an RNA-seq experiment and which differential expression tool should you use? *RNA*, **22**, 839–51.
- ~~10–15.~~ Storey,J.D. (2003) The positive false discovery rate: A bayesian interpretation and the q-value. *The annals of statistics*, **31**, 2013–2035.
- ~~11–16.~~ Cui,X. and Churchill,G.A. (2003) Statistical tests for differential expression in cDNA microarray experiments. *Genome Biol*, **4**, 210.1–210.10.
- ~~12–17.~~ Zhang,S. and Cao,J. (2009) A close examination of double filtering with fold change and t test in microarray analysis. *BMC Bioinformatics*, **10**, 402.
- ~~13–18.~~ Ebrahimpour,M. and Goeman,J.J. (2021) Inflated false discovery rate due to volcano plots: Problem and solutions. *Brief Bioinform*, **22**.
- ~~14–~~ Simmons,J.P., Nelson,L.D. and Simonsohn,U. (2011) False-positive psychology: Undisclosed flexibility in data collection and analysis allows presenting anything as significant. *Psychological science*, **22**, 1359–1366.
- ~~15–19.~~ Wu,J.R., Macklaim,J.M., Genge,B.L. and Gloor,G.B. (2021) Finding the centre: Compositional asymmetry in high-throughput sequencing datasets. In Filzmoser,P., Hron,K., Martín-Fernández,J.A., Palarea-Albaladejo,J. (eds), *Advances in compositional data analysis: Festschrift in honour of vera pawlowsky-glahn*. Springer International Publishing, Cham, pp. 329–346.
- ~~16–~~ Nishijima,S., Stankevic,E., Aasmets,O., Schmidt,T.S.B., Nagata,N., Keller,M.I., Ferretti,P., Juel,H.B., Fullam,A., Robbani,S.M., *et al.* (2024) Fecal microbial load is a major determinant of gut microbiome variation and a confounder for disease associations. *Cell*, 10.1016/j.cell.2024.10.022.
- ~~17–~~ Bullard,J.H., Purdom,E., Hansen,K.D. and Dudoit,S. (2010) Evaluation of statistical methods for normalization and differential expression in mRNA-seq experiments. *BMC Bioinformatics*, **11**, 94.
- ~~18–~~ Jiang,L., Schlesinger,F., Davis,C.A., Zhang,Y., Li,R., Salit,M., Gingeras,T.R. and Oliver,B. (2011) Synthetic spike-in standards for RNA-seq experiments. *Genome Res*, **21**, 1543–51.
- ~~19–~~ Lovén,J., Orlando,D.A., Sigova,A.A., Lin,C.Y., Rahl,P.B., Burge,C.B., Levens,D.L., Lee,T.I. and Young,R.A. (2012) Revisiting global gene expression analysis. *Cell*, **151**, 476–82.
- ~~20–~~ Vandeputte,D., Kathagen,G., D’hoë,K., Vieira-Silva,S., Valles-Colomer,M., Sabino,J., Wang,J., Tito,R.Y., De Commer,L., Darzi,Y., *et al.* (2017) Quantitative microbiome profiling links gut community variation to microbial load. *Nature*, **551**, 507–511.
- ~~21–~~ Props,R., Kerekhof,F.-M., Rubbens,P., De Vrieze,J., Hernandez-Sanabria,E., Waegeman,W., Monsieus,P., Hammes,F. and Boon,N. (2017) Absolute quantification of microbial taxon abundances.

- ~~22.~~20. Nixon,M.P., Gloor,G.B. and Silverman,J.D. (2024) Beyond normalization: Incorporating scale uncertainty in microbiome and gene expression analysis. *bioRxiv*, 10.1101/2024.04.01.587602.
- ~~23.~~21. Lovell,D., Pawlowsky-Glahn,V., Egozcue,J.J., Marguerat,S. and Bähler,J. (2015) Proportionality: A valid alternative to correlation for relative data. *PLoS Comput Biol*, **11**, e1004075.
- ~~24.~~22. Nie,Z., Hu,G., Wei,G., Cui,K., Yamane,A., Resch,W., Wang,R., Green,D.R., Tessarollo,L., Casellas,R., *et al.* (2012) C-Myc is a universal amplifier of expressed genes in lymphocytes and embryonic stem cells. *Cell*, **151**, 68–79.
- ~~25.~~23. Scott,M., Gunderson,C.W., Mateescu,E.M., Zhang,Z. and Hwa,T. (2010) Interdependence of cell growth and gene expression: Origins and consequences. *Science*, **330**, 1099–102.
- ~~26.~~24. Yoshikawa,K., Tanaka,T., Ida,Y., Furusawa,C., Hirasawa,T. and Shimizu,H. (2011) Comprehensive phenotypic analysis of single-gene deletion and overexpression strains of *saccharomyces cerevisiae*. *Yeast*, **28**, 349–61.
- ~~27.~~25. Lin,J. and Amir,A. (2018) Homeostasis of protein and mRNA concentrations in growing cells. *Nat Commun*, **9**, 4496.
- ~~28.~~26. Zozaya-Hinchliffe,M., Lillis,R., Martin,D.H. and Ferris,M.J. (2010) Quantitative PCR assessments of bacterial species in women with and without bacterial vaginosis. *J Clin Microbiol*, **48**, 1812–9.
- ~~29.~~27. Ravel,J., Gajer,P., Abdo,Z., Schneider,G.M., Koenig,S.S.K., McCulle,S.L., Karlebach,S., Gorle,R., Russell,J., Tacket,C.O., *et al.* (2011) Vaginal microbiome of reproductive-age women. *Proc Natl Acad Sci U S A*, doi/10.1073/pnas.100611107.
- ~~30.~~28. Hummelen,R., Fernandes,A.D., Macklaim,J.M., Dickson,R.J., Changalucha,J., Gloor,G.B. and Reid,G. (2010) Deep sequencing of the vaginal microbiota of women with HIV. *PLoS One*, **5**, e12078.
29. Bullard,J.H., Purdom,E., Hansen,K.D. and Dudoit,S. (2010) Evaluation of statistical methods for normalization and differential expression in mRNA-seq experiments. *BMC Bioinformatics*, **11**, 94.
- ~~31.~~30. Dillies,M.-A., Rau,A., Aubert,J., Hennequet-Antier,C., Jeanmougin,M., Servant,N., Keime,C., Marot,G., Castel,D., Estelle,J., *et al.* (2013) A comprehensive evaluation of normalization methods for Illumina high-throughput RNA sequencing data analysis. *Brief Bioinform*, **14**, 671–83.
- ~~32.~~31. Weiss,S., Xu,Z.Z., Peddada,S., Amir,A., Bittinger,K., Gonzalez,A., Lozupone,C., Zaneveld,J.R., Vázquez-Baeza,Y., Birmingham,A., *et al.* (2017) Normalization and microbial differential abundance strategies depend upon data characteristics. *Microbiome*, **5**, 27.
- ~~33.~~32. Hughes,J.B. and Hellmann,J.J. (2005) The application of rarefaction techniques to molecular inventories of microbial diversity. *Methods Enzymol*, **397**, 292–308.
- ~~34.~~33. Mortazavi,A., Williams,B.A., McCue,K., Schaeffer,L. and Wold,B. (2008) Mapping and quantifying mammalian transcriptomes by RNA-seq. *Nat Methods*, **5**, 621–8.
- ~~35.~~34. Wagner,G.P., Kin,K. and Lynch,V.J. (2012) Measurement of mRNA abundance using RNA-seq data: RPKM measure is inconsistent among samples. *Theory Biosci*, **131**, 281–5.
- ~~36.~~35. Robinson,M.D. and Oshlack,A. (2010) A scaling normalization method for differential expression analysis of RNA-seq data. *Genome Biol*, **11**, R25.1–R25.9.

- ~~37. Vandesompele, J., De Preter, K., Pattyn, F., Poppe, B., Van Roy, N., De Paepe, A. and Speleman, F. (2002) Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes. *Genome Biol*, **3**, RESEARCH0034.~~
- ~~38. 36.~~ Aitchison, J. (1982) The statistical analysis of compositional data. *Journal of the Royal Statistical Society: Series B (Methodological)*, **44**, 139–160.
- ~~39. 37.~~ Anders, S. and Huber, W. (2010) Differential expression analysis for sequence count data. *Genome Biol*, **11**, R106.
- ~~40. 38.~~ Aitchison, J. (1986) The statistical analysis of compositional data Chapman & Hall, London, England.
- ~~41. 39.~~ Fernandes, A.D., Macklaim, J.M., Linn, T.G., Reid, G. and Gloor, G.B. (2013) ~~ANOVA-like~~ ANOVA-like differential expression (ALDEx) analysis for mixed population ~~RNA-seq~~ RNA-seq. *PLoS One*, **8**, e67019.
- ~~42. 40.~~ Yerke, A., Fry Brumit, D. and Fodor, A.A. (2024) Proportion-based normalizations outperform compositional data transformations in machine learning applications. *Microbiome*, **12**, 45.
- ~~43. Dos Santos, S.J., Copeland, C., Macklaim, J.M., Reid, G. and Gloor, G.B. (2024) Vaginal metatranscriptome meta-analysis reveals functional BVsubgroups and novel colonisation strategies. *Microbiome*, **12**, 271.~~
- ~~44. 41.~~ Love, M.I., Huber, W. and Anders, S. (2014) Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol*, **15**, 550.1–550.21.
- ~~45. 42.~~ Paulson, J.N., Stine, O.C., Bravo, H.C. and Pop, M. (2013) Differential abundance analysis for microbial marker-gene surveys. *Nat Methods*, **10**, 1200–2.
- ~~46. 43.~~ Fernandes, A.D., Reid, J.N., Macklaim, J.M., McMurrough, T.A., Edgell, D.R. and Gloor, G.B. (2014) Unifying the analysis of high-throughput sequencing datasets: Characterizing RNA-seq, 16S rRNA gene sequencing and selective growth experiments by compositional data analysis. *Microbiome*, **2**, 15.1–15.13.
- ~~47. 44.~~ Gierliński, M., Cole, C., Schofield, P., Schurch, N.J., Sherstnev, A., Singh, V., Wrobel, N., Gharbi, K., Simpson, G., Owen-Hughes, T., *et al.* (2015) Statistical models for RNA-seq data derived from a two-condition 48-replicate experiment. *Bioinformatics*, **31**, 3625–3630.
- ~~48. 45.~~ Benjamini, Y. and Hochberg, Y. (1995) Controlling the false discovery rate: A practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society. Series B (Methodological)*, **57**, 289–300.
- ~~49. 46.~~ Efron, B. (2008) Microarrays, empirical bayes and the two-groups model. *Statist. Sci.*, **23**, 1–22.
- ~~50. 47.~~ Gloor, G., Macklaim, J. and Fernandes, A. (2016) Displaying variation in large datasets: Plotting a visual summary of effect sizes. *Journal of Computational and Graphical Statistics*, **25**, 971–979.
- ~~51. Witten~~
- ~~48. Rocha, D. and Tibshirani, R. (2007) A comparison of fold-change and the t-statistic for microarray data analysis. *Analysis* **J.P.G.**, **1776**, 58–85.~~
- ~~52. Liu, X., Zhao, J., Xue, L., Zhao, T., Ding, W., Han, Y. and Ye, H. (2022) A comparison of transcriptome analysis methods with reference genome in bacteria by RT-qPCR. *BMC Genomics* **Methods Mol Biol**, **232065**, 232. 119–137.~~
- ~~53. Riaz, N., Havel, J~~

49. [Taniguchi,Y., Choi,P.J., Makarov,V.,Li,G.-W., Chen,H., Babu,M., Desrichard,A.,—Urba,W.J., SimsHearn,J.S.,—Hodi,F. S.,Martín-Algarra,S., Mandal,R., Sharfman,W.H.,—et al. \(2017\) Tumor-, Emili,A. and Xie,X.S. \(2010\) Quantifying e. Coli proteome and microenvironment evolution during immunotherapy transcriptome with nivolumab](#) [single-molecule sensitivity in single cells. *CellScience*, **171**329, 934–949. e16–533–8.](#)
54. [Gerard,D. \(2020\) Data-based RNA-seq simulations by binomial thinning. *BMC Bioinformatics*, **21**, 206.](#)
55. [50. Dos Dos Santos,S.J., Copeland,C., Macklaim,J.M., Reid,G. and Gloor,G.B. \(2024\) Vaginal meta-transcriptome meta-analysis reveals functional BV subgroups and novel colonisation strategies. *bioRxiv*, 10.1101/2024.04.24.590967.](#)
56. [—](#)
51. [Macklaim,J.M., Fernandes,A.D., Di Bella,J.M., Hammond,J.-A., Reid,G. and Gloor,G.B. \(2018\) From 2013\) Comparative meta-RNA-seq to biological inference: Using compositional data analysis of the vaginal microbiota and differential expression by lactobacillus iners in meta-transcriptomics](#) [health and dysbiosis. *Methods Mol BiolMicrobiome*, **184**91, 193–213–12.](#)
57. [52. Deng,Z.-L., Gottschick,C., Bhuju,S., Masur,C., Abels,C. and Wagner-Döbler,I. \(2018\) Metatranscriptome analysis of the vaginal microbiota reveals potential mechanisms for protection against metronidazole in bacterial vaginosis. *mSphere*, **3**.](#)
58. [Zhang,Y., Parmigiani,G. and Johnson,W.E. \(2020\) ComBat-seq: Batch effect adjustment for RNA-seq count data. *NAR Genom Bioinform*, **2**, lqaa078.—](#)
59. [53. Fettweis,J.M., Serrano,M.G., Brooks,J.P., Edwards,D.J., Girerd,P.H., Parikh,H.I., Huang,B., Ar-odz,T.J., Edupuganti,L., Glascock,A.L., et al. \(2019\) The vaginal microbiome and preterm birth. *Nat Med*, **25**, 1012–1021.](#)
60. [—Macklaim,J.M., Fernandes,A.D., Di Bella,J.M., Hammond,J.-A., Reid,G. and Gloor,G.B. \(2013\) Comparative meta-RNA-seq of the vaginal microbiota and differential expression by *Lactobacillus iners* in health and dysbiosis. *Microbiome*, **1**, 12.—](#)
61. [54. Okuda,S., Yamada,T., Hamajima,M., Itoh,M., Katayama,T., Bork,P., Goto,S. and Kanehisa,M. \(2008\) KEGG atlas mapping for global analysis of metabolic pathways. *Nucleic Acids Res*, **36**, W423–6.](#)
62. [Rocha,D.J.P.G., Castro,T.L.P., Aguiar,E.R.G.R. and Pacheco,L.G.C. \(2020\) Gene expression analysis in bacteria by RT-qPCR. *Methods Mol Biol*, **2065**, 119–137.—](#)
63. [—Taniguchi,Y., Choi,P.J., Li,G.-W., Chen,H., Babu,M., Hearn,J., Emili,A. and Xie,X.S. \(2010\) Quantifying e. Coli proteome and transcriptome with single-molecule sensitivity in single cells. *Science*, **329**, 533–8.—](#)
64. [55. Marguerat,S., Schmidt,A., Codlin,S., Chen,W., Aebersold,R. and Bähler,J. \(2012\) Quantitative analysis of fission yeast transcriptomes and proteomes in proliferating and quiescent cells. *Cell*, **151**, 671–83.](#)
65. [—McGovern,K.C., Nixon,M.P. and Silverman,J.D. \(2023\) Addressing erroneous scale assumptions in microbe and gene set enrichment analysis. *PLoS Comput Biol*, **19**, e1011659.—](#)
66. [—Zhang,Y., Thompson,K.N., Huttenhower,C. and Franzosa,E.A. \(2021\) Statistical approaches for differential expression analysis in metatranscriptomics. *Bioinformatics*, **37**, i34–i41.—](#)
67. [56. Sonesson,C. and Delorenzi,M. \(2013\) A comparison of methods for differential expression analysis of](#)

- RNA-seq data. *BMC Bioinformatics*, **14**, 91.
- ~~68.~~57. McMurdie,P.J. and Holmes,S. (2014) Waste not, want not: Why rarefying microbiome data is inadmissible. *PLoS Comput Biol*, **10**, e1003531.
- ~~69.~~58. Hawinkel,S., Mattiello,F., Bijmens,L. and Thas,O. (2018) A broken promise : Microbiome differential abundance methods do not control the false discovery rate. *Briefings in Bioinformatics*.
- ~~70.~~Zhang,Y., Thompson,K.N., Branek,T., Yan,Y., Nguyen,L.H., Franzosa,E.A. and Huttenhower,C. (2021) Metatranscriptomics for the human microbiome and microbial community functional profiling. ~~*Annu Rev Biomed Data Sci*, **4**, 279–311.~~
- ~~71.~~59. Brooks,T.G., Lahens,N.F., Mrčela,A. and Grant,G.R. (2024) Challenges and best practices in omics benchmarking. *Nat Rev Genet*, **25**, 326–339.
- ~~72.~~60. Gustafson,P. (2015) Bayesian inference for partially identified models: Exploring the limits of limited data CRC Press.
- ~~73.~~61. Thellin,O., Zorzi,W., Lakaye,B., De Borman,B., Coumans,B., Hennen,G., Grisar,T., Igout,A. and Heinen,E. (1999) Housekeeping genes as internal standards: Use and limits. *J Biotechnol*, **75**, 291–5.
- ~~74.~~62. SEQC/MAQC-III Consortium (2014) A comprehensive assessment of RNA-seq accuracy, reproducibility and information content by the sequencing quality control consortium. *Nat Biotechnol*, **32**, 903–14.
- ~~75.~~63. Song,H., Ling,W., Zhao,N., Plantinga,A.M., Broedlow,C.A., Klatt,N.R., Hensley-McBain,T. and Wu,M.C. (2023) Accommodating multiple potential normalizations in microbiome associations studies. *BMC Bioinformatics*, **24**, 22.