

Exploratory analysis of RNA-seq dataset

gg

2016-01-18

To run this file: Rscript -e "rmarkdown::render('biplots.Rmd')"

Setup

```
library(compositions)
library(zCompositions)
library(ALDEx2)
# you can get this from:
# https://github.com/DavidRLovell/propr
source("~/git/proprBayes/R/propr-functions.R")

library(igraph)

#####
# these are functins that we need for phi.
sma.df <- function(df){
df.cor <- stats::cor(df, use="pairwise.complete.obs")
df.var <- stats::cov(df, use="pairwise.complete.obs")
df.sd <- sqrt(diag(df.var))
r.rf2 <-
  (outer(diag(df.var), diag(df.var), "-")^2 ) /
  (outer(diag(df.var), diag(df.var), "+")^2 - 4 * df.var^2 )
diag(r.rf2) <- 0
res.dof <- nrow(df) - 2
F <- r.rf2/(1 - r.rf2) * res.dof
list(b=sign(df.cor) * outer(df.sd, df.sd, "/"),
p=1 - pf(F, 1, res.dof),
r2=df.cor^2)
}

lt.row.min <- function(X){

  result <- numeric(nrow(X) - 1)

  for(i in 2:nrow(X)){

    result[i-1] <- min(X[i, 1:i-1])

  }

  result
}

propr.phisym <- function (X)
{
```

```

Cov      <- stats::var(X)
tmp      <- 2 * Cov / outer(diag(Cov), diag(Cov), "+")
return((1-tmp)/(1+tmp))
}

##### incorporating phi into ALDEx
# calculate phi on Dirichlet log-ratio distributions
# returns dataframe of the lower-triangle of symmetrical phi metric
# requires David Lovell's propr.phisym function
# this depends in turn on the composits R package
# turtles all the way down!

# we ignore all the other measures that are used for trouble-shooting phi
# the sma.df function in particular is very time and memory intensive

aldex.phi <- function(aldex.clr){

  # calculate expected value of phi
  # a single high phi value will push the component out of consideration
  # a median is right out for memory considerations

  # get first value
  sym.phi <- propr.phisym(t(sapply(getMonteCarloInstances(aldex.clr), function(y){y[,1]})))

  # sum the rest of the values as we proceed through the DIR MC instances
  for(i in 2:numMCInstances(aldex.clr)){
    #print(i)
    sym.phi <- sym.phi + propr.phisym(t(sapply(getMonteCarloInstances(aldex.clr), function(y){y[,i]})))
  }

  ##### Done ALDEx2 stuff

  # make indices of the correct size
  lt <- which(col(sym.phi)<row(sym.phi), arr.ind=FALSE)
  lt.ind <- which(col(sym.phi)<row(sym.phi), arr.ind=TRUE)

  # dataframe to hold the info,
  # data is a set of vectors where only the lower triangle is kept, because the matrix is symmetrical
  # this is needed so the subset function works properly
  sma.df <- data.frame(row=factor(rownames(sym.phi)[lt.ind[, "row"]]),
    col=factor(colnames(sym.phi)[lt.ind[, "col"]]))

  #save the lower triangle as an expected value
  sma.df$phi <- sym.phi[lt] / numMCInstances(aldex.clr)

  return(sma.df)
}

##### end functions

```

First let us explore the data. The idea here is to identify those samples that are closely related

Filter the refseq to a mean count across samples of greater than 5

This is a form biplot that represents distances between samples with variances of components mapped onto

the plot.

```
# this is the taxon table
o <- read.table(
  "~/git/twntyfr/data/merged_readcounts_taxonomy_qiime_L6_16Sorder_color_sum.txt", header=T, row.names=
)

# this is the refseq set
d <- read.table("data/merged_readcounts_subsys4.txt", header=T,
  row.names=1, check.names=F, sep="\t", comment.char="", quote="")

sub4 <- d$subsys4
d$subsys4 <- NULL

refseq <- d$refseqID
d$refseqID <- NULL

len <- d$length
d$length <- NULL

d.min <- d[which(apply(d,1,mean)> 5),]

# we need to replace 0 values with a best estimate
# use zCompositions CZM by default
# but samples must be by row, so use t()

d.n0.CZM <- cmultRepl(t(d.min), label=0, method="CZM")

## No. corrected values: 20044

# turn this into a centered log-ratio transform
# samples are by row
# remember that apply by row rotates the data
# R is terrible, so we need to use t() again

d.clr <- t( apply(d.n0.CZM, 1, function(x){log(x) - mean(log(x))}) )

d.mvar.clr <- mvar(d.clr)

d.pcx <- prcomp(d.clr)

# TODO color samples by most abundant organism
biplot(d.pcx, cex=c(0.5,0.05), col=c("black",rgb(1,0,0,0.2)), var.axes=F,
  xlab=paste("PC1: ", round(sum(d.pcx$sdev[1]^2)/d.mvar.clr, 3)),
  ylab=paste("PC2: ", round(sum(d.pcx$sdev[2]^2)/d.mvar.clr, 3)),
  scale=0
)

# filter the taxon table by what is in e.min
o.d <- o[,colnames(d)]
o.d.CZM <- cmultRepl(t(o.d), label=0, method="CZM")
o.d.clr <- t( apply(o.d.CZM, 1, function(x){log(x) - mean(log(x))}) )
```


in different genomes.

To get around the gene occurrence problem, we aggregate the data to the SEED subsystem 4 level.

```
e <- read.table("data/AitchisonTransformedDataForALDExInput.txt", header=T,
  row.names=1, check.names=F, sep="\t", comment.char="", quote="")

e.n0.CZM <- cmultRepl(t(e), label=0, method="CZM")

## No. corrected values: 2869


# turn this into a centered log-ratio transform
# samples are by row
# remember that apply by row rotates the data
# R is terrible, so we need to use t() again

e.clr <- t( apply(e.n0.CZM, 1, function(x){log(x) - mean(log(x))}) )

e.mvar.clr <- mvar(e.clr)

e.pcx <- prcomp(e.clr)

# TODO color samples by most abundant organism
biplot(e.pcx, cex=c(0.5,0.05), col=c("black",rgb(1,0,0,0.2)), var.axes=F,
  xlab=paste("PC1: ", round(sum(e.pcx$sdev[1]^2)/e.mvar.clr, 3)),
  ylab=paste("PC2: ", round(sum(e.pcx$sdev[2]^2)/e.mvar.clr, 3)),
  scale=0
)
```



This is a much better representation of the data. We did not bother filtering by abundance since there is a much smaller number of features (3874). Now we can see a fairly clear split in the data that is less driven by

gene occurrence and more by functions within the microbiome (cite Beiko review regarding ecosystems of genes rather than organisms here!).

```
mydata <- as.data.frame(e.clr)

# calculate the within group sum of squares
wss <- (nrow(mydata) - 1) * sum(apply(mydata, 2, var))

# summarize the fit for the first 15 groups
for (i in 2:15) wss[i] <- sum(kmeans(mydata,
  centers=i)$withinss)

# the plot suggests a reasonable fit can be had with about 2-3 clusters
# and after that there are only small improvements with additional clusters
layout(matrix(c(1,2,3,4), 2, 2, byrow=T), widths=c(10,10), height=c(10,10))


par(mar=c(5,4,0,5)+0.1)

# make the sum of squares distance plot
plot(1:15, wss[1:15], type="b", xlab="Number of Clusters",
  ylab="Win Grp SS")

par(mar=c(2,0,2,0)+0.1)

# plot the data for cluster sizes 2-4
for(i in 2:4){
  fit <- kmeans(mydata,i)
  mydata$fit.cluster <- NULL # clear any previous data
  mydata <- data.frame(mydata, fit$cluster) # add the cluster data

  coloredBiplot(e.pcx,col=rgb(0,0,0,0.1),cex=c(0.8,0.05),
    xlab=paste("PC1: ", round(sum(e.pcx$sdev[1]^2)/e.mvar.clr, 3), sep=""),
    ylab=paste("PC2: ", round(sum(e.pcx$sdev[2]^2)/e.mvar.clr, 3), sep=""),
    xlab.col=mydata$fit.cluster, arrow.len=0.05, var.axes=F, expand=0.8, scale=0)
}
```



K-means clustering supports only 2 groups. In fact, only the first plot needs to be shown to support this: maybe include the barplot?

Samples 3A, 31S, 13B and 15B are near the midpoint on the right, and 19A and 8B are near the midpoint on the left. Five of these six microbiomes are atypical, and the other 19A is exclusively *L. iners*, but phenotypically is BV. We next exclude these six samples and re-examine the partitioning.

```
e.min <- e

e.min[, "003A"] <- NULL
e.min[, "31S"] <- NULL
e.min[, "013B"] <- NULL
e.min[, "015B"] <- NULL
e.min[, "019A"] <- NULL
e.min[, "008B"] <- NULL

e.min.n0.CZM <- cmultRepl(t(e.min), label=0, method="CZM")
```

```
## No. corrected values: 545
```

```
# turn this into a centered log-ratio transform
# samples are by row
# remember that apply by row rotates the data
# R is terrible, so we need to use t() again

e.min.clr <- t( apply(e.min.n0.CZM, 1, function(x){log(x) - mean(log(x))}) ) )
```

```

e.min.mvar.clr <- mvar(e.min.clr)

e.min.pcx <- prcomp(e.min.clr)

# TODO color samples by most abundant organism
biplot(e.min.pcx, cex=c(0.5,0.05), col=c("black",rgb(1,0,0,0.2)), var.axes=F,
       xlab=paste("PC1: ", round(sum(e.min.pcx$sdev[1]^2)/e.min.mvar.clr, 3)),
       ylab=paste("PC2: ", round(sum(e.min.pcx$sdev[2]^2)/e.min.mvar.clr, 3)),
       scale=0
)

# filter the taxon table by what is in e.min
o.min <- o[,colnames(e.min)]
o.CZM <- cmultRepl(t(o.min), label=0, method="CZM")
o.clr <- t( apply(o.CZM, 1, function(x){log(x) - mean(log(x))}) )

# overplot taxon correlations with first two components
cor.scale.0 <- max(abs(1/range(e.min.pcx$rotation[,1:2])))

points(cor(e.min.pcx$x[,1], o.clr[,1], method="kendall") /cor.scale.0,
       cor(e.min.pcx$x[,2], o.clr[,1], method="kendall")/cor.scale.0, pch="I",
       col="DodgerBlue")
points(cor(e.min.pcx$x[,1], o.clr[,6], method="kendall") /cor.scale.0,
       cor(e.min.pcx$x[,2], o.clr[,6], method="kendall")/cor.scale.0, pch="L",
       col="DarkBlue")

points(cor(e.min.pcx$x[,1], o.clr[,2], method="kendall") /cor.scale.0,
       cor(e.min.pcx$x[,2], o.clr[,2], method="kendall")/cor.scale.0, pch="C", col="cyan")

points(cor(e.min.pcx$x[,1], o.clr[,10], method="kendall") /cor.scale.0,
       cor(e.min.pcx$x[,2], o.clr[,10], method="kendall")/cor.scale.0, pch="M", col="green")


points(cor(e.min.pcx$x[,1], o.clr[,8], method="kendall") /cor.scale.0,
       cor(e.min.pcx$x[,2], o.clr[,8], method="kendall")/cor.scale.0, pch="P", col="purple")

points(cor(e.min.pcx$x[,1], o.clr[,7], method="kendall") /cor.scale.0,
       cor(e.min.pcx$x[,2], o.clr[,7], method="kendall")/cor.scale.0, pch="G", col="red")

points(cor(e.min.pcx$x[,1], o.clr[,9], method="kendall") /cor.scale.0,
       cor(e.min.pcx$x[,2], o.clr[,9], method="kendall")/cor.scale.0, pch="A",
       col="Goldenrod")

points(cor(e.min.pcx$x[,1], o.clr[,12], method="kendall") /cor.scale.0,
       cor(e.min.pcx$x[,2], o.clr[,12], method="kendall")/cor.scale.0, pch="S", col="coral3")

```

```
##### phi with Bayesian estimation
```

```
# the problem is that a low phi can easily arise between functions
# that have the same distribution of 0 values across samples
# so we need to estimate the value of 0 (and other low-count functions)
```

```
# generate random DIR clr instances with ALDEx2
e.x <- aldex.clr(e.min)
```

```
## [1] "operating in serial mode"
```

```
# calculate phi divided by number of random instances
e.min.sma.df <- aldex.phi(e.x)
```


```
# find the set of connections with phi less than some value
# we choose an arbitrary cutoff, but it is higher after Bayesian estimation
# obviously
phi.cutoff <- 0.025
```

```
e.min.sma.lo.phi <- subset(e.min.sma.df, phi < phi.cutoff)
```

```
# generate a graphical object
g <- graph.data.frame(e.min.sma.lo.phi, directed=FALSE)
```

```
# overview of all the proportional relationships
# this can take a long time!!!
```

```
plot(g, layout=layout.fruchterman.reingold.grid(g, weight=0.05/E(g)$phi), vertex.size=1,
     vertex.color="black", vertex.label=NA)
```



```
# # get the clusters from the graph object
g.clust <- clusters(g)

# write them to a file

# # data frame containing the names and group memberships of each cluster
g.df <- data.frame(Systematic.name=V(g)$name, cluster=g.clust$membership,
  cluster.size=g.clust$size[g.clust$membership])
write.table(g.df, file="g.df.txt", sep="\t", quote=F, col.names=NA)

max.clust <- induced.subgraph(g, which(g.clust$membership %in% which(g.clust$size ==
  max(g.clust$size))))
max.clust.names <- V(max.clust)$name

par(mfrow=c(1,1))

biplot(e.min.pcx,col=c("black", rgb(1,0,0,0.2)),cex=c(0.8,0.05),
var.axes=F, scale=0)

#nms <- rownames(g.df)[g.df$cluster==max.clust.names]

#points(e.min.pcx$rotation[max.clust.names,][,1],e.min.pcx$rotation[max.clust.names,][,2],
# col=rgb(0,0,1,0.2),pch=19,cex=0.2)

colours <- c("steelblue3", "skyblue1", "indianred1", "mediumpurple1", "olivedrab3",
  "pink", "#FFED6F", "mediumorchid3", "ivory2", "tan1", "aquamarine3", "#C0C0C0",
  "royalblue4", "mediumvioletred", "#999933", "#666699", "#CC9933", "#006666", "#3399FF",
  "#993300", "#CCCC99", "#666666", "#FFCC66", "#6699CC", "#663366", "#9999CC", "#CCCCCC",
  "#669999", "#CCCC66", "#CC6600", "#9999FF", "#0066CC", "#99CCCC", "#999999", "#FFCC00",
  "#009999", "#FF9900", "#999966", "#66CCCC", "#339966", "#CCCC33", "#EDED"
)


big <- g.df[which(g.df$cluster.size >= 2),]
colnames(big) <- colnames(g.df)

lev <- factor(big$cluster)
for(i in as.numeric(levels(lev))) {
  nms <- rownames(big)[big$cluster==i]
```

```

#print(rownames(big)[big$cluster==i])
#print("")
points(e.min.pcx$rotation[nms,][,1],e.min.pcx$rotation[nms,][,2], col=colours[i],
       pch=19, cex=0.3)
}

```



Here the vast majority of low phi values come from core functions that are constant ratios in all samples, but not differential. A few smaller groups are both differential and compositionally associated.

Incorporating the Bayesian estimate of 0 gives a much clearer picture, than using point estimates. When point estimates are used, then the majority of the clusters are near 0,0 - simply because of sets of functions that share the same distribution of 0 across samples.