

Meta Data Key

Column	Meaning
Orig.ident	Which pool the individual it was sequenced in
Tank	Which tank the individual was housed in
Trigger	Weather sex change was triggered by male pairing or female removal.
Prev_tank	The previous tank the males that became dominants were in
Condition	Experiment (dominants) vs Control (males and females)
Status	The sex of the individual, with levels NRM, M, D, E, NF, F. When doing stats, subset to m, d, and f
Status_Long	The full name of status, "Non reproductive male", 'Male' etc.
Time_Day_2	Amount of time the individual spent in the nest during parental care assay on day 2 (day 1 is acclimation). Here, a higher number indicates a more male-like behavior
Behaviors_Day_2	Same as above except with parental care behaviors
Change_Length	Change in length from start of experiment to 6-mos later. Data only recorded for dominants, expandeds, and new female
Change_Mass	Same as above except with weight
Log_11KT	Log of serum 11kt recorded in each individual. Missing values in 1M and 1F
Average_Area_2.5x	Average area of the gonad at 2.5x magnification in pixels. Data not recorded for females
Total_Slides_With_Gonad	Total number of slides that contained gonad. Data not recorded for females
Estimated_Volume_2.5x	Gonadal volume estimate given average area and number of slides in pixels. Not present in females
Log10_Volume	Log10 of volume estimate
Percent_Testicular	Percent of gonad made up of testicular tissue. Not recorded for females
Percent_Ovarian	Same as above with ovarian tissue
Testicular estimate	Percent testicular * estimated volume

Log10_Testicular_Estimate	Log10 of testicular estimate
---------------------------	------------------------------

This is excluding things you already know e.g., (nCount_RNA etc.)

A note on status:

The experiment began with 3 statuses, Males (M), females (F), and dominants (D).

Dominants are previous males that had been paired with another male for 6 months, a timepoint at which they will be somewhere in the middle of sex change. After gonadal analysis, we found 2 dominants had distinct expanded gonads with no testicular tissue, these dominants were labeled as expandeds (E). We hypothesize these fish are very late in sex change. We also found one dominant with vitellogenic oocytes, meaning it had fully completed sex change. We relevelled this one to be NF (new female). We also have a non reproductive male (NRM) in the dataset. In all stats, I only use M, D, and F. We can discuss if this is right or not but these are the 3 largest groups. In clustering, I use all individuals because I don't think it makes a difference.