

Quantum Transport in Nanoporous Graphene

Bachelor defense

CHRISTOFFER VENDELBO SØRENSEN (s163965)



JUNE 25, 2019

Outline

Introduction

Project aim and nanoporous graphene

Tight Binding

π -orbitals, π -electrons and the TB approximation

The Hamiltonian

Onsites, hops and the full TB Hamiltonian

Green's functions and recursion

Green's matrix, recursion and LDOS

Transmission

Device Green's functions, left/right geometry, rate matrices and spectral functions.

Transmission in 2D


Exploring GNR bridges

Para-O₄-NPG, Para-(OH)₄-NPG, Meta-O₂-NPG,
Meta-(OH)₂-NPG

“Development of tight-binding routines in Python in order to understand electron transport in novel nanoporous graphene devices (NPGs)”


Nanoporous graphene

- ▶ Planar graphene sheets
- ▶ Periodically removed atoms
- ▶ Ribbons and bridges
- ▶ Ballistic electron movement
- ▶ Potential for controlling currents on nanoscale



π -orbitals and π -electrons

- ▶ In plane electrons are bound
- ▶ 1 p_z -electron per site
- ▶ “Tightly bound” hops between sites



TB approximation


- ▶ Electrons tightly bound to sites
- ▶ Hops with potential
- ▶ Average electron energy on site
- ▶ The Hamiltonian is a hop matrix

$$V_{pp\pi} = \langle \phi_{\pi}(m) | \hat{\mathbf{H}} | \phi_{\pi}(n) \rangle$$

$$\epsilon_0 = \langle \phi_{\pi}(i) | \hat{\mathbf{H}} | \phi_{\pi}(i) \rangle$$


Hamiltonian for benzene

$$\mathbf{H} = V_{pp\pi} \begin{array}{c} \\ 1 \\ 2 \\ 3 \\ 4 \\ 5 \\ 6 \end{array} \begin{array}{cccccc} 1 & 2 & 3 & 4 & 5 & 6 \\ \left(\begin{array}{cccccc} 0 & 1 & 0 & 0 & 0 & 1 \\ 1 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 & 0 & 1 \\ 1 & 0 & 0 & 0 & 1 & 0 \end{array} \right) \end{array}$$



Creating the first Hamiltonian


- Atom coordinates
- Interatomic distances
- Applying potential



```
33     h = np.zeros((xyz.shape[0], xyz.shape[0])) # Empty matrix
34     for i in range(xyz.shape[0]): # Take an atomic coordinate
35         for j in range(xyz.shape[0]): # Take another atomic coordinate
36             h[i, j] = LA.norm(np.subtract(xyz[i], xyz[j])) # Measure distances
37     h = np.where(h < 1.6, Vppi, 0) # Replace distances under 1.6 angstrom with Vppi
38     h = np.subtract(h, Vppi * np.identity(xyz.shape[0])) # Remove the diagonal
```


Hopping matrices

- Shift by lattice vector
- Resulting matrices: $\mathbf{h}_0, \mathbf{V}, \mathbf{V}^\dagger$




Full Hamiltonian and first band plots

$$\mathbf{H}(k_x, k_y)\phi_k = \epsilon_n(k_x, k_y)\phi_k$$
$$\mathbf{H}(k_x, k_y) = \mathbf{h}_0 + (\mathbf{V}_x e^{-ik_x} + \mathbf{V}_x^\dagger e^{ik_x} + \mathbf{V}_y e^{-ik_y} + \mathbf{V}_y^\dagger e^{ik_y} \\ + \mathbf{V}_{xy} e^{-ik_x} e^{-ik_y} + \mathbf{V}_{xy}^\dagger e^{ik_x} e^{ik_y})$$

```
73     Ham = Ham + (V1 * np.exp(-1.0j * x)  # Onsite hops and hops in x
74               + np.transpose(V1) * np.exp(1.0j * x)  # Hops in -x
75               + V2 * np.exp(-1.0j * y)  # Hops in y
76               + np.transpose(V2) * np.exp(1.0j * y)  # Hops in -y
77               + V3 * np.exp(-1.0j * x) * np.exp(-1.0j * y)  # Hops in both x and y
78               + np.transpose(V3) * np.exp(1.0j * x) * np.exp(1.0j * y))  # Hops in b
79     e = LA.eigh(Ham)[0]  # Eigen energies from the Hamiltonian
80     v = LA.eigh(Ham)[1]  # Eigen vectors from the Hamiltonian
```

Full Hamiltonian and first band plots

$$\begin{aligned} \text{X: } \mathbf{H}_X &= \mathbf{h}_0 + (\mathbf{V}_x e^{ik_x} + \mathbf{V}_x^\dagger e^{-ik_x} + \mathbf{V}_y + \mathbf{V}_y^\dagger \\ &\quad + \mathbf{V}_{xy} e^{ik_x} + \mathbf{V}_{xy}^\dagger e^{-ik_x}) \\ \text{Y: } \mathbf{H}_Y &= \mathbf{h}_0 + (\mathbf{V}_x + \mathbf{V}_x^\dagger + \mathbf{V}_y e^{-ik_y} + \mathbf{V}_y^\dagger e^{ik_y} \\ &\quad + \mathbf{V}_{xy} e^{-ik_y} + \mathbf{V}_{xy}^\dagger e^{ik_y}) \end{aligned}$$



- Solution to the Scödinger Equation
- Propagator

$$[(E + i\eta)\mathbf{1} - \mathbf{H}]\mathbf{G}(E) = \mathbf{1}$$

↓

$$\mathbf{G}(E) = \mathbf{1}([(E + i\eta)\mathbf{1} - \mathbf{H}])^{-1}$$

► Semi-infinite chain

$$\begin{pmatrix} z\mathbf{1} - \mathbf{H}_c & -\mathbf{V}^\dagger \\ -\mathbf{V} & (z - \varepsilon')\mathbf{1} \end{pmatrix} \begin{pmatrix} \mathbf{X} & \mathbf{G}_{0c} \\ \mathbf{G}_{c0} & \mathbf{G}_{00} \end{pmatrix} = \begin{pmatrix} \mathbf{1} & \mathbf{0} \\ \mathbf{0} & \mathbf{1} \end{pmatrix}$$

$$\begin{aligned} \mathbf{G}_{00}(z) &= \left[(z - \varepsilon') - \mathbf{V}(z\mathbf{1} - \mathbf{H}_c)\mathbf{V}^\dagger \right]^{-1} \\ &= (z - \varepsilon' - \Sigma(z))^{-1} \end{aligned}$$

where

$$z = E + i\eta$$

$$a_0 = \mathbf{V}^\dagger, \quad b_0 = \mathbf{V}$$

$$e_{s0} = \mathbf{h}_s, \quad e_0 = \mathbf{h}$$

in loop:

$$a_1 = a_0 \times g_0 \times a_0$$

$$b_1 = b_0 \times g_0 \times b_0$$

$$e_1 = e_0 + a_0 \times g_0 \times b_0 + b_0 \times g_0 \times a_0$$

$$e_{1s} = e_{0s} + a_0 \times g_0 \times b_0$$

$$g_1 = (z - e_1)^{-1}$$


$$\Sigma_R = e_s - h$$

$$\Sigma_L = e - h - \Sigma_R$$

$$\mathbf{G}_{00} = (z - e_s)^{-1}$$

Recursion

```
92     while np.max(np.abs(a0)) > 1e-6: # Loop with hop matrix as threshold
93         ag = a0 @ g0 # Product defined here and used multiple times
94         a1 = ag @ a0 # New hop matrix (transposed)
95         bg = b0 @ g0 # Product defined here and used multiple times
96         b1 = bg @ b0 # New hop matrix
97         e1 = e0 + ag @ b0 + bg @ a0 # New onsite for "other cells"
98         es1 = es0 + ag @ b0 # New onsite
99         g1 = LA.inv(z - e1) # New Green's function
100        a0 = a1 # Overwrite old variable
101        b0 = b1 # Overwrite old variable
102        e0 = e1 # Overwrite old variable
103        es0 = es1 # Overwrite old variable
104        g0 = g1 # Overwrite old variable
105        q = q + 1 # Counter (for diagnostic purposes)
106    e, es = e0, es0 # Define the onsite Hamiltonians
107    SelfER = es - h # Self-energy from the right
108    SelfEL = e - h - SelfER # Self-energy from the left
109    G00 = LA.inv(z - es) # Green's functions
```




```

64 G00 = np.zeros((En.shape[0]), dtype=complex) # Empty data matrix for Green's functions
65 for i in range(En.shape[0]): # Loop iterating over energies
66     G, SelfER, SelfEL = RecursionRoutine(En[i], h, V, eta) # Invoking the RecursionRoutine
67     G = np.diag(G) # The Green's functions for each site is in the diagonal of the G matrix
68     G00[i] = G[4] # Chosen Green's function (here the 4th site)


```

Translating from system to matrices

Transmission is the probability of an electron being transported through a specific region for a specific range of energies.



The left and right self-energy



```

210     for i in En: # Iteration over each energy
211         gl, scrap, SEL = RecursionRoutine(i, HL, VL, eta=eta) # From the left
212         gr, SER, scrap = RecursionRoutine(i, HR, VR, eta=eta) # From the right
    
```

$$\mathbf{G}_D = [\mathbf{1}(E + i\eta) - \mathbf{H}_D - \mathbf{\Sigma}_L(E) - \mathbf{\Sigma}_R(E)]^{-1}$$
$$\mathbf{\Gamma}_{L,R} = i \left(\mathbf{\Sigma}_{L,R} - \mathbf{\Sigma}_{L,R}^\dagger \right)$$

```
225     GD["GD{:d}".format(q)] = scp.linalg.inv( # Device Green's functions are saved
226         scp.identity(HD.shape[0]) * (i + eta) - HD - SEL - SER) # in a dictionary.
227     GammaL["GammaL{:d}".format(q)] = 1j * (SEL - SEL.conj().transpose()) # Rate ma
228     GammaR["GammaR{:d}".format(q)] = 1j * (SER - SER.conj().transpose()) # likewis
```


Getting Transmission

$$T(E) = \text{Tr} \left[\Gamma_R \mathbf{G}_D \Gamma_L \mathbf{G}^\dagger \right] (E)$$

```
240     for i in range(En.shape[0]): # Iteration for every energy point.
241         T[i] = np.trace((GammaR["GammaR{:d}".format(i)] @ GD["GD{:d}".format(
242             i)] @ GammaL["GammaL{:d}".format(i)] @ GD["GD{:d}".format(i)].conj(
243             ).transpose()).todense()) # Calculate transmission (equation V.9).
```

Getting Transmission

- Text1
- Text2
- Text3



Introduction
oo

Tight Binding
ooo

The Hamiltonian
ooo

Green's functions and recursion
ooo

Transmission
oo●
ooo

Exploring GNR bridges
ooooo

Questions
o


Transmission in 2D

$$\mathbf{H} = \mathbf{h} + \mathbf{V}e^{ik_{\perp}} + \mathbf{V}^{\dagger}e^{i(-k_{\perp})}$$

```



50 h, p = Onsite(xyz=xyz, Vppi=-1, f=1) # Calculate hopping
51 V = Hop(xyz=xyz, xyz1=xyz + np.array([0, UY, 0]), k_perp=-1) # Calculate hops
52 print('Number of hopping elements: {}'.format(np.sum(np.abs(V)))) # Number of hops
53 Ham = h + V * np.exp(1j * i) + np.transpose(V) * np.exp(-1j * i) # Hamiltonian for ith k-point

```



Code validity

$$k = \frac{\pi}{2}$$



Introduction
○○

Tight Binding
○○○

The Hamiltonian
○○○

Green's functions and recursion
○○○



Transmission
○○○
●●○

Exploring GNR bridges
○○○○○

Questions
○

Code validity

$$k = \pi$$



Introduction
○○

Tight Binding
○○○

The Hamiltonian
○○○

Green's functions and recursion
○○○



Transmission
○○○
●●●

Exploring GNR bridges
○○○○○

Questions
○

Code validity

$$k = \text{AVG}\left(0, \frac{\pi}{2}, \pi\right)$$



Introduction
○○

Tight Binding
○○○

The Hamiltonian
○○○


Green's functions and recursion
○○○

Transmission
○○○
●●●



Exploring GNR bridges
○○○○○

Questions
○

Summary of code structure




Para and meta bridges



► Test


Para and meta bridges

► Text1





Para-O₄-NPG

► text



(a)



(e)

Introduction
○○

Tight Binding
○○○

The Hamiltonian
○○○

Green's functions and recursion
○○○


Transmission
○○○
○○○

Exploring GNR bridges
○●○○○



Questions
○

Para-(OH)₄-NPG


► text





(a)



Meta-O₂-NPG



(a)



(e)

► text

Introduction
○○

Tight Binding
○○○

The Hamiltonian
○○○

Green's functions and recursion
○○○


Transmission
○○○
○○○

Exploring GNR bridges
○○○●○○



Questions
○

Meta-(OH)₂-NPG

► text



(a)



“Development of tight-binding routines in Python in order to understand electron transport in novel nanoporous graphene devices (NPGs)”

Questions

CHRISTOFFER VENDELBO SØRENSEN (s163965)



JUNE 25, 2019

Introduction
oo

Tight Binding
ooo

The Hamiltonian
ooo

Green's functions and recursion
ooo

Transmission
ooo
ooo

Exploring GNR bridges
oooooo

Questions
●

► Appendix