

# VHEGEN: A vibronic Hamiltonian expansion generator for trigonal and tetragonal polyatomic systems

Robert A. Lang

Riley J. Hickman

Tao Zeng

Thank you for using **VHEGEN**, the **V**-ibronic **H**-amiltonian **E**-xpansion **GEN**-erator for trigonal and tetragonal polyatomic systems. This is a **VHEGEN** output file compiled by **pdf<sub>l</sub>at<sub>e</sub>x**. If the **VHEGEN** package was used in research resulting in a publication, please reference the article in *Computer Physics Communications* which describes the program ([doi here]). Additional information regarding the matrix element expansion process, including the independent matrix element eigenvalues, their root formulas and constraints, and their transformation to the real basis (if applicable), can be found in the **log** output file. For questions, bugs, or comments, please contact robert.lang@mail.utoronto.ca.

## Contents

<b>1</b>	<b>Vibronic interaction</b>	<b>2</b>
<b>2</b>	<b>Vibronic Hamiltonian operator in the complex <math>E</math> basis</b>	<b>2</b>
<b>3</b>	<b>Matrix element expansions in the complex <math>E</math> basis</b>	<b>2</b>
3.1	Order: 2 . . . . .	2
3.2	Order: 3 . . . . .	2
<b>4</b>	<b>Vibronic Hamiltonian operator in the real <math>E</math> basis</b>	<b>3</b>
<b>5</b>	<b>Matrix element expansions in the real <math>E</math> basis</b>	<b>3</b>
5.1	Order: 2 . . . . .	3
5.2	Order: 3 . . . . .	4

# 1 Vibronic interaction

$(E + A) \otimes (e + a)$  in  $C_3$

## 2 Vibronic Hamiltonian operator in the complex $E$ basis

$$\hat{H} = (|+\rangle \quad |-\rangle \quad |A\rangle) \begin{pmatrix} 0 & 0 & H_{+A} \\ 0 & 0 & H_{-A} \\ H_{A+} & H_{A-} & 0 \end{pmatrix} \begin{pmatrix} \langle +| \\ \langle -| \\ \langle A| \end{pmatrix}$$

## 3 Matrix element expansions in the complex $E$ basis

### 3.1 Order: 2

Number of terms:  $H_{+A}$ : 4.

**Polar e-coordinates:**

$$H_{+A}^{(2)} = -b_{0,0,2}^i \rho^2 \sin(2\phi) + ib_{0,0,2}^i \rho^2 \cos(2\phi) + b_{1,0,-1}^i \rho z \sin(\phi) + ib_{1,0,-1}^i \rho z \cos(\phi) + ib_{0,0,2}^r \rho^2 \sin(2\phi) + b_{0,0,2}^r \rho^2 \cos(2\phi) - ib_{1,0,-1}^r \rho z \sin(\phi) + b_{1,0,-1}^r \rho z \cos(\phi)$$

$$H_{A+}^{(2)} = -b_{0,0,2}^i \rho^2 \sin(2\phi) - ib_{0,0,2}^i \rho^2 \cos(2\phi) + b_{1,0,-1}^i \rho z \sin(\phi) - ib_{1,0,-1}^i \rho z \cos(\phi) - ib_{0,0,2}^r \rho^2 \sin(2\phi) + b_{0,0,2}^r \rho^2 \cos(2\phi) + ib_{1,0,-1}^r \rho z \sin(\phi) + b_{1,0,-1}^r \rho z \cos(\phi)$$

$$H_{-A}^{(2)} = -b_{0,0,2}^i \rho^2 \sin(2\phi) - ib_{0,0,2}^i \rho^2 \cos(2\phi) + b_{1,0,-1}^i \rho z \sin(\phi) - ib_{1,0,-1}^i \rho z \cos(\phi) - ib_{0,0,2}^r \rho^2 \sin(2\phi) + b_{0,0,2}^r \rho^2 \cos(2\phi) + ib_{1,0,-1}^r \rho z \sin(\phi) + b_{1,0,-1}^r \rho z \cos(\phi)$$

$$H_{A-}^{(2)} = -b_{0,0,2}^i \rho^2 \sin(2\phi) + ib_{0,0,2}^i \rho^2 \cos(2\phi) + b_{1,0,-1}^i \rho z \sin(\phi) + ib_{1,0,-1}^i \rho z \cos(\phi) + ib_{0,0,2}^r \rho^2 \sin(2\phi) + b_{0,0,2}^r \rho^2 \cos(2\phi) - ib_{1,0,-1}^r \rho z \sin(\phi) + b_{1,0,-1}^r \rho z \cos(\phi)$$

**Cartesian e-coordinates:**

$$H_{+A}^{(2)} = -2b_{0,0,2}^i xy + ib_{0,0,2}^i (x-y)(x+y) + ib_{1,0,-1}^i xz + b_{1,0,-1}^i yz + 2ib_{0,0,2}^r xy + b_{0,0,2}^r (x-y)(x+y) + b_{1,0,-1}^r xz - ib_{1,0,-1}^r yz$$

$$H_{A+}^{(2)} = -2b_{0,0,2}^i xy - ib_{0,0,2}^i (x-y)(x+y) - ib_{1,0,-1}^i xz + b_{1,0,-1}^i yz - 2ib_{0,0,2}^r xy + b_{0,0,2}^r (x-y)(x+y) + b_{1,0,-1}^r xz + ib_{1,0,-1}^r yz$$

$$H_{-A}^{(2)} = -2b_{0,0,2}^i xy - ib_{0,0,2}^i (x-y)(x+y) - ib_{1,0,-1}^i xz + b_{1,0,-1}^i yz - 2ib_{0,0,2}^r xy + b_{0,0,2}^r (x-y)(x+y) + b_{1,0,-1}^r xz + ib_{1,0,-1}^r yz$$

$$H_{A-}^{(2)} = -2b_{0,0,2}^i xy + ib_{0,0,2}^i (x-y)(x+y) + ib_{1,0,-1}^i xz + b_{1,0,-1}^i yz + 2ib_{0,0,2}^r xy + b_{0,0,2}^r (x-y)(x+y) + b_{1,0,-1}^r xz - ib_{1,0,-1}^r yz$$

### 3.2 Order: 3

Number of terms:  $H_{+A}$ : 6.

### Polar e-coordinates:

$$H_{+A}^{(3)} = b_{0,2,-1}^i \rho^3 \sin(\phi) + ib_{0,2,-1}^i \rho^3 \cos(\phi) - b_{1,0,2}^i \rho^2 z \sin(2\phi) + ib_{1,0,2}^i \rho^2 z \cos(2\phi) + b_{2,0,-1}^i \rho z^2 \sin(\phi) \\ + ib_{2,0,-1}^i \rho z^2 \cos(\phi) - ib_{0,2,-1}^r \rho^3 \sin(\phi) + b_{0,2,-1}^r \rho^3 \cos(\phi) + ib_{1,0,2}^r \rho^2 z \sin(2\phi) + b_{1,0,2}^r \rho^2 z \cos(2\phi) \\ - ib_{2,0,-1}^r \rho z^2 \sin(\phi) + b_{2,0,-1}^r \rho z^2 \cos(\phi)$$

$$H_{+A}^{(3)} = b_{0,2,-1}^i \rho^3 \sin(\phi) - ib_{0,2,-1}^i \rho^3 \cos(\phi) - b_{1,0,2}^i \rho^2 z \sin(2\phi) - ib_{1,0,2}^i \rho^2 z \cos(2\phi) + b_{2,0,-1}^i \rho z^2 \sin(\phi) \\ - ib_{2,0,-1}^i \rho z^2 \cos(\phi) + ib_{0,2,-1}^r \rho^3 \sin(\phi) + b_{0,2,-1}^r \rho^3 \cos(\phi) - ib_{1,0,2}^r \rho^2 z \sin(2\phi) + b_{1,0,2}^r \rho^2 z \cos(2\phi) \\ + ib_{2,0,-1}^r \rho z^2 \sin(\phi) + b_{2,0,-1}^r \rho z^2 \cos(\phi)$$

$$H_{-A}^{(3)} = b_{0,2,-1}^i \rho^3 \sin(\phi) - ib_{0,2,-1}^i \rho^3 \cos(\phi) - b_{1,0,2}^i \rho^2 z \sin(2\phi) - ib_{1,0,2}^i \rho^2 z \cos(2\phi) + b_{2,0,-1}^i \rho z^2 \sin(\phi) \\ - ib_{2,0,-1}^i \rho z^2 \cos(\phi) + ib_{0,2,-1}^r \rho^3 \sin(\phi) + b_{0,2,-1}^r \rho^3 \cos(\phi) - ib_{1,0,2}^r \rho^2 z \sin(2\phi) + b_{1,0,2}^r \rho^2 z \cos(2\phi) \\ + ib_{2,0,-1}^r \rho z^2 \sin(\phi) + b_{2,0,-1}^r \rho z^2 \cos(\phi)$$

$$H_{A-}^{(3)} = b_{0,2,-1}^i \rho^3 \sin(\phi) + ib_{0,2,-1}^i \rho^3 \cos(\phi) - b_{1,0,2}^i \rho^2 z \sin(2\phi) + ib_{1,0,2}^i \rho^2 z \cos(2\phi) + b_{2,0,-1}^i \rho z^2 \sin(\phi) \\ + ib_{2,0,-1}^i \rho z^2 \cos(\phi) - ib_{0,2,-1}^r \rho^3 \sin(\phi) + b_{0,2,-1}^r \rho^3 \cos(\phi) + ib_{1,0,2}^r \rho^2 z \sin(2\phi) + b_{1,0,2}^r \rho^2 z \cos(2\phi) \\ - ib_{2,0,-1}^r \rho z^2 \sin(\phi) + b_{2,0,-1}^r \rho z^2 \cos(\phi)$$

### Cartesian e-coordinates:

$$H_{+A}^{(3)} = ib_{0,2,-1}^i x(x^2 + y^2) + b_{0,2,-1}^i y(x^2 + y^2) - 2b_{1,0,2}^i xyz + ib_{1,0,2}^i z(x - y)(x + y) + ib_{2,0,-1}^i xz^2 + b_{2,0,-1}^i yz^2 \\ + b_{0,2,-1}^r x(x^2 + y^2) - ib_{0,2,-1}^r y(x^2 + y^2) + 2ib_{1,0,2}^r xyz + b_{1,0,2}^r z(x - y)(x + y) + b_{2,0,-1}^r xz^2 - ib_{2,0,-1}^r yz^2$$

$$H_{+A}^{(3)} = -ib_{0,2,-1}^i x(x^2 + y^2) + b_{0,2,-1}^i y(x^2 + y^2) - 2b_{1,0,2}^i xyz - ib_{1,0,2}^i z(x - y)(x + y) - ib_{2,0,-1}^i xz^2 + b_{2,0,-1}^i yz^2 \\ + b_{0,2,-1}^r x(x^2 + y^2) + ib_{0,2,-1}^r y(x^2 + y^2) - 2ib_{1,0,2}^r xyz + b_{1,0,2}^r z(x - y)(x + y) + b_{2,0,-1}^r xz^2 + ib_{2,0,-1}^r yz^2$$

$$H_{-A}^{(3)} = -ib_{0,2,-1}^i x(x^2 + y^2) + b_{0,2,-1}^i y(x^2 + y^2) - 2b_{1,0,2}^i xyz - ib_{1,0,2}^i z(x - y)(x + y) - ib_{2,0,-1}^i xz^2 + b_{2,0,-1}^i yz^2 \\ + b_{0,2,-1}^r x(x^2 + y^2) + ib_{0,2,-1}^r y(x^2 + y^2) - 2ib_{1,0,2}^r xyz + b_{1,0,2}^r z(x - y)(x + y) + b_{2,0,-1}^r xz^2 + ib_{2,0,-1}^r yz^2$$

$$H_{A-}^{(3)} = ib_{0,2,-1}^i x(x^2 + y^2) + b_{0,2,-1}^i y(x^2 + y^2) - 2b_{1,0,2}^i xyz + ib_{1,0,2}^i z(x - y)(x + y) + ib_{2,0,-1}^i xz^2 + b_{2,0,-1}^i yz^2 \\ + b_{0,2,-1}^r x(x^2 + y^2) - ib_{0,2,-1}^r y(x^2 + y^2) + 2ib_{1,0,2}^r xyz + b_{1,0,2}^r z(x - y)(x + y) + b_{2,0,-1}^r xz^2 - ib_{2,0,-1}^r yz^2$$

## 4 Vibronic Hamiltonian operator in the real $E$ basis

$$\hat{H} = (|X\rangle \quad |Y\rangle \quad |A\rangle) \begin{pmatrix} 0 & 0 & H_{XA} \\ 0 & 0 & H_{YA} \\ H_{AX} & H_{AY} & 0 \end{pmatrix} \begin{pmatrix} \langle X| \\ \langle Y| \\ \langle A| \end{pmatrix}$$

## 5 Matrix element expansions in the real $E$ basis

### 5.1 Order: 2

Number of terms:  $H_{XA}$ : 4 (all from  $H_{+A}$ ),  $H_{YA}$ : 4 (all from  $H_{+A}$ ).

**Polar e-coordinates:**

$$H_{YA}^{(2)} = -\sqrt{2} (b_{0,0,2}^i \rho^2 \cos(2\phi) + b_{1,0,-1}^i \rho z \cos(\phi) + b_{0,0,2}^r \rho^2 \sin(2\phi) - b_{1,0,-1}^r \rho z \sin(\phi))$$

$$H_{AX}^{(2)} = \sqrt{2} (-b_{0,0,2}^i \rho^2 \sin(2\phi) + b_{1,0,-1}^i \rho z \sin(\phi) + b_{0,0,2}^r \rho^2 \cos(2\phi) + b_{1,0,-1}^r \rho z \cos(\phi))$$

$$H_{XA}^{(2)} = \sqrt{2} (-b_{0,0,2}^i \rho^2 \sin(2\phi) + b_{1,0,-1}^i \rho z \sin(\phi) + b_{0,0,2}^r \rho^2 \cos(2\phi) + b_{1,0,-1}^r \rho z \cos(\phi))$$

$$H_{AY}^{(2)} = -\sqrt{2} (b_{0,0,2}^i \rho^2 \cos(2\phi) + b_{1,0,-1}^i \rho z \cos(\phi) + b_{0,0,2}^r \rho^2 \sin(2\phi) - b_{1,0,-1}^r \rho z \sin(\phi))$$

**Cartesian e-coordinates:**

$$H_{YA}^{(2)} = -\sqrt{2} (b_{0,0,2}^i (x-y)(x+y) + b_{1,0,-1}^i xz + 2b_{0,0,2}^r xy - b_{1,0,-1}^r yz)$$

$$H_{AX}^{(2)} = \sqrt{2} (-2b_{0,0,2}^i xy + b_{1,0,-1}^i yz + b_{0,0,2}^r (x-y)(x+y) + b_{1,0,-1}^r xz)$$

$$H_{XA}^{(2)} = \sqrt{2} (-2b_{0,0,2}^i xy + b_{1,0,-1}^i yz + b_{0,0,2}^r (x-y)(x+y) + b_{1,0,-1}^r xz)$$

$$H_{AY}^{(2)} = -\sqrt{2} (b_{0,0,2}^i (x-y)(x+y) + b_{1,0,-1}^i xz + 2b_{0,0,2}^r xy - b_{1,0,-1}^r yz)$$

## 5.2 Order: 3

Number of terms:  $H_{XA}$ : 6 (all from  $H_{+A}$ ),  $H_{YA}$ : 6 (all from  $H_{+A}$ ).

**Polar e-coordinates:**

$$H_{YA}^{(3)} = -\sqrt{2} (b_{0,2,-1}^i \rho^3 \cos(\phi) + b_{1,0,2}^i \rho^2 z \cos(2\phi) + b_{2,0,-1}^i \rho z^2 \cos(\phi) - b_{0,2,-1}^r \rho^3 \sin(\phi) + b_{1,0,2}^r \rho^2 z \sin(2\phi) - b_{2,0,-1}^r \rho z^2 \sin(\phi))$$

$$H_{AX}^{(3)} = \sqrt{2} (b_{0,2,-1}^i \rho^3 \sin(\phi) - b_{1,0,2}^i \rho^2 z \sin(2\phi) + b_{2,0,-1}^i \rho z^2 \sin(\phi) + b_{0,2,-1}^r \rho^3 \cos(\phi) + b_{1,0,2}^r \rho^2 z \cos(2\phi) + b_{2,0,-1}^r \rho z^2 \cos(\phi))$$

$$H_{XA}^{(3)} = \sqrt{2} (b_{0,2,-1}^i \rho^3 \sin(\phi) - b_{1,0,2}^i \rho^2 z \sin(2\phi) + b_{2,0,-1}^i \rho z^2 \sin(\phi) + b_{0,2,-1}^r \rho^3 \cos(\phi) + b_{1,0,2}^r \rho^2 z \cos(2\phi) + b_{2,0,-1}^r \rho z^2 \cos(\phi))$$

$$H_{AY}^{(3)} = -\sqrt{2} (b_{0,2,-1}^i \rho^3 \cos(\phi) + b_{1,0,2}^i \rho^2 z \cos(2\phi) + b_{2,0,-1}^i \rho z^2 \cos(\phi) - b_{0,2,-1}^r \rho^3 \sin(\phi) + b_{1,0,2}^r \rho^2 z \sin(2\phi) - b_{2,0,-1}^r \rho z^2 \sin(\phi))$$

**Cartesian e-coordinates:**

$$H_{YA}^{(3)} = -\sqrt{2} \left( b_{0,2,-1}^i x (x^2 + y^2) + b_{1,0,2}^i z (x - y) (x + y) + b_{2,0,-1}^i x z^2 - b_{0,2,-1}^r y (x^2 + y^2) + 2b_{1,0,2}^r x y z - b_{2,0,-1}^r y z^2 \right)$$

$$H_{AX}^{(3)} = \sqrt{2} \left( b_{0,2,-1}^i y (x^2 + y^2) - 2b_{1,0,2}^i x y z + b_{2,0,-1}^i y z^2 + b_{0,2,-1}^r x (x^2 + y^2) + b_{1,0,2}^r z (x - y) (x + y) + b_{2,0,-1}^r x z^2 \right)$$

$$H_{XA}^{(3)} = \sqrt{2} \left( b_{0,2,-1}^i y (x^2 + y^2) - 2b_{1,0,2}^i x y z + b_{2,0,-1}^i y z^2 + b_{0,2,-1}^r x (x^2 + y^2) + b_{1,0,2}^r z (x - y) (x + y) + b_{2,0,-1}^r x z^2 \right)$$

$$H_{AY}^{(3)} = -\sqrt{2} \left( b_{0,2,-1}^i x (x^2 + y^2) + b_{1,0,2}^i z (x - y) (x + y) + b_{2,0,-1}^i x z^2 - b_{0,2,-1}^r y (x^2 + y^2) + 2b_{1,0,2}^r x y z - b_{2,0,-1}^r y z^2 \right)$$