

A sample computation

Altan Erdnigor

March 21, 2024

1 Notation

- p a prime number.
- $\mathbf{SL}_3(\mathbb{Z})$ the special linear group over \mathbb{Z} .
- Γ_p the p th congruence subgroup of $\mathbf{SL}_3(\mathbb{Z})$.
- $Z_G(x)$ the centralizer of $x \in G$.

•

$$A = \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & p+2 \\ 0 & 1 & 2p \end{pmatrix} \in \mathbf{SL}_3(\mathbb{Z}) \quad (1)$$

•

$$\tilde{A} = \mathbf{Id} + pA = \begin{pmatrix} 1 & 0 & p \\ p & 1 & 2p+p^2 \\ 0 & p & 1+2p^2 \end{pmatrix} \in \Gamma_p \quad (2)$$

- If C is a matrix, $\chi_C(\lambda) := \det(\lambda \mathbf{Id} - C)$ is the characteristics polynomial.
- $f(t) := \chi_A(t) = t^3 - 2pt^2 - (p+2)t - 1$.

2 Intro

Recall Lucas primes (see https://en.wikipedia.org/wiki/Lucas_number#Lucas_primes, <https://t5k.org/top20/page.php?id=48>)

$$2, 3, 7, 11, 29, 47, 199, 521, 2207, 3571, 9349, 3010349, 54018521, 370248451, 6643838879, \dots \quad (3)$$

In this notes we establish the following results:

1. The regularor of \tilde{A} grows as $\approx \ln^2(p)$.
That is, $\text{Reg}(\mathbb{Q}(\alpha)/\mathbb{Q}) \approx \ln^2(p)$ for α a root of $\chi_{\tilde{A}}$.
2. The index of the centralizers $[Z_{\mathbf{SL}_3(\mathbb{Z})}(\tilde{A}) : Z_{\Gamma_p}(\tilde{A})]$ grows as $O(\ln p)$ for p a Lucas prime.

This might be interesting.

3 Regulator

We refer to Keith Conrad's write-up on Dirichlet's unit theorem and regulators [1] for the definitions. The current proof mimics the proof of Theorem 5.12 of Conrad.

Notice that

$$\begin{aligned}\chi_{\tilde{A}}(\lambda) &= \det(\lambda \mathbf{Id} - \tilde{A}) \\ &= \det(\lambda \mathbf{Id} - (\mathbf{Id} + pA)) = \det((\lambda - 1)\mathbf{Id} - pA) \\ &= p^3 \det\left(\frac{\lambda - 1}{p} \mathbf{Id} - A\right) = p^3 \chi_A\left(\frac{\lambda - 1}{p}\right).\end{aligned}\quad (4)$$

Hence adding the root of χ_A or $\chi_{\tilde{A}}$ result in the same field; therefore we reduce to showing that $\text{Reg}(\mathbb{Q}(\alpha)/\mathbb{Q}) \approx \ln^2(p)$ for α a root of $\chi_A(t) = f(t) = t^3 - (2pt^2 + (p+2)t + 1)$.

Lemma 3.1. $\mathbb{Q}(\alpha)/\mathbb{Q}$ is totally real of degree 3 for primes $p \neq 2$.

Proof. $f(t)$ is irreducible over \mathbb{Q} ; indeed, by the rational roots theorem it's sufficient to check ± 1 :

$$f(1) = -3p - 2, f(-1) = -p.$$

A simple computation shows that the discriminant of $f(t)$ equals

$$\text{disc}_f(p) = 4p^4 - 12p^3 + 4p^2 - 24p + 5.$$

If $p > 4$ the discriminant $\text{disc}_f(p) > 0$ is positive. Therefore the cubic extension is totally real. \square

Proposition 3.2. $\mathbb{Z}[\alpha]^* = \{\pm \alpha^a (2\alpha + 1)^b \mid a, b \in \mathbb{Z}\}$.

Note that $\alpha, 2\alpha + 1$ are not necessarily fundamental units in $\mathbb{Q}(\alpha)/\mathbb{Q}$ as we don't claim that the ring of integers of $\mathbb{Q}(\alpha)/\mathbb{Q}$ coincides with $\mathbb{Z}[\alpha]$.

Proof. Note that $f(\alpha) = 0$ implies

$$\alpha(\alpha^2 - 2p\alpha - (p+2)) = 1 \quad (5)$$

$$(1 + 2\alpha)(1 + p\alpha) = \alpha^3 \quad (6)$$

It shows that $\alpha, 1 + 2\alpha$ are indeed units.

Let $\alpha_1 > \alpha_2 > \alpha_3 \in \mathbb{R}$ be the three different roots of f . We shall compute them approximately.

$$\begin{aligned}\alpha_1 &= 2p + \frac{1}{2} + O\left(\frac{1}{p}\right), \\ \alpha_2 &= -\frac{1}{p} + O\left(\frac{1}{p^4}\right), \\ \alpha_3 &= -\frac{1}{2} + O\left(\frac{1}{p}\right).\end{aligned}$$

Remark 3.3. A computation shows that for $p = 10000$ we have

$$\alpha_1 = 20000.5000874981, \alpha_2 = -0.000100000000000100, \alpha_3 = -0.499987498124648.$$

It is not important that p is not a prime in this case as the estimate works for any sufficiently large p .

By the definition of the regulator we have

$$\begin{aligned} \text{Reg}(\alpha, 2\alpha + 1) &= \left| \begin{array}{cc} \ln |\alpha_1| & \ln |\alpha_2| \\ \ln |2\alpha_1 + 1| & \ln |2\alpha_2 + 1| \end{array} \right| \\ &\approx \left| \begin{array}{cc} \ln |2p + \frac{1}{2}| & \ln |\frac{-1}{p}| \\ \ln |4p + 2| & \ln |\frac{-2}{p} + 1| \end{array} \right| = \ln(2p + \frac{1}{2})(\ln(p - 2) - \ln(p)) + \ln(4p + 2) \ln(p) \\ &= \ln(2p + \frac{1}{2}) \ln(p - 2) - \ln(2p + \frac{1}{2}) \ln(p) + \ln(4p + 2) \ln(p). \quad (7) \end{aligned}$$

Therefore $\text{Reg}(\alpha, 2\alpha + 1) > 0$ for all prime p .

Hence $\alpha, 2\alpha + 1$ are independent units.

It is left to prove that they are fundamental units in $\mathbb{Z}[\alpha]$. By Corollary 5.9 from Conrad it is sufficient to check

$$\frac{16 \text{Reg}(\alpha, 2\alpha + 1)}{(\ln(\text{disc}_f / 4))^2} < 2.$$

Substituting, we obtain

$$\frac{16 \text{Reg}(\alpha, 2\alpha + 1)}{(\ln(\text{disc}_f / 4))^2} \approx \frac{16(\ln(2p + \frac{1}{2}) \ln(p - 2) - \ln(2p + \frac{1}{2}) \ln(p) + \ln(4p + 2) \ln(p))}{(\ln((p^4 + 2p^3 - 5p^2 - 6p - 23)/4))^2}.$$

Asymptotically, the latter equals

$$\xrightarrow{p \rightarrow \infty} \frac{16 \ln(p)^2}{(\ln(p^4))^2} = 1.$$

Therefore it is < 2 for big enough p , QED. \square

Remark 3.4. One can do the estimate more carefully, but for now postpone it.

Remark 3.5. We just proved that the regulator is approximately

$$\ln(2p + \frac{1}{2}) \ln(p - 2) - \ln(2p + \frac{1}{2}) \ln(p) + \ln(4p + 2) \ln(p),$$

which is close to $\ln^2 p$ we wanted from the beginning.

4 Centralizers

Proposition 4.1. The centralizer of \tilde{A} in $\mathbf{SL}_3(\mathbb{Z})$ is generated by $A, A + \mathbf{Id}$.

$$Z_{\mathbf{SL}_3(\mathbb{Z})}(\tilde{A}) = \{\pm A^a (2A + \mathbf{Id})^b \mid a, b \in \mathbb{Z}\}.$$

Proof. Since \tilde{A} is regular, its centralizer in $\mathbf{Mat}_3(\mathbb{C})$ is $\mathbb{C}\langle \mathbf{Id}, A, A^2 \rangle$. Now,

$$\mathbb{C}\langle \mathbf{Id}, A, A^2 \rangle \cap \mathbf{SL}_3(\mathbb{Z}) \subset \mathbb{Z}\langle \mathbf{Id}, A, A^2 \rangle.$$

Indeed,

$$\mathbf{Id} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, A = \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & p+2 \\ 0 & 1 & 2p \end{pmatrix}, A^2 = \begin{pmatrix} 0 & 1 & 2p \\ 0 & p+1 & 2p^2+4p+1 \\ 1 & p & 4p^2+p+2 \end{pmatrix}, \quad (8)$$

Considering the first matrix column we see that if a complex combination has integer coefficients, it is in fact integer combination.

Moreover, the centralizer of \tilde{A} is a group, therefore it lies inside the multiplicative group of $\mathbb{Z}[A]$

$$Z_{\mathbf{SL}_3(\mathbb{Z})}(\tilde{A}) \subset \mathbb{C}\langle \mathbf{Id}, A, A^2 \rangle \cap \mathbf{SL}_3(\mathbb{Z}) \subset \mathbb{Z}[A]^*.$$

There is an isomorphism of \mathbb{Z} -algebras $\mathbb{Z}[A] \simeq \mathbb{Z}[x]/(f(x)) = \mathbb{Z}[\alpha]$. Applying Proposition 3.2 end the proof

$$Z_{\mathbf{SL}_3(\mathbb{Z})}(\tilde{A}) \subset \mathbb{Z}[A]^* = \{\pm A^a(2A + \mathbf{Id})^b \mid a, b \in \mathbb{Z}\}.$$

□

We are to study the centralizer of \tilde{A} in Γ_p .

$$Z_{\Gamma_p}(\tilde{A}) \subset Z_{\mathbf{SL}_3(\mathbb{Z})}(\tilde{A}) \cong \mathbb{Z}^2.$$

In general this subgroup can be difficult to describe; that leads to considering *Lucas primes*.

Proposition 4.2. *Let p be a Lucas prime. Let k be the integer part of $\log_\phi p$ where ϕ is the golden ratio.*

The centralizer $Z_{\Gamma_p}(\tilde{A})$ contains \tilde{A} and A^{4k} .

Proof. The only thing to prove is that $A^{4k} \in \Gamma_p$.

It suffices to show that the eigenvalues of $A \pmod{p}$ are $4k$ -th roots of unity. Computing

$$\chi_A(t) = f(t) \equiv t^3 - 2t - 1 = (t+1)(t^2 - t - 1) \pmod{p},$$

shows that it is left to work with the golden ratio in \mathbb{F}_p which we denote by ϕ_p . That is, $\phi_p \in \mathbb{F}_{p^2}$ satisfies $\phi_p^2 - \phi_p - 1 = 0$.

By the definition of a Lucas number we have

$$p = \phi^k + (-\phi)^{-k} \quad (9)$$

where ϕ is a root of $x^2 - x - 1$.

The RHS of (9) being invariant under the change $\phi \rightarrow (-\phi)^{-1}$ manifests it as a symmetric polynomial in the roots of $x^2 - x - 1$, thus having a presentation

$$\phi^k + (-\phi)^{-k} = P(\phi, (-\phi)^{-1}),$$

where P is a *universal* polynomial. This observation justifies that the Equation (9) can be taken modulo p to have the form

$$0 = \phi_p^k + (-\phi_p)^{-k},$$

which implies

$$1 = \phi_p^{4k}.$$

□

Theorem 4.3. *The index of the centralizers is bounded by $4 \log_\phi p$*

$$[Z_{\mathbf{SL}_3(\mathbb{Z})}(\tilde{A}) : Z_{\Gamma_p}(\tilde{A})] \leq 4 \log_\phi p.$$

In particular, it grows as $O(\ln p)$.

Proof. Identify $Z_{\mathbf{SL}_3(\mathbb{Z})}(\tilde{A}) \cong \mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}^2$ as in Proposition 4.1.

Observe that by (6) we have

$$1 + p\alpha = \alpha^3(2\alpha + 1)^{-1}.$$

By the previous Proposition $Z_{\Gamma_p}(\tilde{A})$ contains $\begin{pmatrix} 3 \\ -1 \end{pmatrix}, \begin{pmatrix} 4k \\ 0 \end{pmatrix}$; Clearly, the index

$$\mathbb{Z} \left\langle \begin{pmatrix} 3 \\ -1 \end{pmatrix}, \begin{pmatrix} 4k \\ 0 \end{pmatrix} \right\rangle \subset \mathbb{Z}^2,$$

equals $4k \approx 4 \log_\phi p$ and the index $[Z_{\Gamma_p}(\tilde{A}) : Z_{\mathbf{SL}_3(\mathbb{Z})}(\tilde{A})]$ has to divide it. □

References

- [1] Keith Conrad. Dirichlet's unit theorem. <https://kconrad.math.uconn.edu/blurbs/gradnumthy/unittheorem.pdf>.