

A sample computation

Altan Erdnigor

June 27, 2024

Contents

1 Notation

- p a prime number.
- $\mathbf{SL}_3(\mathbb{Z})$ the special linear group over \mathbb{Z} .
- Γ_p the p th congruence subgroup of $\mathbf{SL}_3(\mathbb{Z})$.
- $Z_G(x)$ the centralizer of $x \in G$.

•

$$A = \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & p+2 \\ 0 & 1 & 2p \end{pmatrix} \in \mathbf{SL}_3(\mathbb{Z}) \quad (1)$$

•

$$\tilde{A} = \mathbf{Id} + pA = \begin{pmatrix} 1 & 0 & p \\ p & 1 & 2p+p^2 \\ 0 & p & 1+2p^2 \end{pmatrix} \in \Gamma_p \quad (2)$$

- If C is a matrix, $\chi_C(\lambda) := \det(\lambda \mathbf{Id} - C)$ is the characteristics polynomial.
- $f(t) := \chi_A(t) = t^3 - 2pt^2 - (p+2)t - 1$.

2 Intro

Recall the Lucas primes (see https://en.wikipedia.org/wiki/Lucas_number#Lucas_primes, <https://t5k.org/top20/page.php?id=48>)

$$2, 3, 7, 11, 29, 47, 199, 521, 2207, 3571, 9349, 3010349, 54018521, 370248451, 6643838879, \dots \quad (3)$$

In this notes we establish the following results:

1. The regularor of \tilde{A} grows as $\approx \ln^2(p)$.
That is, $\text{Reg}(\mathbb{Q}(\alpha)/\mathbb{Q}) \approx \ln^2(p)$ for α a root of $\chi_{\tilde{A}}$.
2. The index of the centralizers $[Z_{\mathbf{SL}_3(\mathbb{Z})}(\tilde{A}) : Z_{\Gamma_p}(\tilde{A})]$ grows as $O(\ln p)$ for p a Lucas prime.

This might be interesting.

3 Regulator

We refer to Keith Conrad's write-up on Dirichlet's unit theorem and regulators [?] for the definitions. The current proof mimics the proof of Theorem 5.12 of Conrad.

Notice that

$$\begin{aligned}\chi_{\tilde{A}}(\lambda) &= \det(\lambda \mathbf{Id} - \tilde{A}) \\ &= \det(\lambda \mathbf{Id} - (\mathbf{Id} + pA)) = \det((\lambda - 1)\mathbf{Id} - pA) \\ &= p^3 \det\left(\frac{\lambda - 1}{p} \mathbf{Id} - A\right) = p^3 \chi_A\left(\frac{\lambda - 1}{p}\right).\end{aligned}\quad (4)$$

Hence adding the root of χ_A or $\chi_{\tilde{A}}$ result in the same field; therefore we reduce to showing that $\text{Reg}(\mathbb{Q}(\alpha)/\mathbb{Q}) \approx \ln^2(p)$ for α a root of $\chi_A(t) = f(t) = t^3 - (2pt^2 + (p+2)t + 1)$.

Lemma 3.1. $\mathbb{Q}(\alpha)/\mathbb{Q}$ is totally real of degree 3 for primes $p \neq 2$.

Proof. $f(t)$ is irreducible over \mathbb{Q} ; indeed, by the rational roots theorem it's sufficient to check ± 1 :

$$f(1) = -3p - 2, f(-1) = -p.$$

A simple computation shows that the discriminant of $f(t)$ equals

$$\text{disc}_f(p) = 4p^4 - 12p^3 + 4p^2 - 24p + 5.$$

If $p > 4$ the discriminant $\text{disc}_f(p) > 0$ is positive. Therefore the cubic extension is totally real. \square

Proposition 3.2. $\mathbb{Z}[\alpha]^* = \{\pm \alpha^a (2\alpha + 1)^b \mid a, b \in \mathbb{Z}\}$.

Note that $\alpha, 2\alpha + 1$ are not necessarily fundamental units in $\mathbb{Q}(\alpha)/\mathbb{Q}$ as we don't claim that the ring of integers of $\mathbb{Q}(\alpha)/\mathbb{Q}$ coincides with $\mathbb{Z}[\alpha]$.

Proof. Note that $f(\alpha) = 0$ implies

$$\alpha(\alpha^2 - 2p\alpha - (p+2)) = 1 \quad (5)$$

$$(1 + 2\alpha)(1 + p\alpha) = \alpha^3 \quad (6)$$

It shows that $\alpha, 1 + 2\alpha$ are indeed units.

Let $\alpha_1 > \alpha_2 > \alpha_3 \in \mathbb{R}$ be the three different roots of f . We shall compute them approximately.

$$\begin{aligned}\alpha_1 &= 2p + \frac{1}{2} + O\left(\frac{1}{p}\right), \\ \alpha_2 &= -\frac{1}{p} + O\left(\frac{1}{p^4}\right), \\ \alpha_3 &= -\frac{1}{2} + O\left(\frac{1}{p}\right).\end{aligned}$$

Remark 3.3. A computation shows that for $p = 10000$ we have

$$\alpha_1 = 20000.5000874981, \alpha_2 = -0.000100000000000100, \alpha_3 = -0.499987498124648.$$

It is not important that p is not a prime in this case as the estimate works for any sufficiently large p .

By the definition of the regulator we have

$$\begin{aligned} \text{Reg}(\alpha, 2\alpha + 1) &= \left| \begin{array}{cc} \ln |\alpha_1| & \ln |\alpha_2| \\ \ln |2\alpha_1 + 1| & \ln |2\alpha_2 + 1| \end{array} \right| \\ &\approx \left| \begin{array}{cc} \ln |2p + \frac{1}{2}| & \ln |\frac{-1}{p}| \\ \ln |4p + 2| & \ln |\frac{-2}{p} + 1| \end{array} \right| = \ln(2p + \frac{1}{2})(\ln(p - 2) - \ln(p)) + \ln(4p + 2) \ln(p) \\ &= \ln(2p + \frac{1}{2}) \ln(p - 2) - \ln(2p + \frac{1}{2}) \ln(p) + \ln(4p + 2) \ln(p). \quad (7) \end{aligned}$$

Therefore $\text{Reg}(\alpha, 2\alpha + 1) > 0$ for all prime p .

Hence $\alpha, 2\alpha + 1$ are independent units.

It is left to prove that they are fundamental units in $\mathbb{Z}[\alpha]$. By Corollary 5.9 from Conrad it is sufficient to check

$$\frac{16 \text{Reg}(\alpha, 2\alpha + 1)}{(\ln(\text{disc}_f / 4))^2} < 2.$$

Substituting, we obtain

$$\frac{16 \text{Reg}(\alpha, 2\alpha + 1)}{(\ln(\text{disc}_f / 4))^2} \approx \frac{16(\ln(2p + \frac{1}{2}) \ln(p - 2) - \ln(2p + \frac{1}{2}) \ln(p) + \ln(4p + 2) \ln(p))}{(\ln((p^4 + 2p^3 - 5p^2 - 6p - 23)/4))^2}.$$

Asymptotically, the latter equals

$$\xrightarrow{p \rightarrow \infty} \frac{16 \ln(p)^2}{(\ln(p^4))^2} = 1.$$

Therefore it is < 2 for big enough p , QED. \square

Remark 3.4. One can do the estimate more carefully, but for now postpone it.

Remark 3.5. We just proved that the regulator is approximately

$$\ln(2p + \frac{1}{2}) \ln(p - 2) - \ln(2p + \frac{1}{2}) \ln(p) + \ln(4p + 2) \ln(p),$$

which is close to $\ln^2 p$ we wanted from the beginning.

4 Centralizers

Proposition 4.1. The centralizer of \tilde{A} in $\mathbf{SL}_3(\mathbb{Z})$ is generated by $A, 2A + \mathbf{Id}$.

$$Z_{\mathbf{SL}_3(\mathbb{Z})}(\tilde{A}) = \{\pm A^a (2A + \mathbf{Id})^b \mid a, b \in \mathbb{Z}\}.$$

Proof. Since \tilde{A} is regular, its centralizer in $\mathbf{Mat}_3(\mathbb{C})$ is $\mathbb{C}\langle \mathbf{Id}, A, A^2 \rangle$. Now,

$$\mathbb{C}\langle \mathbf{Id}, A, A^2 \rangle \cap \mathbf{SL}_3(\mathbb{Z}) \subset \mathbb{Z}\langle \mathbf{Id}, A, A^2 \rangle.$$

Indeed,

$$\mathbf{Id} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, A = \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & p+2 \\ 0 & 1 & 2p \end{pmatrix}, A^2 = \begin{pmatrix} 0 & 1 & 2p \\ 0 & p+1 & 2p^2+4p+1 \\ 1 & p & 4p^2+p+2 \end{pmatrix}, \quad (8)$$

Considering the first matrix column we see that if a complex combination has integer coefficients, it is in fact integer combination.

Moreover, the centralizer of \tilde{A} is a group, therefore it lies inside the multiplicative group of $\mathbb{Z}[A]$

$$Z_{\mathbf{SL}_3(\mathbb{Z})}(\tilde{A}) \subset \mathbb{C}\langle \mathbf{Id}, A, A^2 \rangle \cap \mathbf{SL}_3(\mathbb{Z}) \subset \mathbb{Z}[A]^*.$$

There is an isomorphism of \mathbb{Z} -algebras $\mathbb{Z}[A] \simeq \mathbb{Z}[x]/(f(x)) = \mathbb{Z}[\alpha]$. Applying Proposition ?? end the proof

$$Z_{\mathbf{SL}_3(\mathbb{Z})}(\tilde{A}) \subset \mathbb{Z}[A]^* = \{\pm A^a(2A + \mathbf{Id})^b \mid a, b \in \mathbb{Z}\}.$$

□

We are to study the centralizer of \tilde{A} in Γ_p .

$$Z_{\Gamma_p}(\tilde{A}) \subset Z_{\mathbf{SL}_3(\mathbb{Z})}(\tilde{A}) \cong \mathbb{Z}^2.$$

In general this subgroup can be difficult to describe; that leads to considering *Lucas primes*.

4.1 Lucas primes give small degree $\log(p)$

Proposition 4.2. *Let p be a Lucas prime. Let k be the integer part of $\log_\phi p$ where ϕ is the golden ratio.*

The centralizer $Z_{\Gamma_p}(\tilde{A})$ contains \tilde{A} and A^{4k} .

Proof. The only thing to prove is that $A^{4k} \in \Gamma_p$.

It suffices to show that the eigenvalues of $A \pmod{p}$ are $4k$ -th roots of unity. Computing

$$\chi_A(t) = f(t) \equiv t^3 - 2t - 1 = (t+1)(t^2 - t - 1) \pmod{p},$$

shows that it is left to work with the golden ratio in \mathbb{F}_p which we denote by ϕ_p . That is, $\phi_p \in \mathbb{F}_{p^2}$ satisfies $\phi_p^2 - \phi_p - 1 = 0$.

By the definition of a Lucas number we have

$$p = \phi^k + (-\phi)^{-k} \quad (9)$$

where ϕ is a root of $x^2 - x - 1$.

The RHS of (9) being invariant under the change $\phi \rightarrow (-\phi)^{-1}$ manifests it as a symmetric polynomial in the roots of $x^2 - x - 1$, thus having a presentation

$$\phi^k + (-\phi)^{-k} = P(\phi, (-\phi)^{-1}),$$

where P is a *universal* polynomial. This observation justifies that the Equation (??) can be taken modulo p to have the form

$$0 = \phi_p^k + (-\phi_p)^{-k},$$

which implies

$$1 = \phi_p^{4k}.$$

□

Theorem 4.3. *The index of the centralizers is bounded by $4 \log_\phi p$*

$$[Z_{\mathbf{SL}_3(\mathbb{Z})}(\tilde{A}) : Z_{\Gamma_p}(\tilde{A})] \leq 4 \log_\phi p.$$

In particular, it grows as $O(\ln p)$.

Proof. Identify $Z_{\mathbf{SL}_3(\mathbb{Z})}(\tilde{A}) \cong \mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}^2$ as in Proposition ??.

Observe that by (??) we have

$$1 + p\alpha = \alpha^3(2\alpha + 1)^{-1}.$$

By the previous Proposition $Z_{\Gamma_p}(\tilde{A})$ contains $\begin{pmatrix} 3 \\ -1 \end{pmatrix}, \begin{pmatrix} 4k \\ 0 \end{pmatrix}$; Clearly, the index

$$\mathbb{Z} \left\langle \begin{pmatrix} 3 \\ -1 \end{pmatrix}, \begin{pmatrix} 4k \\ 0 \end{pmatrix} \right\rangle \subset \mathbb{Z}^2,$$

equals $4k \approx 4 \log_\phi p$ and the index $[Z_{\Gamma_p}(\tilde{A}) : Z_{\mathbf{SL}_3(\mathbb{Z})}(\tilde{A})]$ has to divide it. □

4.2 Bounds on the degree $\log(p) \prec \deg \prec p$

Motivated by geometric considerations, we call the degree \deg the index $[Z_{\Gamma_p}(\tilde{A}) : Z_{\mathbf{SL}_3(\mathbb{Z})}(\tilde{A})]$.

Proposition 4.4. *We have $O(\log(p)) \leq \deg \leq O(p)$.*

Proof. The proof of the previous Theorem shows that $Z_{\mathbf{SL}_3(\mathbb{Z})}(\tilde{A})$ is generated by $\alpha^3(2\alpha + 1)^{-1}, \alpha^{\deg}$.

So, the goal is to show that the multiplicative order of ϕ_p is bounded by $\log(p)$ from below and by p from above.

First let us show $\log(p) < \deg$. Indeed, lifting ϕ_p to ϕ we notice that

$$\phi_p^k = 1 \Rightarrow (-\phi_p^{-1})^k = (-1)^k \Rightarrow \phi^k + (-\phi^{-1})^k - (1 + (-1)^k) \in p\mathbb{Z}.$$

Thus the degree $\deg > \log(p) - \epsilon$ for p big enough and some small ϵ .

Finally, we shall show $\deg \leq O(p)$. What we show in reality is that if $\sqrt{5} \in \mathbb{F}_p$, the degree \deg divides $p - 1$, $\deg | p - 1$ and if $\sqrt{5} \notin \mathbb{F}_p$, $\deg | 2p + 2$.

Indeed, the first part is clear by Fermat's little theorem. To show the second, let's decompose

$$x^2 - x - 1 = (x - \alpha_1)(x - \alpha_2),$$

where $\alpha_1, \alpha_2 \in \mathbb{F}_{p^2}$. The Frobenius automorphism $\text{Frob} : t \mapsto t^p$ acts on the roots α_1, α_2 by permuting them; thus we have $\alpha_1^p = \alpha_2 = -1/\alpha_1$ implying $\alpha_1^{2p+2} = (-1)^2 = 1$. □

5 Enter family

Let $f_{a,b}(t) = t^3 - apt^2 - (bp + a)t - b$. Here $a, b \in \mathbb{Z}$. Note that $f_{2,1}(t) = f(t)$ is the characteristic polynomial of A .

We can now define

$$A_{a,b} = \begin{pmatrix} 0 & 0 & b \\ 1 & 0 & bp + a \\ 0 & 1 & ap \end{pmatrix} \in \mathbf{Mat}_3(\mathbb{Z}), \quad (10)$$

$$\tilde{A}_{a,b} = \mathbf{Id} + pA_{a,b} = \begin{pmatrix} 1 & 0 & bp \\ p & 1 & bp^2 + ap \\ 0 & p & 1 + ap^2 \end{pmatrix} \in \Gamma_p. \quad (11)$$

Here $\tilde{A}_{a,b}$ is indeed invertible thanks to

$$\begin{aligned} \det(\tilde{A}_{a,b}) &= p^3 \det\left(\frac{1}{p}\mathbf{Id} + A_{a,b}\right) = -p^3 \det\left(\frac{-1}{p}\mathbf{Id} - A_{a,b}\right) \\ &= -p^3 f_{a,b}\left(\frac{-1}{p}\right) = 1 + ap^2 - (bp + a)p^2 + bp^3 = 1. \end{aligned} \quad (12)$$

Actually, a generic element in Γ_p is conjugate to $\tilde{A}_{a,b}$.

Proposition 5.1. *An element $\tilde{B} \in \Gamma_p$ is either*

1. \mathbf{Id} ;
2. conjugate to

$$\begin{pmatrix} 1 & p & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

over \mathbb{Q} ;

3. conjugate to $\tilde{A}_{a,b}$ over \mathbb{Q} for some $a, b \in \mathbb{Z}$.

The proof goes by applying the Frobenius normal form over \mathbb{Q} .

Proof. Define $B = \frac{1}{p}(\tilde{B} - \mathbf{Id}) \in \mathbf{Mat}_3(\mathbb{Z})$.

There are three cases: the minimal polynomial minpoly_B of B is of degree 1, 2 or 3.

1. $\deg \text{minpoly}_B = 1$. Hence \tilde{B} is scalar, hence \mathbf{Id} ;
2. $\deg \text{minpoly}_B = 2$. Hence $\frac{\chi_B}{\text{minpoly}_B}$ is linear, thus dividing minpoly_B ; we obtain χ_B decomposes into a product of linear multiples over \mathbb{Z}

$$\chi_B(\lambda) = (\lambda - a)^2(\lambda - b), a, b \in \mathbb{Z}.$$

The condition $-p^3 \chi_B\left(\frac{-1}{p}\right) = \chi_{\tilde{B}}(0) = 1$ now reads

$$(1 + ap)^2(1 + bp) = 1,$$

thus $a, b = 0$ if $p > 2$ and B is nilpotent. Therefore \tilde{B} is conjugate to

$$\begin{pmatrix} 1 & p & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}.$$

3. The Frobenius normal form of B consists of the unique 3×3 block.

Pick $C \in \mathbf{GL}_3(\mathbb{Q})$ such that

$$C^{-1}BC = \begin{pmatrix} 0 & 0 & r_0 \\ 1 & 0 & r_1 \\ 0 & 1 & r_2 \end{pmatrix} \in \mathbf{SL}_3(\mathbb{Q}).$$

This matrix is actually integer $C^{-1}BC \in \mathbf{Mat}_3(\mathbb{Z})$. Indeed, the coefficients in the last column are the coefficients of the characteristic polynomial. Therefore, $C^{-1}\tilde{B}C \in \Gamma_p$.

We now show that $C^{-1}\tilde{B}C = \tilde{A}_{a,b}$. This follows from the equation $\det(C^{-1}\tilde{B}C) = 1$ which reads

$$1 + r_2p - r_1p^2 + r_0p^3 = 1,$$

implying

$$r_0 = b, r_1 = bp + a, r_2 = ap,$$

for some $a, b \in \mathbb{Z}$.

□

6 The Unknown

Problem 6.1. *What is the behavior of Reg, \deg for an arbitrary member of the family $\tilde{A}_{a,b}$?*

TODO: * Study the problem for fixed p and changing a, b ; here a, b are such that $\text{Reg } f_{a,b} < p/5$?

$$\text{disc } f_{a,b} = a^2 * p^4 * b^2 - 2 * a^3 * p^3 * b + a^4 * p^2 + 4 * p^3 * b^3 - 6 * a * p^2 * b^2 - 6 * a^2 * p * b + 4 * a^3 - 27 * b^2.$$

7 A Strange example

Let p a prime number with $p \equiv 1 \pmod{3}$. Define

$$A = \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 2p \\ 0 & 1 & p^2 \end{pmatrix} \in \mathbf{SL}_3(\mathbb{Z}) \quad (13)$$

$$\tilde{A} = \mathbf{Id} + pA = \begin{pmatrix} 1 & 0 & p \\ p & 1 & 2p^2 \\ 0 & p & 1 + p^3 \end{pmatrix} \in \Gamma_p \quad (14)$$

The characteristic poly of A is $f(t) := \chi_A(t) = t^3 - p^2t^2 - 2pt - 1$.

Let $\mathbb{Q}(\alpha)$ be the number field obtain by attaching the root of $f(t)$ to \mathbb{Q} .

Proposition 7.1. $\mathbb{Q}(\alpha)/\mathbb{Q}$ is a complex number field of degree 3.

This is terrible as the unit group then is of rank one.

Proof. The discriminant of $f(t)$ equals

$$\text{disc}_f(p) = -4p^3 - 27.$$

The discriminant is always negative $\text{disc}_f(p) < 0$, thus f has only one real root and the other roots are complex and conjugate to each other. \square

Proposition 7.2. $\mathbb{Z}[\alpha]^* = \{\pm \alpha^a (1 + p\alpha)^b \mid a, b \in \mathbb{Z}\}$.

Note that $\alpha, 1 + p\alpha$ are not necessarily fundamental units in $\mathbb{Q}(\alpha)/\mathbb{Q}$ as we don't claim that the ring of integers of $\mathbb{Q}(\alpha)/\mathbb{Q}$ coincides with $\mathbb{Z}[\alpha]$.

Note that $f(\alpha) = 0$ implies

$$\alpha(\alpha^2 - p^2\alpha - 2p) = 1 \quad (15)$$

$$(1 + p\alpha)^2 = \alpha^3 \quad (16)$$

It shows that $\alpha, 1 + p\alpha$ are indeed units.

Let $\alpha_1 \in \mathbb{R}, \alpha_2 = \bar{\alpha}_3 \in \mathbb{C} \setminus \mathbb{R}$ be the three different roots of f . We shall compute them approximately.

$$\begin{aligned} \alpha_1 &= p^2 + \frac{2}{p} + O\left(\frac{1}{p}\right), \\ \alpha_2 &= -p^{-1} + i p^{-\frac{5}{2}} + \frac{3}{2}p^{-4} + O\left(\frac{1}{p^4}\right), \\ \alpha_3 &= -p^{-1} - i p^{-\frac{5}{2}} + \frac{3}{2}p^{-4} + O\left(\frac{1}{p^4}\right). \end{aligned}$$

Remark 7.3. A computation shows that for $p = 1000$ we have

$$\begin{aligned} \alpha_1 &= 1.0000000020000000\text{e}+6, \\ \alpha_2 &= -0.00099999998500000 - i \ 3.162277652\text{e}-8, \\ \alpha_3 &= -0.00099999998500000 + i \ 3.162277652\text{e}-8. \end{aligned}$$

It is not important that p is not a prime in this case as the estimate works for any sufficiently large p .

$$\begin{aligned} f(t) &= t^3 - ap^2t^2 - (a+1)pt - 1. \\ 1 &= (t^2 - ap^2t - (a+1)p)t, t^3 = (pt+1)(apt+1). \end{aligned}$$

8 A Strange example v2

Let p a prime number with $p \equiv 1 \pmod{3}$. Define

$$A = \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & -p^2 \end{pmatrix} \in \mathbf{SL}_3(\mathbb{Z}) \quad (17)$$

$$\tilde{A} = \mathbf{Id} + pA = \begin{pmatrix} 1 & 0 & p \\ p & 1 & 0 \\ 0 & p & 1 - p^3 \end{pmatrix} \in \Gamma_p \quad (18)$$

The characteristic poly of A is $f(t) := \chi_A(t) = t^3 + p^2t^2 - 1$.

Let $\mathbb{Q}(\alpha)$ be the number field obtain by attaching the root of $f(t)$ to \mathbb{Q} .

Proposition 8.1. $\mathbb{Q}(\alpha)/\mathbb{Q}$ is a totally real number field of degree 3.

Proof. The discriminant of $f(t)$ equals

$$\text{disc}_f(p) = 4p^6 - 27.$$

The discriminant is always positive $\text{disc}_f(p) > 0$, thus the number field is totally real. \square

Proposition 8.2. $\mathbb{Z}[\alpha]^* = \{\pm \alpha^a (1 + p\alpha)^b \mid a, b \in \mathbb{Z}\}$.

Note that $\alpha, 1 + p\alpha$ are not necessarily fundamental units in $\mathbb{Q}(\alpha)/\mathbb{Q}$ as we don't claim that the ring of integers of $\mathbb{Q}(\alpha)/\mathbb{Q}$ coincides with $\mathbb{Z}[\alpha]$.

Proof. We have

$$\alpha(\alpha^2 + p^2\alpha) = 1 \tag{19}$$

$$(1 - p\alpha)(1 + p\alpha) = \alpha^3 \tag{20}$$

It shows that $\alpha, 1 + p\alpha$ are indeed units.

Let $\alpha_1 < \alpha_2 < \alpha_3$ be the three different roots of f . We shall compute them approximately.

$$\begin{aligned} \alpha_1 &= -p^2 + O(p^{-4}), \\ \alpha_2 &= -p^{-1} + O(p^{-4}), \\ \alpha_3 &= p^{-1} + O(p^{-4}). \end{aligned}$$

Remark 8.3. A computation shows that for $p = 100$ we have

$$\begin{aligned} \alpha_1 &= -9999.99999999000, \\ \alpha_2 &= -0.01000000500000625, \\ \alpha_3 &= 0.00999999500000625. \end{aligned}$$

It is not important that p is not a prime in this case as the estimate works for any sufficiently large p .

By the definition of the regulator we have

$$\begin{aligned} \text{Reg}(\alpha, p\alpha + 1) &= \begin{vmatrix} \ln |\alpha_1| & \ln |\alpha_3| \\ \ln |p\alpha_1 + 1| & \ln |p\alpha_3 + 1| \end{vmatrix} \\ &\approx \begin{vmatrix} \ln |-p^2| & \ln |\frac{1}{p}| \\ \ln |-p^3 + 1| & \ln |2| \end{vmatrix} = \ln(p^2) \ln 2 + \ln(p^3 - 1) \ln(p) \\ &\approx \ln(p)(3 \ln(p) + 2 \ln 2). \end{aligned} \tag{21}$$

Therefore $\text{Reg}(\alpha, p\alpha + 1) > 0$ for all prime p .

Hence $\alpha, p\alpha + 1$ are independent units.

Remark 8.4. For example, for $p = 547$ SageMath computes the regulator to be approximately $\text{Reg} = 127.978045931846$.

```

p = Primes().unrank(100) # p = 547
R.<x> = PolynomialRing(QQ)
P = x^3 + p^2 * x^2 - 1
if not P.is_irreducible():
    print("NOT IRREDUCIBLE")
    break
K.<a> = QQ.extension(P)
print(K.regulator())

```

Listing 1: SageMath code example

Whereas the above estimate gives 127.97804593184651.

These numbers coincide up to 10^{-12} .

It is left to prove that they are fundamental units in $\mathbb{Z}[\alpha]$. By Corollary 5.9 from Conrad it is sufficient to check

$$\frac{16 \operatorname{Reg}(\alpha, p\alpha + 1)}{(\ln(\operatorname{disc}_f/4))^2} < 2.$$

Substituting, we obtain

$$\frac{16 \operatorname{Reg}(\alpha, 2\alpha + 1)}{(\ln(\operatorname{disc}_f/4))^2} \approx \frac{16 \ln(p)(3 \ln(p) + 2 \ln 2)}{(\ln((4p^6 - 27)/4))^2}.$$

Asymptotically, the latter equals

$$\xrightarrow{p \rightarrow \infty} \frac{48 \ln(p)^2}{(\ln(p^6))^2} = \frac{4}{3}.$$

Therefore it is < 2 for big enough p , QED. \square

Remark 8.5. We just proved that the regulator is approximately

$$\ln(p)(3 \ln(p) + 2 \ln 2),$$

which is close to $\ln^2 p$ we wanted from the beginning.

9 Centralizers

Proposition 9.1. The centralizer of \tilde{A} in $\mathbf{SL}_3(\mathbb{Z})$ is generated by $A, pA + \mathbf{Id}$.

$$Z_{\mathbf{SL}_3(\mathbb{Z})}(\tilde{A}) = \{A^a(pA + \mathbf{Id})^b \mid a, b \in \mathbb{Z}\}.$$

Proof. Since \tilde{A} is regular, its centralizer in $\mathbf{Mat}_3(\mathbb{C})$ is $\mathbb{C}\langle \mathbf{Id}, A, A^2 \rangle$. Now,

$$\mathbb{C}\langle \mathbf{Id}, A, A^2 \rangle \cap \mathbf{SL}_3(\mathbb{Z}) \subset \mathbb{Z}\langle \mathbf{Id}, A, A^2 \rangle.$$

Indeed,

$$\mathbf{Id} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, A = \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & -p^2 \end{pmatrix}, A^2 = \begin{pmatrix} 0 & 1 & -p^2 \\ 0 & 0 & 1 \\ 1 & -p^2 & p^4 \end{pmatrix}, \quad (22)$$

Considering the first matrix column we see that if a complex combination has integer coefficients, it is in fact integer combination.

Moreover, the centralizer of \tilde{A} is a group, therefore it lies inside the multiplicative group of $\mathbb{Z}[A]$

$$Z_{\mathbf{SL}_3(\mathbb{Z})}(\tilde{A}) \subset \mathbb{C} \langle \mathbf{Id}, A, A^2 \rangle \cap \mathbf{SL}_3(\mathbb{Z}) \subset \mathbb{Z}[A]^*.$$

There is an isomorphism of \mathbb{Z} -algebras $\mathbb{Z}[A] \simeq \mathbb{Z}[x]/(f(x)) = \mathbb{Z}[\alpha]$. Applying Proposition ?? end the proof

$$Z_{\mathbf{SL}_3(\mathbb{Z})}(\tilde{A}) \subset \mathbb{Z}[A]^* = \{\pm A^a(pA + \mathbf{Id})^b \mid a, b \in \mathbb{Z}\}.$$

□

We are to study the centralizer of \tilde{A} in Γ_p .

$$Z_{\Gamma_p}(\tilde{A}) \subset Z_{\mathbf{SL}_3(\mathbb{Z})}(\tilde{A}) \cong \mathbb{Z}^2.$$

Proposition 9.2. *The index of the one centralizer inside the other does not depend on p and equals 3.*

Proof. As A is equivalent to the cyclic permutation matrix \pmod{p} it follows $A^3 \in \Gamma_p$.

Thus the smaller centralizer is generated A^3, \tilde{A}

$$Z_{\Gamma_p}(\tilde{A}) = \{A^{3a}(pA + \mathbf{Id})^b \mid a, b \in \mathbb{Z}\},$$

and it's clear that the index is 3.

□