
NATIONAL HURRICANE CENTER TROPICAL CYCLONE REPORT[1]

TROPICAL STORM JULIETTE

(EP102025)

24 – 28 August 2025

John P. Cangialosi
National Hurricane Center
15 September 2025



GOES-18 VISIBLE SATELLITE IMAGE OF TROPICAL STORM JULIETTE AT 1520 UTC 26 AUGUST. IMAGE COURTESY OF NOAA/NESDIS/STAR.

Juliette was a tropical storm that remained over the open waters of the eastern North Pacific Ocean.

[1] This is an abbreviated Tropical Cyclone Report since there were no coastal watches or warnings issued and no direct fatalities reported in association with Juliette.

Tropical Storm Juliette

24 – 28 AUGUST 2025

BEST TRACK

The "best track[2]" positions and intensities are listed in Table 1. The best track chart of Juliette's path is given in Fig. 1, with the wind and pressure histories along with available observations[3] shown in Figs. 2 and 3, respectively.

A station on Clarion Island reported a sustained wind of 37 kt with a gust to 46 kt at 1130 UTC 26 August when Juliette passed the island.

Origin

Juliette formed around 1800 UTC 24 August about 450 n mi south of the southern tip of the Baja California peninsula. Juliette appears to have formed from the combination of a tropical wave that crossed Central America on 19 and 20 August and a disturbance in the monsoon trough.

Peak Intensity and Minimum Pressure

Juliette's peak intensity of 60 kt at 0600 and 1200 UTC 26 August is based on a blend of the satellite intensity estimates from TAFB, SAB, and UW-CIMSS. The minimum central pressure of 994 mb is primarily based on the Knaff-Zehr-Courtney pressure-wind relationship.

CASUALTY AND DAMAGE STATISTICS

There were no reports of damage or casualties associated with Juliette.

---

[2] A digital record of the complete best track, including wind radii, can be found on line at ftp://ftp.nhc.noaa.gov/atcf. Data for the current year's storms are located in the *btk* directory, while previous years' data are located in the *archive* directory.

[3] Observations include subjective satellite-based Dvorak technique intensity estimates from the Tropical Analysis and Forecast Branch (TAFB)) and the Satellite Analysis Branch (SAB), objective Advanced Dvorak Technique (ADT) estimates and Satellite Consensus (SATCON) estimates from the Cooperative Institute for Meteorological Satellite Studies/University of Wisconsin-Madison (UW-CIMSS). Data and imagery from NOAA polar-orbiting satellites including the Advanced Microwave Sounding Unit (AMSU), the NASA Global Precipitation Mission (GPM), the European Space Agency's Advanced Scatterometer (ASCAT), and Defense Meteorological Satellite Program (DMSP) satellites, and the Time-Resolved Observations of Precipitation structure and storm Intensity with a Constellation of Smallsats (TROPICS) satellites, among others, were also useful in constructing the best track of Juliette.

FORECAST AND WARNING VERIFICATION

Table 2 lists the number of hours prior to formation corresponding to the first issuance of an NHC Tropical Weather Outlook (TWO) forecast in each likelihood category. The formation of Juliette was reasonably well anticipated, but the system formed sooner than expected. The precursor disturbance was first highlighted in the 7-day TWO with a low probability of formation 90 h before development. The likelihood was subsequently raised to the medium category 72 h in advance and to the high category 36 h prior to genesis. Lead times for the 2-day outlooks showed good reliability in the low and medium categories. A low chance of development was first indicated 48 h before formation, increasing to the medium category 30 h in advance. However, probabilities never reached the high category prior to genesis. Figure 4 shows composites of the 7-day TWO genesis areas for each likelihood category leading up to the formation of Juliette. Juliette formed near the middle of the predicted genesis region highlighted in the TWO.

A verification of NHC official track and intensity forecasts for Juliette are given in Table 3a and 4a, respectively. A homogeneous comparison of the official track and intensity errors with selected guidance models are given in Table 3b and 4b, respectively. The mean official track forecast errors for Juliette were close to average, a little above the 5-yr means from 12 to 48 h and a little below the means at 60 and 72 h. HFBI and HCCA were generally the best track models for Juliette, beating the official forecasts at most time periods. The NHC mean official intensity errors for Juliette were quite low, besting the 5-yr means at all verifying times, except 24 h. No intensity model consistently outperformed NHC for Juliette.

There were no coastal watches or warnings issued for Juliette.

Table 1. Best track for Tropical Storm Juliette, 24–28 August 2025.

Date/Time (UTC)	Latitude (°N)	Longitude (°W)	Pressure (mb)	Wind Speed (kt)	Stage
24 / 1800	15.7	109.3	1006	30	tropical depression
25 / 0000	16.1	110.5	1005	30	"
25 / 0600	16.5	111.6	1004	35	tropical storm
25 / 1200	16.9	112.8	1002	40	"
25 / 1800	17.3	113.9	999	50	"
26 / 0000	17.7	114.9	997	55	"
26 / 0600	18.2	115.6	995	60	"
26 / 1200	18.7	116.2	994	60	"
26 / 1800	19.4	116.9	996	55	"
27 / 0000	20.4	117.8	997	55	"
27 / 0600	21.4	118.6	999	50	"
27 / 1200	22.5	119.3	1002	45	"
27 / 1800	23.5	120.0	1005	40	"
28 / 0000	24.4	120.6	1006	40	"
28 / 0600	25.1	120.7	1007	35	low
28 / 1200	25.5	120.7	1007	30	"
28 / 1800	25.8	120.8	1008	30	"
29 / 0000	25.9	121.1	1009	25	"
29 / 0600	25.9	121.4	1010	25	"
29 / 1200	25.8	121.8	1011	20	"
29 / 1800	25.6	122.2	1012	20	"
30 / 0000	25.3	122.7	1013	15	"
30 / 0600					dissipated
26 / 1200	18.7	116.2	994	60	maximum wind and minimum pressure

Table 2. Number of hours in advance of formation of Juliette associated with the first NHC Tropical Weather Outlook forecast in the indicated likelihood category. Note that the timings for the "Low" category do not include forecasts of a 0% chance of genesis.

	Hours Before Genesis	
	48-Hour Outlook	**168-Hour Outlook**
Low (<40%)	48	90
Medium (40%-60%)	30	72
High (>60%)	-	36

Table 3a. NHC official (OFCL) and climatology-persistence skill baseline (OCD5) track forecast errors (n mi) for Tropical Storm Juliette, 24–28 August 2025. Mean errors for the previous 5-yr period are shown for comparison. Official errors that are smaller than the 5-yr means are shown in boldface type.

	Forecast Period (h)							
	12	24	36	48	60	72	96	120
OFCL	23.2	38.1	50.8	54.0	**63.9**	**76.4**		
OCD5	40.4	86.0	132.3	172.9	205.1	242.0		
Forecasts	12	10	8	6	4	2		
OFCL (2020-24)	22.4	33.6	43.5	53.2	65.1	79.4	108.6	135.2
OCD5 (2020-24)	38.2	75.3	117.2	161.0	206.0	252.0	340.7	422.7

Table 3b. Homogeneous comparison of selected track forecast guidance models (in n mi) for Tropical Storm Juliette, 24–28 August 2025. Errors smaller than the NHC official forecast are shown in boldface type.

Model ID	Forecast Period (h)							
	12	24	36	48	60	72	96	120
OFCL	23.2	38.1	50.8	54.0	63.9	76.4		
OCD5	40.4	86.0	132.3	172.9	205.1	242.0		
GFSI	**22.0**	**34.2**	55.7	89.6	131.5	155.6		
EMXI	24.1	44.8	68.9	72.2	78.4	**69.6**		
HWFI	**21.7**	**28.4**	**44.4**	64.3	93.5	92.8		
HMNI	**23.1**	**33.8**	**39.3**	**52.5**	84.6	109.8		
HFAI	**22.0**	**34.8**	**41.8**	**49.8**	78.2	89.3		
HFBI	24.1	**36.6**	**39.3**	**45.6**	**61.7**	**68.1**		
HCCA	**20.9**	**30.4**	**38.8**	**49.7**	**52.6**	**58.3**		
TVCE	**20.2**	**31.1**	**37.9**	**51.8**	73.6	84.2		
AEMI	**22.6**	**31.4**	**40.9**	65.6	108.5	133.9		
Forecasts	12	10	8	6	4	2		

Table 4a. NHC official (OFCL) and climatology-persistence skill baseline (OCD5) intensity forecast errors (kt) for Tropical Storm Juliette, 24–28 August 2025. Mean errors for the previous 5-yr period are shown for comparison. Official errors that are smaller than the 5-yr means are shown in boldface type.

	Forecast Period (h)							
	12	24	36	48	60	72	96	120
OFCL	**4.2**	10.0	**9.4**	**8.3**	**2.5**	**0.0**		
OCD5	4.6	11.9	14.2	11.0	11.2	16.0		
Forecasts	12	10	8	6	4	2	0	0
OFCL (2020-24)	5.7	8.9	10.8	12.9	14.4	15.5	17.0	18.6
OCD5 (2020-24)	7.4	12.4	16.2	19.2	21.0	21.3	21.5	21.6

Table 4b. Homogeneous comparison of selected intensity forecast guidance models (in kt) for Tropical Storm Juliette, 24–28 August 2025. Errors smaller than the NHC official forecast are shown in boldface type.

Model ID	Forecast Period (h)							
	12	24	36	48	60	72	96	120
OFCL	4.2	10.0	9.4	8.3	2.5	0.0		
OCD5	4.6	11.9	14.2	11.0	11.2	16.0		
HWFI	7.6	10.8	12.4	**7.7**	2.5	9.0		
HMNI	4.6	10.9	12.4	**6.2**	5.5	8.0		
HFAI	5.3	10.7	10.5	**7.7**	8.2	8.5		
HFBI	**3.6**	10.2	10.9	**7.7**	5.8	2.5		
DSHP	4.6	10.4	12.1	10.2	9.0	12.0		
LGEM	4.6	10.4	12.5	11.3	7.2	9.0		
HCCA	**3.8**	**9.7**	11.1	8.3	4.0	1.0		
IVCN	**4.0**	**9.5**	10.9	**7.7**	2.5	2.5		
GFSI	6.2	12.2	14.0	10.7	7.8	6.5		
EMXI	7.2	13.5	14.6	12.5	11.8	15.5		
Forecasts	12	10	8	6	4	2		




Figure 1. Best track positions for Tropical Storm Juliette, 24–28 August 2025.




Figure 2. Selected wind observations and best track maximum sustained surface wind speed curve for Tropical Storm Juliette, 24–28 August 2025. Advanced Dvorak Technique estimates represent the Current Intensity at the nominal observation time. SATCON intensity estimates are from the Cooperative Institute for Meteorological Satellite Studies. Dashed vertical lines correspond to 0000 UTC.




Figure 3. Selected pressure observations and best track minimum central pressure curve for Tropical Storm Juliette, 24–28 August 2025. Advanced Dvorak Technique estimates represent the Current Intensity at the nominal observation time. SATCON intensity estimates are from the Cooperative Institute for Meteorological Satellite Studies. KZC P-W refers to pressure estimates derived using the Knaff-Zehr-Courtney pressure-wind relationship. Dashed vertical lines correspond to 0000 UTC.




Figure 4. Composites of 7-day tropical cyclone genesis areas depicted in NHC's Tropical Weather Outlooks prior to the formation of Juliette for (a) all probabilistic genesis categories, (b) the low (<40%) category, (c) medium (40–60%) category, and (d) high (>60%) category. The location of genesis is indicated by the black star.