

machine learning ~~12~~ formulae and expressions

linear regression

hypothesis:

$$h_{\theta}(x) = \theta_0 + \theta_1 x + \dots + \theta_n$$

parameters:

$$\theta_0, \theta_1, \dots, \theta_n$$

cost function:

$$J(\theta_0, \theta_1, \dots, \theta_n) = \frac{1}{2m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})^2$$

goal:

$$\min_{\theta_0, \theta_1, \dots, \theta_n} J(\theta_0, \theta_1, \dots, \theta_n)$$

partial derivative gradient descent for multivariate linear regression:

repeat until convergence {

$$\theta_j := \theta_j - a \frac{\partial}{\partial \theta_j} J(\theta)$$

$$\theta_j := \theta_j - a \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)}) x_j^{(i)}$$

(update θ_j for $j = 0, \dots, n$ simultaneously)

}

normal equation:

$$\theta = (X^T X)^{-1} X^T y$$

regularized multivariate linear regression:

$$J(\theta) = \frac{1}{2m} \left[\sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)})^2 + \lambda \sum_{j=1}^n \theta_j^2 \right]$$

$$\min_{\theta} J(\theta)$$

regularized gradient descent for multivariate linear regression: (for all j)

repeat until convergence {

$$\theta_0 := \theta_0 - a \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)}) x_0^{(i)}$$

$$\theta_j := \theta_j - a \left[\frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)}) x_j^{(i)} + \frac{\lambda}{m} \theta_j \right]$$

(update θ_j for $j = \textcolor{red}{X}$ 1, 2, 3 ..., n simultaneously)

}

alternative notation for θ_j update: $\theta_j := \theta_j \left(1 - a \frac{\lambda}{m} \right) - a \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)}) x_j^{(i)}$

regularized normal equation:

$$\theta = \left(X^T X + \lambda \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & \ddots & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \right)^{-1} X^T y$$

logistic regression

hypothesis:

$$h_{\theta}(x) = g(\theta^T X) \rightarrow g(z) = \frac{1}{1 + e^{-z}}$$

$$h_{\theta}(x) = \frac{1}{1 + e^{-\theta^T X}} \rightarrow \text{sigmoid/logistic function}$$

interpretation:

$$h_{\theta}(x) = P(y = 1|x; \theta)$$

cost function:

$$\text{cost}(h_{\theta}(x^{(i)}), y^{(i)}) = \frac{1}{2} (h_{\theta}(x^{(i)}) - y^{(i)})^2$$

$$\text{cost}(h_{\theta}(x^{(i)}), y^{(i)}) = \begin{cases} -\log(h_{\theta}(x^{(i)})) & \text{if } y = 1 \\ -\log(1 - h_{\theta}(x^{(i)})) & \text{if } y = 0 \end{cases}$$

$$\begin{aligned} J(\theta) &= \frac{1}{m} \sum_{i=1}^m \text{cost}(h_{\theta}(x^{(i)}), y^{(i)}) \\ &= \frac{1}{m} \left[\sum_{i=1}^m y^{(i)} \log h_{\theta}(x^{(i)}) + (1 - y^{(i)}) \log (1 - h_{\theta}(x^{(i)})) \right] \end{aligned}$$

partial derivative gradient descent for multivariate logistic regression:

repeat until convergence {

$$\theta_j := \theta_j - a \frac{\partial}{\partial \theta_j} J(\theta)$$

$$\theta_j := \theta_j - a \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)}) x_j^{(i)}$$

(update θ_j for $j = 0, \dots, n$ simultaneously)

}

$$h_{\theta}(x) = \theta^T X \text{ to } h_{\theta}(x) = \frac{1}{1 + e^{-\theta^T X}}$$

regularized multivariate logistic regression:

$$J(\theta) = \frac{1}{m} \left[\sum_{i=1}^m y^{(i)} \log h_{\theta}(x^{(i)}) + (1 - y^{(i)}) \log (1 - h_{\theta}(x^{(i)})) \right] + \frac{\lambda}{2m} \sum_{j=1}^u \theta_j^2$$

regularized gradient descent for multivariate logistic regression: (for all j)

repeat until convergence {

$$\theta_0 := \theta_0 - a \frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)}) x_0^{(i)}$$

$$\theta_j := \theta_j - a \left[\frac{1}{m} \sum_{i=1}^m (h_{\theta}(x^{(i)}) - y^{(i)}) x_j^{(i)} + \frac{\lambda}{m} \theta_j \right]$$

(update θ_j for $j = \text{X}, 1, 2, 3 \dots, n$ simultaneously)

}

$$h_{\theta}(x) = \theta^T X \text{ to } h_{\theta}(x) = \frac{1}{1+e^{-\theta^T x}}$$

neural networks

backpropagation algorithm

gradient computation

$$J(\theta) = -\frac{1}{m} \left[\sum_{i=1}^m \sum_{k=1}^K y_k^{(i)} \log(h_{\theta}(x^{(i)}))_k + (1 - y_k^{(i)}) \log(1 - (h_{\theta}(x^{(i)}))_k) \right] \\ + \frac{\lambda}{2m} \sum_{l=1}^{L-1} \sum_{i=1}^{s_l} \sum_{j=1}^{s_{l+1}} (\theta_{ji}^{(l)})^2$$

$$\min_{\theta} J(\theta)$$

gradient checking

$$\frac{\partial}{\partial \theta} J(\theta) \approx \frac{J(\theta + \varepsilon) - J(\theta - \varepsilon)}{2\varepsilon}$$

train · validation · test error

training error:

$$J_{train}(\theta) = \frac{1}{2m} \sum_{i=1}^m (h_{\theta}(x^{(i)}), y^{(i)})^2$$

cross validation error:

$$J_{cv}(\theta) = \frac{1}{2m_{cv}} \sum_{i=1}^{m_{cv}} (h_{\theta}(x_{cv}^{(i)}), y_{cv}^{(i)})^2$$

test error:

$$J_{test}(\theta) = \frac{1}{2m_{test}} \sum_{i=1}^{m_{test}} (h_{\theta}(x_{test}^{(i)}), y_{test}^{(i)})^2$$

precision:

$$\frac{\text{true positives}}{\# \text{ **predicted** positives}} = \frac{\text{true positives}}{\text{true positives} + \text{**false positives**}}$$

recall:

$$\frac{\text{true positives}}{\# \text{ **actual** positives}} = \frac{\text{true positives}}{\text{true positives} + \text{**false negatives**}}$$

support vector machines

optimization objective:

$$\min_{\theta} \frac{1}{m} C \sum_{i=1}^m y^{(i)} \text{cost}_1(\theta^T x^{(i)}) + (1 - y^{(i)}) \text{cost}_0(\theta^T x^{(i)}) + \frac{1}{2} \sum_{j=1}^n \theta_j^2$$

hypothesis output:

$$h_{\theta}x = \begin{cases} 1 & \text{if } \theta^T x \text{ is } \geq 0 \\ 0 & \text{otherwise} \end{cases}$$

decision boundary:

$$\min_{\theta} \frac{1}{2} \sum_{j=1}^n \theta_j^2 = \frac{1}{2} (\theta_1^2 + \theta_2^2) = \frac{1}{2} \left(\sqrt{\theta_1^2 + \theta_2^2} \right)^2 = \frac{1}{2} \|\theta\|^2$$

kernels

similarity:

$$f_1 = \text{similarity}(x, \ell^{(1)}) = \exp\left(-\frac{\|x - \ell^{(1)}\|^2}{2\sigma^2}\right) = \exp\left(-\frac{\sum_{j=1}^n (x_j - \ell_j^{(1)})^2}{2\sigma^2}\right)$$

k-means algorithm

optimization objective:

$$J(c^{(1)}, \dots, c^{(m)}, \mu_1, \dots, \mu_K) = \frac{1}{m} \sum_{i=1}^m \|x^{(i)} - \mu_c^{(i)}\|^2$$

anomaly detection algorithm

gaussian distribution:

$$p(x; \mu, \sigma^2) = \frac{1}{\sqrt{2\pi}\sigma} \exp\left(-\frac{(x - \mu)^2}{2\sigma^2}\right)$$

$$p(x) = \prod_{j=1}^n p(x_j; \mu_j, \sigma_j^2) = \prod_{j=1}^n \frac{1}{\sqrt{2\pi}\sigma_j} \exp\left(-\frac{(x_j - \mu_j)^2}{2\sigma_j^2}\right)$$

multivariate gaussian distribution:

$$p(x; \mu, \Sigma) = \frac{1}{(2\pi)^{\frac{n}{2}} |\Sigma|^{\frac{1}{2}}} \exp\left(-\frac{1}{2} (x - \mu)^T \Sigma^{-1} (x - \mu)\right)$$

recommender systems

optimization objective:

to learn $\theta^{(j)}$ (parameter for user j):

$$\min_{\theta^{(j)}} \frac{1}{2} \sum_{i:r(i,j)=1} \left((\theta^{(j)})^T (x^{(i)}) - y^{(i,j)} \right)^2 + \frac{\lambda}{2} \sum_{k=1}^n (\theta_k^{(j)})^2$$

to learn $\theta^{(1)}, \theta^{(2)}, \dots, \theta^{(n_u)}$:

$$\min_{\theta^{(j)}, \dots, \theta^{(n_u)}} \frac{1}{2} \sum_{j=1}^{n_u} \sum_{i:r(i,j)=1} \left((\theta^{(j)})^T (x^{(i)}) - y^{(i,j)} \right)^2 + \frac{\lambda}{2} \sum_{j=1}^{n_u} \sum_{k=1}^n (\theta_k^{(j)})^2$$

simultaneous gradient descent update:

(for $k = 0$)

$$\theta_k^{(j)} := \theta_k^{(j)} - \alpha \sum_{i:r(i,j)=1} \left((\theta^{(j)})^T (x^{(i)}) - y^{(i,j)} \right) x_k^{(i)}$$

(for $k \neq 0$)

$$\theta_k^{(j)} := \theta_k^{(j)} - \alpha \left(\sum_{i:r(i,j)=1} \left((\theta^{(j)})^T (x^{(i)}) - y^{(i,j)} \right) x_k^{(i)} + \lambda \theta_k^{(j)} \right)$$

collaborative filtering optimization algorithm:

given $\theta^{(1)}, \dots, \theta^{(n_u)}$, to learn $x^{(i)}$:

$$\min_{x^{(i)}} \frac{1}{2} \sum_{j:r(i,j)=1} \left((\theta^{(j)})^T (x^{(i)}) - y^{(i,j)} \right)^2 + \frac{\lambda}{2} \sum_{k=1}^n \left(x_k^{(i)} \right)^2$$

given $\theta^{(1)}, \dots, \theta^{(n_u)}$, to learn $x^{(1)}, \dots, x^{(n_m)}$:

$$\min_{x^{(1)}, \dots, x^{(n_m)}} \frac{1}{2} \sum_{i=1}^{n_m} \sum_{j:r(i,j)=1} \left((\theta^{(j)})^T (x^{(i)}) - y^{(i,j)} \right)^2 + \frac{\lambda}{2} \sum_{i=1}^{n_m} \sum_{k=1}^n \left(x_k^{(i)} \right)^2$$

collaborative filtering algorithm:

given features $x^{(1)}, \dots, x^{(n_m)}$, estimate parameters $\theta^{(1)}, \dots, \theta^{(n_u)}$:

$$\min_{\theta^{(1)}, \dots, \theta^{(n_u)}} \frac{1}{2} \sum_{j=1}^{n_u} \sum_{i:r(i,j)=1} \left((\theta^{(j)})^T x^{(i)} - y^{(i,j)} \right)^2 + \frac{\lambda}{2} \sum_{j=1}^{n_u} \sum_{k=1}^n \left(\theta_k^{(j)} \right)^2$$

given parameters $\theta^{(1)}, \dots, \theta^{(n_u)}$, estimate features $x^{(1)}, \dots, x^{(n_m)}$:

$$\min_{x^{(1)}, \dots, x^{(n_m)}} \frac{1}{2} \sum_{i=1}^{n_m} \sum_{j:r(i,j)=1} \left((\theta^{(j)})^T x^{(i)} - y^{(i,j)} \right)^2 + \frac{\lambda}{2} \sum_{i=1}^{n_m} \sum_{k=1}^n \left(x_k^{(i)} \right)^2$$

minimizing $x^{(1)}, \dots, x^{(n_m)}$ and $\theta^{(1)}, \dots, \theta^{(n_u)}$ simultaneously:

$$J(x^{(1)}, \dots, x^{(n_m)}, \theta^{(1)}, \dots, \theta^{(n_u)})$$

$$= \frac{1}{2} \sum_{(i,j):r(i,j)=1} \left((\theta^{(j)})^T x^{(i)} - y^{(i,j)} \right)^2 + \frac{\lambda}{2} \sum_{i=1}^{n_m} \sum_{k=1}^n \left(x_k^{(i)} \right)^2 + \frac{\lambda}{2} \sum_{j=1}^{n_u} \sum_{k=1}^n \left(\theta_k^{(j)} \right)^2$$

collaborative filtering algorithm update:

$$x_k^{(i)} := x_k^{(i)} - \alpha \left(\sum_{j:r(i,j)=1} \left((\theta^{(j)})^T x^{(i)} - y^{(i,j)} \right) \theta_k^{(j)} + \lambda x_k^{(i)} \right)$$

$$\theta_k^{(j)} := \theta_k^{(j)} - \alpha \left(\sum_{i:r(i,j)=1} \left((\theta^{(j)})^T x^{(i)} - y^{(i,j)} \right) x_k^{(i)} + \lambda \theta_k^{(j)} \right)$$