

MELANOBS: forest cover change, carbon, and biodiversity data in Melanesia



Ghislain VIEILLEDENT¹ Philippe BIRNBAUM¹
David BRUY² Thomas IBANEZ²

[1] Cirad UMR AMAP, [2] IRD UMR AMAP



Outline

1 Introduction

- Context
- Objectives

2 Forest cover maps

- FAO FRA estimates
- Global maps
- National data

3 Carbon maps

- Global maps
- National maps

4 Biodiversity maps

- Global biodiversity maps
- Global biodiversity data-sets
- National data-sets

5 Conclusion

- Summary
- Perspectives

Plan

1 Introduction

- Context
- Objectives

2 Forest cover maps

- FAO FRA estimates
- Global maps
- National data

3 Carbon maps

- Global maps
- National maps


4 Biodiversity maps

- Global biodiversity maps
- Global biodiversity data-sets
- National data-sets

5 Conclusion


- Summary
- Perspectives

Context



- Tropical forests : 50% of terrestrial biodiversity.
- Tropical deforestation : 15% of anthropogenic carbon emissions.
- Mapping forest cover, carbon stocks and biodiversity is essential for conservation planning.

Objectives



- MELANOBS project : building a Melanesian forest observatory.
- Which data on forest cover, carbon stock and biodiversity are available for Melanesian countries ?

Plan

1 Introduction

- Context
- Objectives

2 Forest cover maps

- FAO FRA estimates
- Global maps
- National data

3 Carbon maps

- Global maps
- National maps

4 Biodiversity maps

- Global biodiversity maps
- Global biodiversity data-sets
- National data-sets

5 Conclusion

- Summary
- Perspectives

FAO FRA estimates, forest cover

Table – Forest cover estimates (in % of land area).

Country	Area (km ²)	FRA 2015
PNG	462840	78%
Solomon Islands	28896	87%
Vanuatu	12189	36%
Fiji	18272	61%
New Caledonia	18575	45%

- Forest Ressources Assessment (FRA) from the Food and Agriculture Organization (FAO).
- Estimates are reported by countries to FAO.
- Differentiate forest types : forest, primary forest, plantations.
- Not frequently updated.
- Information is not spatialized.

FAO FRA estimates, deforestation

Table – Mean annual deforestation (in ha).

Country	FRA 2015–2020
PNG	34000
Solomon Islands	
Vanuatu	
Fiji	
New Caledonia	

- Rather good estimates of forest cover but poor estimates of deforestation/regrowth.

Global Forest Change (GFC)



- **Global Forest Change** (Hansen et al. 2013, Univ. of Maryland).
- Used by **Global Forest Watch** (GFW) : platform about the world forests. GFW releases the **Global Forest Review**.
- It is in fact a tree cover change product. User must define a tree cover threshold to define the forest (e.g. 30%).
- Derive from Landsat images from 2000. 30m resolution. One mosaic per year.

Global Forest Change (GFC)

Table – Forest cover estimates (in % of land area).

Country	Area (km ²)	FRA 2015	GFC30 2020
PNG	462840	78%	84%
Solomon Islands	28896	87%	81%
Vanuatu	12189	36%	81%
Fiji	18272	61%	57%
New Caledonia	18575	45%	62%

Table – Mean annual deforestation (in ha).

Country	FRA	GFC30
	2015–2020	2010–2020
PNG	34000	104678
Solomon Islands		13460
Vanuatu		939
Fiji		2663
New Caledonia		1328

- Overestimate forest cover if low tree cover threshold (e.g. 30%).
- Underestimate small-scale deforestation (e.g. logging).

Tropical Moist Forests (TMF)

SCIENCE ADVANCES | RESEARCH ARTICLE

ENVIRONMENTAL STUDIES

Long-term (1990–2019) monitoring of forest cover changes in the humid tropics

C. Vancutsem^{1*}, F. Achard¹, J.-F. Pekel¹, G. Vieilledent^{1,2,3,4}, S. Carboni⁵, D. Simonetti¹, J. Gallego¹, L. E.O. C. Aragão⁶, R. Nasi⁷

- Tropical Moist Forests (Vancutsem et al. 2021, from Joint Research Center).
- Only consider evergreen tropical forests (tropical moist forests, mangroves, evergreen dry tropical forests). Cannot be used to monitor deciduous dry forests.
- Derive from Landsat images from 1990. 30m resolution. Time-series at the pixel scale.

Tropical Moist Forests (TMF)

Table – Forest cover estimates (in % of land area).

Country	Area (km2)	FRA 2015	GFC30 2020	TMF 2020
PNG	462840	78%	84%	85%
Solomon Islands	28896	87%	81%	95%
Vanuatu	12189	36%	81%	95%
Fiji	18272	61%	57%	NA
New Caledonia	18575	45%	62%	46%

Table – Mean annual deforestation (in ha).


Country	FRA 2015–2020	GFC30 2010–2020	TMF 2010–2020
PNG	34000	104678	48691
Solomon Islands		13460	1751
Vanuatu		939	564
Fiji		2663	
New Caledonia		1328	2425

- Fiji is not entirely available (beyond the 180th meridian).
- Overestimate forest cover in some areas (e.g. Vanuatu).

National data

- There is room to improve forest cover change maps at the national scale.
- MELANOBS objectives :
 - Which forest cover change data is available at the national scale ?
 - Derive up to date forest cover change maps for participating countries.

In New-Caledonia



- Coarse vegetation maps from IRD (Jaffre, Morat).
- Forest cover change map for 2000-2010-2020 derived from TMF.
- Natural forest cover map for year ~2020 derived from photo-interpretation of aerial images.

Plan

1 Introduction

- Context
- Objectives

2 Forest cover maps

- FAO FRA estimates
- Global maps
- National data

3 Carbon maps

- Global maps
- National maps

4 Biodiversity maps

- Global biodiversity maps
- Global biodiversity data-sets
- National data-sets

5 Conclusion

- Summary
- Perspectives

Global maps

Name	Resolution	Reference	Epoch	Method
Saatchi	1 km	Saatchi 2011	2000	GLAS, MODIS, QSCAT, SRTM
WHRC-Baccini	500 m	Baccini 2012	2008	GLAS, MODIS, SRTM
Avitabile	1 km	Avitabile 2016	2008	fusion of Saatchi and Baccini
GFW-Baccini	30 m	Baccini 2017	2000	GLAS, Landsat, SRTM
CCI Biomass	100 m	Santoro 2019	2020	ALOS2, PALSAR 2, Sentinel 1
GEDI*	1 km	Dubayah 2023	2020	LiDAR GEDI 2, ALS
more...				

- Usually a three step approach : field data, LiDAR, satellite images.
- Satellites images (optical or radar) are used for extrapolation.
- * different approach for GEDI products (see below).

Disadvantages of global products





Figure – Field plots used in Saatchi et al. 2011

- Some countries might be absent from the final map (eg. New Caledonia for Saatchi, WHRC-Baccini and Avitabile's maps).
- Global models might not be accurate for countries with no field data for calibration.
- High discrepancies between maps.
- Resolutions might be coarse : ≥ 500 m.

GEDI derived AGB map




- No extrapolation using satellite images and SRTM.
- GEDI footprints are aggregated within 1 km grid cells.
- Low resolution : 1 km, location uncertainty of about 25 m.
- Same problem as for other data-sets : no field data from Melanesia for calibration.

National maps

- There is room to improve forest carbon stock maps at the national scale.
- MELANOBS objectives :
 - Which forest carbon data is available at the national scale ?
 - Derive up to date forest carbon stock maps for participating countries.

In New-Caledonia



- No existing national forest carbon stock map.
- NC-PIPPN forest inventory network + MELANOBS network of 1ha permanent forest plots.
- LiDAR data.
- Good cover by GEDI.

Plan

1 Introduction

- Context
- Objectives

2 Forest cover maps

- FAO FRA estimates
- Global maps
- National data

3 Carbon maps

- Global maps
- National maps

4 Biodiversity maps

- Global biodiversity maps
- Global biodiversity data-sets
- National data-sets

5 Conclusion

- Summary
- Perspectives

Global biodiversity maps



- As a first approximation of biodiversity in forests, we can focus on tree diversity.
- One objective would be to obtain tree community maps (β diversity).
- More detailed forest typology than dichotomous dry/moist forests or low-elevation/high-elevation forests.
- Global maps often represent species richness (α -diversity). A few examples below.

Global biodiversity maps



RESEARCH ARTICLE

ENVIRONMENTAL SCIENCES
SUSTAINABILITY SCIENCE



High exposure of global tree diversity to human pressure

Wen-Yong Guo (郭文永)^{a,b,c,d,1}, Josep M. Serra-Diaz^e, Franziska Schrotter^f, Wolf L. Eiserhardt^b, Brian S. Maitner^g, Cory Merow^{h,i}, Cyrille Violette, Madhur Anandⁱ, Michaël Belluard^j, Hans Henrik Bruun^m, Chae-ho Byun^k, Jane A. Catford^o, Bruno E. L. Cerabolini^p, Eduardo Chacón-Madrigal^q, Daniela Ciccarelli^r, J. Hans C. Cornelissen^s, Anh Tuan Dang-Le^{t,u}, Angel de Frutos^v, Arildo S. Dias^w, Aelton B. Giroldo^x, Kun Guo^{c,y}, Alvaro G. Gutiérrez^{y,z}, Wesley Hattingh^{a,b}, Tianhua He (何田华)^{b,c,c}, Peter Hietz^{dd}, Nate Hough-Snee^{ee}, Steven Jansen^{ff}, Jens Katte^{v,g,h}, Tamir Klein^{hh}, Benjamin Komacⁱ, Nathan J. B. Kraft^{ll}, Koen Kramer^{kk,l}, Sandra Lavorel^{mm}, Christopher H. Luskⁿⁿ, Adam R. Martin^{oo}, Maurizio Mencuccini^{pp,qq}, Sean T. Michaletz^{rr,ss}, Vanessa Minden^{tt,uu}, Akira S. Mori^{vv}, Ülo Niinemets^{ww}, Yusuke Onoda^{aa}, Josep Peñuelas^{yy,zz}, Valério D. Pillar^{aa,bb}, Jan Pisek^{bb,cc}, Bjorn J. M. Robroek^{cc,cc}, Brandon Schamp^{dd,dd}, Martijn Slottee^{cc}, Énio Egon Sosinski Jr.^{ff}, Nadejda A. Soudzilovskaia^{gg}, Nelson Thiffault^{hh}, Peter van Bodegomⁱⁱ, Fons van der Plasⁱⁱ, Ian J. Wright^{kk,ll}, Wu-Bing Xu^{a,b,v}, Jingming Zheng^{mm,nn}, Brian J. Enquist^{gg,nnn}, and Jens-Christian Svenning^{a,b,1}

Edited by Hugh Possingham, The Nature Conservancy, Sherwood, QLD, Australia; received December 29, 2020; accepted April 13, 2022


- SDM for 46,752 tree species from GBIF, BIEN, DRYFLOR, RAINBIO, and ALA datasets.
- They considered taxonomic, phylogenetic, and functional diversity but disregarded β -diversity.

Global biodiversity maps




Keil & Chase. (2019). Global patterns and drivers of tree diversity integrated across a continuum of spatial grains. *Nature Ecology & Evolution*.

Continental biodiversity maps



Réjou-Méchain et al. (2021). Unveiling African rainforest composition and vulnerability to global change. *Nature*.

Continental biodiversity maps



ter Steege et al. (2003). A spatial model of tree α -diversity and tree density for the Amazon. *Biodiversity and Conservation*.

Global or continental tree data-sets

- GBIF : Global Biodiversity Information Facility.
- BIEN : Botanical Information and Ecology Network.
- DRYFLOR : Latin American Seasonally Dry Tropical Forest Floristic Network.
- RAINBIO : mega-database of tropical African vascular plants distributions.

National data-sets


- No global or continental tree community maps that could be used at national scales.
- But there are global datasets (GBIF, BIEN) that could be used at national scales.
- MELANOBS objectives :
 - Which tree diversity data is available for each country ?
 - Derive first tree community maps for participating countries.

Tree data in New-Caledonia



- NOU herbarium data.
- NC-PIPPN forest plot network with floristic data.
- Endemia (Red List Authority) data.

Mapping tree communities in New Caledonia



- Use of joint species distribution models for 878 species and 554 sites.
- JSMDs : account for species co-occurrence.
- Predicting species probability of presence for each 1km pixel.
- Clustering species to obtain tree communities.

Plan

- 1 Introduction
 - Context
 - Objectives
 - 2 Forest cover maps
 - FAO FRA estimates
 - Global maps
 - National data
 - 3 Carbon maps
 - Global maps
 - National maps
 - 4 Biodiversity maps
 - Global biodiversity maps
 - Global biodiversity data-sets
 - National data-sets
 - 5 Conclusion
 - Summary
 - Perspectives

Summary

- Melanesia is often absent from global forest cover change, carbon, or biodiversity maps.
- For carbon and biodiversity, global maps are not derived using data from Melanesia (or only a few). They do not have a high accuracy if used at the national scale.
- **Objectives** : deriving accurate maps of forest cover change, carbon stocks, and tree communities based on field data from each participating country.

Perspectives

Several perspectives for forest monitoring and conservation planning :

- How current deforestation impact carbon emissions and biodiversity loss ?
- Identifying areas of high conservation values with regards to carbon and biodiversity.
- Anticipating impact on carbon and biodiversity of various deforestation scenarios.
- Carbon and biodiversity credits associated with avoided deforestation.



... Thank you for attention ...

<https://ecology.ghislainv.fr/presentations>



AMAP
lab



cirad