

# Probability and Wiener process

Aakash Ghosh

19MS129

1<sup>st</sup> November, 2022

We are going to look at:

1. Probability from a measure theoretic viewpoint

We are going to look at:

1. Probability from a measure theoretic viewpoint
2. The Law of Large Numbers (LLN)

We are going to look at:

1. Probability from a measure theoretic viewpoint
2. The Law of Large Numbers (LLN)
3. The Central Limit Theorem (CTL)

We are going to look at:

1. Probability from a measure theoretic viewpoint
2. The Law of Large Numbers (LLN)
3. The Central Limit Theorem (CTL)
4. Weiner Processes

# Probability from a measure theoretic viewpoint



## Terminology

We look at some terminology used by probabilists and their analysis counterparts.

We look at some terminology used by probabilists and their analysis counterparts.

1. A **Sample Space** is a **Measure Space**



## Terminology

We look at some terminology used by probabilists and their analysis counterparts.

1. A Sample Space is a Measure Space
2. A Measurable set is an Event

We look at some terminology used by probabilists and their analysis counterparts.

1. A **Sample Space** is a **Measure Space**
2. A **Measurable set** is an **Event**
3. A **Measurable function** is a **Random Variable**

We look at some terminology used by probabilists and their analysis counterparts.

1. A Sample Space is a Measure Space
2. A Measurable set is an Event
3. A Measurable function is a Random Variable
4. The integral  $\int f d\mu$  is called the Expectation or Mean of Random Variable  $f$ .

We look at some terminology used by probabilists and their analysis counterparts.

1. A Sample Space is a Measure Space
2. A Measurable set is an Event
3. A Measurable function is a Random Variable
4. The integral  $\int f d\mu$  is called the Expectation or Mean of Random Variable  $f$ . We will denote expectations with  $E[\cdot]$

1. The **Variance** of a Random Variable  $X$  is defined as

$$\sigma^2(X) = \inf_{a \in \mathbb{R}} E[(X - a)^2].$$

## Terminology

1. The **Variance** of a Random Variable  $X$  is defined as  $\sigma^2(X) = \inf_{a \in \mathbb{R}} E[(X - a)^2]$ . If  $X$  has the second moment, then  $a = E[X]$ .

## Terminology

1. The **Variance** of a Random Variable  $X$  is defined as  $\sigma^2(X) = \inf_{a \in \mathbb{R}} E[(X - a)^2]$ . If  $X$  has the second moment, then  $a = E[X]$ .
2. The **Standard Deviation** is defined as the  $\sigma(X) = \sqrt{\text{Var}(X)}$ .

## Terminology

1. The **Variance** of a Random Variable  $X$  is defined as  $\sigma^2(X) = \inf_{a \in \mathbb{R}} E[(X - a)^2]$ . If  $X$  has the second moment, then  $a = E[X]$ .
2. The **Standard Deviation** is defined as the  $\sigma(X) = \sqrt{\text{Var}(X)}$ .
3. We define the **distribution** of  $X$  to be  $F(t) = P(X < t)$



## Terminology

1. The **Variance** of a Random Variable  $X$  is defined as  $\sigma^2(X) = \inf_{a \in \mathbb{R}} E[(X - a)^2]$ . If  $X$  has the second moment, then  $a = E[X]$ .
2. The **Standard Deviation** is defined as the  $\sigma(X) = \sqrt{\text{Var}(X)}$ .
3. We define the **distribution** of  $X$  to be  $F(t) = P(X \leq t)$
4. For random variables  $\{X_i\}_{i=1}^n$ , we can define a joint distribution by considering a map from  $\Omega \rightarrow \mathbb{R}^n$  and the measure defined by the distribution function given by  $F(t_1, t_2, \dots) = P(\cap \{X_i \leq t_i\})$

# Independence of events

## Independence of Events

A set of events  $\{E_i\}_{i=1}^n$  is called **A set of Independent Events** if for any subset  $\{E_{i_k}\}$  of those events we have:

$$P(\cap E_{i_k}) = \prod P(E_{i_k})$$

## Independence of Events

A set of events  $\{E_i\}_{i=1}^n$  is called **A set of Independent Events** if for any subset  $\{E_{i_k}\}$  of those events we have:

$$P(\cap E_{i_k}) = \prod P(E_{i_k})$$

We can extend this definition to an infinite collection  $\{E_i\}_{i=1}^\infty$  by requiring  $\{E_i\}_{i=1}^k$  to be an independent set of event for all  $k \in \mathbb{N}$ .

## Extension of independence to sub sigma algebras

Let  $\{\mathcal{A}_i\}_{i=1}^n$  be a collection of sub sigma algebras. We call this set independent if for any choice of  $A_i \in \mathcal{A}_i$  we have  $\{A_i\}_{i=1}^n$  to be an independent set of events.

## Extension of independence to sub sigma algebras

Let  $\{\mathcal{A}_i\}_{i=1}^n$  be a collection of sub sigma algebras. We call this set independent if for any choice of  $A_i \in \mathcal{A}_i$  we have  $\{A_i\}_{i=1}^n$  to be an independent set of events. A thing to note is if we define  $\mathcal{A}_i$  to be the sigma algebra defined by  $E_i$ , then we get the definition given before this.

## Independence of Random variables

We call a set of Random variables  $\{X_i\}_{i=1}^n$  to be an **Independent set of random variables** if for any collection of measurable sets  $B_i \in \mathcal{B}_{\mathbb{R}}$  we have  $\{X_i^{-1}(B_i)\}$  to be a set of independent events.

## Independence of Random variables

We call a set of Random variables  $\{X_i\}_{i=1}^n$  to be an **Independent set of random variables** if for any collection of measurable sets  $B_i \in \mathcal{B}_{\mathbb{R}}$  we have  $\{X_i^{-1}(B_i)\}$  to be a set of independent events.

Such a collection is uniquely characterized by the fact that the joint distribution is the product of the individual distributions, i.e.

$$P_{(X_1, X_2, X_3 \dots)} = \prod P_{(X_i)}$$



## Properties of Independent Random Variables

Independence is a rather strong property, and we get some non-trivial results.

## Properties of Independent Random Variables

Independence is a rather strong property, and we get some non-trivial results. For a collection of random variables

$$\{X_i\}_{i=1}^n$$

1.  $P_{X_1+X_2\ldots} = P_{X_1} * P_{X_2} * \ldots$

## Properties of Independent Random Variables

Independence is a rather strong property, and we get some non-trivial results. For a collection of random variables

$$\{X_i\}_{i=1}^n$$

1.  $P_{X_1+X_2\ldots} = P_{X_1} * P_{X_2} * \ldots$
2.  $E(\prod |X_j|) = \prod E(|X_j|)$

## Properties of Independent Random Variables

Independence is a rather strong property, and we get some non-trivial results. For a collection of random variables

$$\{X_i\}_{i=1}^n$$

1.  $P_{X_1+X_2\ldots} = P_{X_1} * P_{X_2} * \ldots$
2.  $E(\prod |X_j|) = \prod E(|X_j|)$
3.  $\sigma^2(\sum X_i) = \sum \sigma^2(X_i)$

## Properties of Independent Random Variables

Independence is a rather strong property, and we get some non-trivial results. For a collection of random variables

$$\{X_i\}_{i=1}^n$$

1.  $P_{X_1+X_2\ldots} = P_{X_1} * P_{X_2} * \ldots$
2.  $E(\prod |X_j|) = \prod E(|X_j|)$
3.  $\sigma^2(\sum X_i) = \sum \sigma^2(X_i)$
4. In general for a measurable  $f$ , and independent random Variables  $X, Y$ , we have
$$E(X.(f \circ Y)) = E(X)E(f \circ Y)$$

## Properties of Independent Random Variables

Independence is a rather strong property, and we get some non-trivial results. For a collection of random variables

$$\{X_i\}_{i=1}^n$$

1.  $P_{X_1+X_2\ldots} = P_{X_1} * P_{X_2} * \ldots$
2.  $E(\prod |X_j|) = \prod E(|X_j|)$
3.  $\sigma^2(\sum X_i) = \sum \sigma^2(X_i)$
4. In general for a measurable  $f$ , and independent random Variables  $X, Y$ , we have
$$E(X.(f \circ Y)) = E(X)E(f \circ Y)$$

# Law of large numbers(LLN)



## The Idea behind LLN

The idea behind LLN is that when an experiment is run multiple times, the resulting result is close to the mean expectations.



## The Idea behind LLN

The idea behind LLN is that when an experiment is run multiple times, the resulting result is close to the mean expectations.

Another way to think about this is to note that the probability (measure) the set taking values away from the expectation is low which is something we expect: points with higher probability gets highlighted when we perform integrations.

## The Idea behind LLN

The idea behind LLN is that when an experiment is run multiple times, the resulting result is close to the mean expectations.

Another way to think about this is to note that the probability (measure) the set taking values away from the expectation is low which is something we expect: points with higher probability gets highlighted when we perform integrations.

There are some restriction on random variables for which this law can be applied, and we look at three variations.

## The Weak Law of large numbers.

The Weak Law of Large Numbers that if  $\{X_j\}_1^\infty$  is a sequence of independent  $L^2$  random variables with means  $\{\mu_j\}$  and variances  $\{\sigma_j^2\}$ , and if  $n^{-2} \sum_1^n \sigma_j^2 \rightarrow 0$  as  $n \rightarrow \infty$ , then  $n^{-1} \sum_1^n (X_j - \mu_j) \rightarrow 0$  in probability as  $n \rightarrow \infty$ .

## The Weak Law of large numbers.

The Weak Law of Large Numbers that if  $\{X_j\}_1^\infty$  is a sequence of independent  $L^2$  random variables with means  $\{\mu_j\}$  and variances  $\{\sigma_j^2\}$ , and if  $n^{-2} \sum_1^n \sigma_j^2 \rightarrow 0$  as  $n \rightarrow \infty$ , then  $n^{-1} \sum_1^n (X_j - \mu_j) \rightarrow 0$  in probability as  $n \rightarrow \infty$ .

✓This is very easy to prove: Only requires Chebyshev's inequality

## The Weak Law of large numbers.

The Weak Law of Large Numbers that if  $\{X_j\}_1^\infty$  is a sequence of independent  $L^2$  random variables with means  $\{\mu_j\}$  and variances  $\{\sigma_j^2\}$ , and if  $n^{-2} \sum_1^n \sigma_j^2 \rightarrow 0$  as  $n \rightarrow \infty$ , then  $n^{-1} \sum_1^n (X_j - \mu_j) \rightarrow 0$  in probability as  $n \rightarrow \infty$ .

✓ This is very easy to prove: Only requires Chebyshev's inequality

✗ As the name of the theorem suggests, the conditions can be relaxed.

## The strong law of natural numbers

Let  $\{X_j\}_1^\infty$  be a sequence of independent  $L^2$  random variables with means  $\{\mu_j\}$  and variances  $\{\sigma_j^2\}$ . If  $n^{-2} \sum_1^n \sigma_j^2 \rightarrow 0$  as  $n \rightarrow \infty$ , then  $n^{-1} \sum_1^n (X_j - \mu_j) \rightarrow 0$  in probability as  $n \rightarrow \infty$ .

✓Relaxes the bounds on  $\lim_{n \rightarrow \infty} n^{-2} \sum_{i=1}^n \sigma_i^2$

## The strong law of natural numbers

Let  $\{X_j\}_1^\infty$  be a sequence of independent  $L^2$  random variables with means  $\{\mu_j\}$  and variances  $\{\sigma_j^2\}$ . If  $n^{-2} \sum_1^n \sigma_j^2 \rightarrow 0$  as  $n \rightarrow \infty$ , then  $n^{-1} \sum_1^n (X_j - \mu_j) \rightarrow 0$  in probability as  $n \rightarrow \infty$ .

✓Relaxes the bounds on  $\lim_{n \rightarrow \infty} n^{-2} \sum_{i=1}^n \sigma_i^2$

✗Requires some work for a full prove. In particular, the Borel-Cantelli lemma and Kolmogorov's Inequality identity is required.

## The strong law of natural numbers for IIDs

If  $\{X_n\}_1^\infty$  is a sequence of independent identically distributed  $L^1$  random variables with mean  $\mu$ , then  $n^{-1} \sum_1^n X_j \rightarrow \mu$  almost surely as  $n \rightarrow \infty$ .



## The strong law of natural numbers for IIDs

If  $\{X_n\}_1^\infty$  is a sequence of independent identically distributed  $L^1$  random variables with mean  $\mu$ , then  $n^{-1} \sum_1^n X_j \rightarrow \mu$  almost surely as  $n \rightarrow \infty$ .

- For IIDs relaxation we get a relaxation with respect to the presence of the second moment.

# The Central Limit Theorem (CLT)

## Weak/Vague convergence of measure

Let  $\{\mu_n\}_{n=1}^{\infty}$  be a sequence of Borel measures on an LCH  $X$ . We say the sequence converges to  $\mu$  vaguely if for any  $f \in C_c(X)$  we have  $\int f d\mu_n \rightarrow \int f d\mu$ .

## Weak/Vague convergence of measure

Let  $\{\mu_n\}_{n=1}^{\infty}$  be a sequence of Borel measures on an LCH  $X$ . We say the sequence converges to  $\mu$  vaguely if for any  $f \in C_c(X)$  we have  $\int f d\mu_n \rightarrow \int f d\mu$ .

If  $X = \mathbb{R}$ , then by Riesz Representation theorem, we have some function  $f_n$  such that  $\mu_n(E) = \int_E f_n dm$ . Then it is sufficient that  $f_n$  converges to  $f$  on the points of continuity of  $f$ .

## The Central Limit Theorem (CLT)

Let  $\{X_j\}$  be a sequence of independent identically distributed  $L^2$  random variables with mean  $\mu$  and variance  $\sigma^2$ . As  $n \rightarrow \infty$ , the distribution of  $(\sigma\sqrt{n})^{-1} \sum_1^n (X_j - \mu)$  converges vaguely to the standard normal distribution  $\nu_0^1$ , and for all  $a \in \mathbb{R}$ ,

$$\lim_{n \rightarrow \infty} P\left(\frac{1}{\sigma\sqrt{n}} \sum_1^n (X_n - \mu) \leq a\right) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^a e^{-t^2/2} dt$$

# Wiener Processes

## A physical background

Wiener Processes occur naturally in nature in the form of Brownian motion. We start with the following assumptions:

1. For an increasing sequence  $\{t_i\}_{i=1}^{\infty}$ , the set of random variables  $\{X(t_i + 1) - X(t_i)\}_{i=1}^{\infty}$  are independent.

## A physical background

Wiener Processes occur naturally in nature in the form of Brownian motion. We start with the following assumptions:

1. For an increasing sequence  $\{t_i\}_{i=1}^{\infty}$ , the set of random variables  $\{X(t_i + 1) - X(t_i)\}_{i=1}^{\infty}$  are independent.
2. As there are no external factors and the process is homogenous, we can assume  $X(t + r) - X(t)$  depends on  $r$  only (and not on  $t$ ).



## A physical background

Wiener Processes occur naturally in nature in the form of Brownian motion. We start with the following assumptions:

1. For an increasing sequence  $\{t_i\}_{i=1}^{\infty}$ , the set of random variables  $\{X(t_i + 1) - X(t_i)\}_{i=1}^{\infty}$  are independent.
2. As there are no external factors and the process is homogenous, we can assume  $X(t + r) - X(t)$  depends on  $r$  only (and not on  $t$ ).
3. Divide  $n$  in small sections of length  $\delta$ . Then we note  $X(t + n) - X(n) = \sum_{i=1}^{t/\delta} X(t + (k + 1)\delta) - X(t + k\delta)$ . We can therefore consider the distribution to be the limiting sum of an infinite series of itself.

## A physical background

Wiener Processes occur naturally in nature in the form of Brownian motion. We start with the following assumptions:

1. For an increasing sequence  $\{t_i\}_{i=1}^{\infty}$ , the set of random variables  $\{X(t_i + 1) - X(t_i)\}_{i=1}^{\infty}$  are independent.
2. As there are no external factors and the process is homogenous, we can assume  $X(t + r) - X(t)$  depends on  $r$  only (and not on  $t$ ).
3. Divide  $n$  in small sections of length  $\delta$ . Then we note  $X(t + n) - X(n) = \sum_{i=1}^{t/\delta} X(t + (k + 1)\delta) - X(t + k\delta)$ . We can therefore consider the distribution to be the limiting sum of and infinite series of itself. Therefore, it stands to reason that the distribution is a normal distribution.

## A physical background

Wiener Processes occur naturally in nature in the form of Brownian motion. We start with the following assumptions:

1. For an increasing sequence  $\{t_i\}_{i=1}^{\infty}$ , the set of random variables  $\{X(t_i + 1) - X(t_i)\}_{i=1}^{\infty}$  are independent.
2. As there are no external factors and the process is homogenous, we can assume  $X(t + r) - X(t)$  depends on  $r$  only (and not on  $t$ ).
3. Divide  $n$  in small sections of length  $\delta$ . Then we note  $X(t + n) - X(n) = \sum_{i=1}^{t/\delta} X(t + (k + 1)\delta) - X(t + k\delta)$ . We can therefore consider the distribution to be the limiting sum of and infinite series of itself. Therefore, it stands to reason that the distribution is a normal distribution.
4. Moreover, it is easy to see from the properties of IID's that  $\sigma^2(X(t + k) - X(t)) = k\sigma^2(X(t + 1) - X(t))$

## The distribution function

From the above heuristic considerations, we guess that  $X$  has the following distribution :

$$X(t+k) - X(k) \sim \nu_0^{C(k)}$$

*...which indeed supports our assumptions.*