

# PLIP Interaction Report

**Compound:** Lyngbyabellin A | **Receptor:** 1MJE (mouse BRCA2 DBD, chain A) | **Best  $\Delta G$ :** -8.903 kcal/mol  
Source: PLIP (plip-tool.biotec.tu-dresden.de) — Track 1, 1MJE receptor

## Hydrophobic Interactions - - - - -

| Index | Residue | AA  | Distance (Å) | Ligand Atom | Protein Atom |
|-------|---------|-----|--------------|-------------|--------------|
| 1     | 2453A   | PRO | 3.54         | 34          | 604          |
| 2     | 2473A   | VAL | 3.83         | 40          | 761          |
| 3     | 2550A   | TRP | 3.87         | 47          | 1510         |
| 4     | 2551A   | LYS | 3.78         | 47          | 1521         |
| 5     | 2557A   | PHE | 3.34         | 39          | 1576         |

## Halogen Bonds ———

| Index | Residue | AA  | Distance (Å) | Donor Angle (°) | Acceptor Angle (°) | Donor Atom | Acceptor Atom |
|-------|---------|-----|--------------|-----------------|--------------------|------------|---------------|
| 1     | 2610A   | CYS | 3.74         | 162.70          | 118.14             | 45 [Cl]    | 2126 [S3]     |
| 2     | 2669A   | GLY | 3.93         | 149.29          | 112.82             | 45 [Cl]    | 2460 [O2]     |

Note: Only compound with halogen bonds, mediated by its two Cl atoms. CYS2610 and GLY2669 are confirmed P1 pocket residues.