

PLIP Interaction Report

Compound: Scytonemin | **Receptor:** 1MJE (mouse BRCA2 DBD, chain A) | **Best ΔG :** -10.99 kcal/mol
Source: PLIP (plip-tool.biotec.tu-dresden.de) — Track 1, 1MJE receptor

Hydrophobic Interactions - - - - -

Index	Residue	AA	Distance (Å)	Ligand Atom	Protein Atom
1	2444A	LEU	3.18	13	527
2	2550A	TRP	3.77	11	1507
3	2551A	LYS	3.41	24	1520
4	2551A	LYS	3.76	7	1518
5	2554A	ALA	3.47	40	1546
6	2579A	TYR	3.74	33	1803
7	2646A	TRP	3.74	31	2249

Hydrogen Bonds —

Index	Residue	AA	Dist H-A (Å)	Dist D-A (Å)	Donor Angle (°)	Protein donor?	Side chain	Donor Atom	Acceptor Atom
1	2551A	LYS	3.16	4.01	141.75	Yes	Yes	1522 [N3+]	16 [N2]

π -Stacking Interactions - - - - -

Index	Residue	AA	Distance (Å)	Angle (°)	Offset (Å)	Stacking Type	Ligand Atoms
1	2550A	TRP	3.89	5.00	1.35	Parallel (P)	37, 38, 39, 40, 41, 42
2	2550A	TRP	3.74	5.19	1.09	Parallel (P)	37, 38, 39, 40, 41, 42

π -Cation Interactions - - - - -

Index	Residue	AA	Distance (Å)	Offset (Å)	Protein charged?	Ligand Group	Ligand Atoms
1	2551A	LYS	3.65	0.65	Yes	Aromatic	16, 18, 19, 20, 23
2	2551A	LYS	3.86	1.40	Yes	Aromatic	17, 18, 19, 21, 22
3	2671A	LYS	3.59	0.47	Yes	Aromatic	6, 9, 10, 12, 13, 14

Note: Richest interaction profile of all four compounds. TRP2550 and LYS2551 are core anchor residues contacted by all four compounds.