HANOI UNIVERSITY OF SCIENCE AND TECHNOLOGY ADVANCED PROGRAMS - PHYSICS HOMEWORK

1. MECHANICS

1.1 PRELIMINARIES

lass:						Stude	ent ID:			• • • • • •			
					\mathbf{G}	rade t	able						
Question:	1	2	3	4	5	6	7	8	9	10	11	12	Total
Points:	4	2	3	3	2	3	4	3	4	5	5	5	43
Score:													
 2 3 4 									• • • • • •				
2. The force charges, between	r is th	e dista	nce be	tween	the cha	arges, a	and $arepsilon$ is						
												• • • • • •	

Н	anoi University of Science and Technology - Advanced Programs - Physics Homework	
3.	Add $(9.2 \times 10^3 \mathrm{s}) + (8.3 \times 10^4 \mathrm{s}) + (0.008 \times 10^6 \mathrm{s})$, taking into account significant figures.	[3]
4.	Multiply $3.079 \times 10^2 \mathrm{m}$ by $0.068 \times 10^{-1} \mathrm{m}$, taking into account significant figures.	[3]
5.	What, approximately, is the percent uncertainty for a measurement given as $1.57\mathrm{m}^2$?	[2
6.	What, roughly, is the percent uncertainty in the volume of a spherical beach ball of radius $r=0.84\pm0.04\mathrm{m}$?	[3]
7.	What is the area, and its approximate uncertainty, of a circle of radius $3.1 \times 10^4 \mathrm{cm}$?	[4]

Page 2 of 4

[Turn over

Н	anoi University of Science and Technology - Advanced Programs - Physics Homework	
8.	Express the following sum with the correct number of significant figures:	[3]
	$1.80\mathrm{m} + 142.5\mathrm{cm} + 5.34 \times 10^5\mathrm{\mu m}$	
9.	A standard baseball has a circumference of approximately $23\mathrm{cm}$. If a baseball had the same mass per unit volume (see Tables in Section $1-5$, Giancoli) as a neutron or a proton, about what would its mass be?	[4]
10.	Grains of fine California beach sand are approximately spheres with an average radius of $50\mathrm{m}$ and are made of silicon dioxide, which has a density of $2600\mathrm{kg}\mathrm{m}^{-3}$. What mass of sand grains would have a total surface area (the total area of all the individual spheres) equal to the surface area of a cube $1.00\mathrm{m}$ on an edge?	[5]

the slope, and 2.0 m deep slips into a valley in a mud slide. Assume that the mud ends up uniformly distributed over a surface area of the valley measuring $0.40\mathrm{km}\times0.40\mathrm{km}$ and that mud has a density of $1900\mathrm{kg}\mathrm{m}^{-3}$. What is the mass of the mud sitting above a $4.0\mathrm{m}^2$ area of the valley floor?	Н	anoi University of Science and Technology - Advanced Programs - Physics Homework	
11. During heavy rain, a section of a mountainside meaesuring 2.5 km horizontally, 0.80 km up along the slope, and 2.0 m deep slips into a valley in a mud slide. Assume that the mud ends up uniformly distributed over a surface area of the valley measuring 0.40 km × 0.40 km and that mud has a density of 1900 kg m ⁻³ . What is the mass of the mud sitting above a 4.0 m ² area of the valley floor? 12. A vertical container with base area measuring 14.0 cm by 17.0 cm is being filled with identical pieces of candy, each with a volume of 50.0 mm ³ and a mass of 0.0200 g. Assume that the volume of the empty spaces between the candies is negligible. If the height of the candies in the container increases at the rate of 0.250 cm s ⁻¹ , at what rate (kilograms per minute) does the mass of the candies in the			
11. During heavy rain, a section of a mountainside meadsuring 2.5 km horizontally, 0.80 km up along the slope, and 2.0 m deep slips into a valley in a mud slide. Assume that the mud ends up uniformly distributed over a surface area of the valley measuring 0.40 km × 0.40 km and that mud has a density of 1900 kg m ⁻³ . What is the mass of the mud sitting above a 4.0 m ² area of the valley floor? 12. A vertical container with base area measuring 14.0 cm by 17.0 cm is being filled with identical pieces of candy, each with a volume of 50.0 mm ³ and a mass of 0.0200 g. Assume that the volume of the empty spaces between the candies is negligible. If the height of the candies in the container increases at the rate of 0.250 cm s ⁻¹ , at what rate (kilograms per minute) does the mass of the candies in the			
the slope, and 2.0 m deep slips into a valley in a mud slide. Assume that the mud ends up uniformly distributed over a surface area of the valley measuring 0.40 km × 0.40 km and that mud has a density of 1900 kg m ⁻³ . What is the mass of the mud sitting above a 4.0 m ² area of the valley floor? 12. A vertical container with base area measuring 14.0 cm by 17.0 cm is being filled with identical pieces of candy, each with a volume of 50.0 mm ³ and a mass of 0.0200 g. Assume that the volume of the empty spaces between the candies is negligible. If the height of the candies in the container increases at the rate of 0.250 cm s ⁻¹ , at what rate (kilograms per minute) does the mass of the candies in the			
the slope, and 2.0 m deep slips into a valley in a mud slide. Assume that the mud ends up uniformly distributed over a surface area of the valley measuring 0.40 km × 0.40 km and that mud has a density of 1900 kg m ⁻³ . What is the mass of the mud sitting above a 4.0 m ² area of the valley floor? 12. A vertical container with base area measuring 14.0 cm by 17.0 cm is being filled with identical pieces of candy, each with a volume of 50.0 mm ³ and a mass of 0.0200 g. Assume that the volume of the empty spaces between the candies is negligible. If the height of the candies in the container increases at the rate of 0.250 cm s ⁻¹ , at what rate (kilograms per minute) does the mass of the candies in the			
12. A vertical container with base area measuring 14.0 cm by 17.0 cm is being filled with identical pieces of candy, each with a volume of 50.0 mm ³ and a mass of 0.0200 g. Assume that the volume of the empty spaces between the candies is negligible. If the height of the candies in the container increases at the rate of 0.250 cm s ⁻¹ , at what rate (kilograms per minute) does the mass of the candies in the	11.	the slope, and $2.0\mathrm{m}$ deep slips into a valley in a mud slide. Assume that the mud ends up uniformly distributed over a surface area of the valley measuring $0.40\mathrm{km} \times 0.40\mathrm{km}$ and that mud has a density	
12. A vertical container with base area measuring 14.0 cm by 17.0 cm is being filled with identical pieces of candy, each with a volume of 50.0 mm ³ and a mass of 0.0200 g. Assume that the volume of the empty spaces between the candies is negligible. If the height of the candies in the container increases at the rate of 0.250 cm s ⁻¹ , at what rate (kilograms per minute) does the mass of the candies in the			
12. A vertical container with base area measuring 14.0 cm by 17.0 cm is being filled with identical pieces of candy, each with a volume of 50.0 mm ³ and a mass of 0.0200 g. Assume that the volume of the empty spaces between the candies is negligible. If the height of the candies in the container increases at the rate of 0.250 cm s ⁻¹ , at what rate (kilograms per minute) does the mass of the candies in the			
12. A vertical container with base area measuring 14.0 cm by 17.0 cm is being filled with identical pieces of candy, each with a volume of 50.0 mm ³ and a mass of 0.0200 g. Assume that the volume of the empty spaces between the candies is negligible. If the height of the candies in the container increases at the rate of 0.250 cm s ⁻¹ , at what rate (kilograms per minute) does the mass of the candies in the			
12. A vertical container with base area measuring 14.0 cm by 17.0 cm is being filled with identical pieces of candy, each with a volume of 50.0 mm ³ and a mass of 0.0200 g. Assume that the volume of the empty spaces between the candies is negligible. If the height of the candies in the container increases at the rate of 0.250 cm s ⁻¹ , at what rate (kilograms per minute) does the mass of the candies in the			
of candy, each with a volume of $50.0 \mathrm{mm^3}$ and a mass of $0.0200 \mathrm{g}$. Assume that the volume of the empty spaces between the candies is negligible. If the height of the candies in the container increases at the rate of $0.250 \mathrm{cm s^{-1}}$, at what rate (kilograms per minute) does the mass of the candies in the			
of candy, each with a volume of $50.0 \mathrm{mm^3}$ and a mass of $0.0200 \mathrm{g}$. Assume that the volume of the empty spaces between the candies is negligible. If the height of the candies in the container increases at the rate of $0.250 \mathrm{cm s^{-1}}$, at what rate (kilograms per minute) does the mass of the candies in the			
of candy, each with a volume of $50.0 \mathrm{mm^3}$ and a mass of $0.0200 \mathrm{g}$. Assume that the volume of the empty spaces between the candies is negligible. If the height of the candies in the container increases at the rate of $0.250 \mathrm{cm s^{-1}}$, at what rate (kilograms per minute) does the mass of the candies in the			
of candy, each with a volume of $50.0 \mathrm{mm^3}$ and a mass of $0.0200 \mathrm{g}$. Assume that the volume of the empty spaces between the candies is negligible. If the height of the candies in the container increases at the rate of $0.250 \mathrm{cm s^{-1}}$, at what rate (kilograms per minute) does the mass of the candies in the			
	12.	of candy, each with a volume of $50.0\mathrm{mm^3}$ and a mass of $0.0200\mathrm{g}$. Assume that the volume of the empty spaces between the candies is negligible. If the height of the candies in the container increases at the rate of $0.250\mathrm{cms^{-1}}$, at what rate (kilograms per minute) does the mass of the candies in the	

Page 4 of 4 End of homework