



Chapter 5. Hypothesis Testing

Nguyễn Văn Hạnh
Department of Applied Mathematics
School of Applied Mathematics and Informatics

First semester, 2023-2024

Contents

- 1 Introduction to hypothesis testing
- 2 Test on a population mean μ .
- 3 Test on a population proportion p .
- 4 Test on the difference between two means $\mu_1 - \mu_2$.
- 5 Test on the difference between two proportions $p_1 - p_2$

Introduction

Example 5.1:

- Past experience indicates that a monthly long-distance telephone bill is normally distributed with $\mu_0 = 17.85$ (USD) and a standard deviation $\sigma = 3.39$.

Introduction

Example 5.1:

- Past experience indicates that a monthly long-distance telephone bill is normally distributed with $\mu_0 = 17.85$ (USD) and a standard deviation $\sigma = 3.39$.
- After an advertising campaign aimed at increasing long-distance telephone usage, the manager of the telephone company took a random sample of 25 household bills and recorded the mean of their monthly usage $\bar{x} = 19.93$.

Introduction

Example 5.1:

- Past experience indicates that a monthly long-distance telephone bill is normally distributed with $\mu_0 = 17.85$ (USD) and a standard deviation $\sigma = 3.39$.
- After an advertising campaign aimed at increasing long-distance telephone usage, the manager of the telephone company took a random sample of 25 household bills and recorded the mean of their monthly usage $\bar{x} = 19.93$.
- Do the data allow us to infer that the campaign was successful?

Introduction

Example 5.1:

- Past experience indicates that a monthly long-distance telephone bill is normally distributed with $\mu_0 = 17.85$ (USD) and a standard deviation $\sigma = 3.39$.
 - After an advertising campaign aimed at increasing long-distance telephone usage, the manager of the telephone company took a random sample of 25 household bills and recorded the mean of their monthly usage $\bar{x} = 19.93$.
 - Do the data allow us to infer that the campaign was successful?
- We need to make a decision of accepting or rejecting the hypothesis that the campaign was successful ($\mu > 17.85$, the mean monthly bill μ after the advertising campaign is greater than 17.85).

Introduction

Example 5.2: It is known that the mean monthly electricity bills of households in a region in March 2022 is 225.5 (thousands dong). Observed the monthly electricity bills of 200 households in April 2022 and the data are given below:


```

196.65 468.75 320.50 300.50 213.05 140.60 290.00 216.95 360.50 317.95 195.55
220.50 255.60 289.00 194.55 374.25 382.05 185.55 219.10 215.60 220.00 186.75
 97.80 340.50  88.50 209.50 234.04 333.00 291.10 108.50 245.00 184.00 153.50
219.50 214.15 155.20 140.40 108.50 410.00 125.50 220.30 160.00 300.50 310.20
244.40 194.50 210.20 360.00 456.50 237.40 235.00 203.25 109.20 240.15 260.50
275.50 101.55 455.50 246.25 291.55 262.00 378.65 194.50 248.00 262.92  85.75
248.00 204.75 310.70 213.10 320.50 125.60 110.25  77.35 119.50 313.50 222.00
388.10 110.50 160.00 210.00 310.30 380.10 281.00 105.35 280.15 188.80 272.50
103.40 213.50 280.50 119.50 166.10 180.50 212.00 154.75 100.50 452.60 436.35
225.00 124.30 170.00 127.35 107.90 140.00 195.00 315.10 241.05 168.00 120.50
223.95 237.05 285.45 100.50 228.55 248.70 175.80 466.05 219.00 216.00 425.50
390.00 176.85 240.50 226.00 108.70 160.00 470.50 225.00 440.00 265.00 162.80
260.50 175.80  73.05 460.50 263.60  59.50 198.00 416.50 315.50 155.00 190.00
158.50 225.00 266.70 153.60 238.00 297.60 201.75 240.50 270.90 196.65 299.20
 70.50 125.60 100.40 240.00 240.00 224.05 194.00 247.00 325.40 102.20 166.10
361.00 430.00 240.00 250.50 470.00 157.75  98.40 236.50 230.85 317.65 200.70
165.00 350.50 319.15 275.88 203.05 234.50 220.75 180.50 436.50 403.00 460.50
220.00 103.50 222.15 170.50 224.15 460.00 260.40 200.50 311.40 260.00 251.55
100.60 212.20

```

Introduction

The distribution of data can be approximated by a normal distribution:



Introduction

- Modelling: We can suppose that the monthly electricity bills X of households in this region in April 2022 follow a normal distribution with parameters $\theta = (\mu, \sigma^2)$.

Introduction

- Modelling: We can suppose that the monthly electricity bills X of households in this region in April 2022 follow a normal distribution with parameters $\theta = (\mu, \sigma^2)$.
- The parameter μ is the population mean (the mean monthly electricity bills of all households in this region in April 2022).

Introduction

- Modelling: We can suppose that the monthly electricity bills X of households in this region in April 2022 follow a normal distribution with parameters $\theta = (\mu, \sigma^2)$.
- The parameter μ is the population mean (the mean monthly electricity bills of all households in this region in April 2022).
- The sample mean $\bar{x} = 236.78 > \mu_0 = 225.5$ (the mean monthly electricity bills of 200 households in April 2022 is greater than that in March 2022).

Introduction

- Modelling: We can suppose that the monthly electricity bills X of households in this region in April 2022 follow a normal distribution with parameters $\theta = (\mu, \sigma^2)$.
- The parameter μ is the population mean (the mean monthly electricity bills of all households in this region in April 2022).
- The sample mean $\bar{x} = 236.78 > \mu_0 = 225.5$ (the mean monthly electricity bills of 200 households in April 2022 is greater than that in March 2022).
- Question: Is there enough evidence to infer that the mean monthly electricity bills μ of all households in April 2022 is greater than that in March 2022?

Introduction

- Modelling: We can suppose that the monthly electricity bills X of households in this region in April 2022 follow a normal distribution with parameters $\theta = (\mu, \sigma^2)$.
- The parameter μ is the population mean (the mean monthly electricity bills of all households in this region in April 2022).
- The sample mean $\bar{x} = 236.78 > \mu_0 = 225.5$ (the mean monthly electricity bills of 200 households in April 2022 is greater than that in March 2022).
- Question: Is there enough evidence to infer that the mean monthly electricity bills μ of all households in April 2022 is greater than that in March 2022?
- We need to make a decision of accepting or rejecting the hypothesis " $\mu > 225.5$ ".

Hypothesis testing: Definitions

- **Research hypothesis:** " $\mu > 225.5$ " is called the research hypothesis: need to be accepted or rejected.

Hypothesis testing: Definitions

- **Research hypothesis:** " $\mu > 225.5$ " is called the research hypothesis: need to be accepted or rejected.
- We state an opposite hypothesis with the research hypothesis and is called the **null hypothesis**, labeled by $H_0 : \mu \leq 225.5$ (the mean monthly electricity bills μ of all households in April 2022 is not greater than that in March 2022).

Hypothesis testing: Definitions

- **Research hypothesis:** " $\mu > 225.5$ " is called the research hypothesis: need to be accepted or rejected.
- We state an opposite hypothesis with the research hypothesis and is called the **null hypothesis**, labeled by $H_0 : \mu \leq 225.5$ (the mean monthly electricity bills μ of all households in April 2022 is not greater than that in March 2022).
- To make a decision of accepting the research hypothesis, we test whether the information of the sample is strong enough to reject the null hypothesis H_0 .

Hypothesis testing: Definitions

- **Research hypothesis:** " $\mu > 225.5$ " is called the research hypothesis: need to be accepted or rejected.
- We state an opposite hypothesis with the research hypothesis and is called the **null hypothesis**, labeled by $H_0 : \mu \leq 225.5$ (the mean monthly electricity bills μ of all households in April 2022 is not greater than that in March 2022).
- To make a decision of accepting the research hypothesis, we test whether the information of the sample is strong enough to reject the null hypothesis H_0 .
- **The alternative hypothesis, labeled by H_1 :** is the research hypothesis ($\mu > 225.5$).

Hypothesis testing: Definitions

- **Research hypothesis:** " $\mu > 225.5$ " is called the research hypothesis: need to be accepted or rejected.
- We state an opposite hypothesis with the research hypothesis and is called the **null hypothesis**, labeled by $H_0 : \mu \leq 225.5$ (the mean monthly electricity bills μ of all households in April 2022 is not greater than that in March 2022).
- To make a decision of accepting the research hypothesis, we test whether the information of the sample is strong enough to reject the null hypothesis H_0 .
- **The alternative hypothesis, labeled by H_1 :** is the research hypothesis ($\mu > 225.5$).
- Problem of testing: test the null hypothesis $H_0 : \mu \leq 225.5$ versus the alternative hypothesis $H_1 : \mu > 225.5$.

Hypothesis testing: Definitions

- 04 outcomes of a test:

Possible Hypothesis Test Outcomes		
Decision	Accept H_0	Reject H_0
H_0 is true	Correct Decision (No error)	Type I Error
	Probability = $1 - \alpha$	Probability = α
H_0 is false	Type II Error	Correct Decision (No error)
	Probability = β	Probability = $1 - \beta$

Hypothesis testing: Definitions

- 04 outcomes of a test:

Possible Hypothesis Test Outcomes		
Decision	Accept H_0	Reject H_0
H_0 is true	Correct Decision (No error)	Type I Error
	Probability = $1 - \alpha$	Probability = α
H_0 is false	Type II Error	Correct Decision (No error)
	Probability = β	Probability = $1 - \beta$

- Type I error:** we reject H_0 when it is true (we infer that $\mu > 225.5$ but μ is really ≤ 225.5)

Hypothesis testing: Definitions

- 04 outcomes of a test:

Possible Hypothesis Test Outcomes		
Decision	Accept H_0	Reject H_0
H_0 is true	Correct Decision (No error)	Type I Error
	Probability = $1 - \alpha$	Probability = α
H_0 is false	Type II Error	Correct Decision (No error)
	Probability = β	Probability = $1 - \beta$

- Type I error:** we reject H_0 when it is true (we infer that $\mu > 225.5$ but μ is really ≤ 225.5)
- Type II error:** we accept H_0 when it is false (we infer that $\mu \leq 225.5$ but μ is really > 225.5)

Hypothesis testing: Definitions

- $P(\text{Type I error}) = P(\text{reject } H_0 \mid H_0 \text{ is true}) = \alpha.$

Hypothesis testing: Definitions

- $P(\text{Type I error}) = P(\text{reject } H_0 \mid H_0 \text{ is true}) = \alpha.$
- $P(\text{Type II error}) = P(\text{accept } H_0 \mid H_0 \text{ is false}) = \beta.$

Hypothesis testing: Definitions

- $P(\text{Type I error}) = P(\text{reject } H_0 \mid H_0 \text{ is true}) = \alpha.$
- $P(\text{Type II error}) = P(\text{accept } H_0 \mid H_0 \text{ is false}) = \beta.$
- $\text{Power of test} = 1 - \beta = P(\text{we accept } H_1 \text{ when it is true}).$

Hypothesis testing: Definitions

- $P(\text{Type I error}) = P(\text{reject } H_0 \mid H_0 \text{ is true}) = \alpha.$
- $P(\text{Type II error}) = P(\text{accept } H_0 \mid H_0 \text{ is false}) = \beta.$
- $\text{Power of test} = 1 - \beta = P(\text{we accept } H_1 \text{ when it is true}).$
- If α decreases, then β will increase and vice versa.

Hypothesis testing: Definitions

- $P(\text{Type I error}) = P(\text{reject } H_0 \mid H_0 \text{ is true}) = \alpha$.
- $P(\text{Type II error}) = P(\text{accept } H_0 \mid H_0 \text{ is false}) = \beta$.
- $\text{Power of test} = 1 - \beta = P(\text{we accept } H_1 \text{ when it is true})$.
- If α decreases, then β will increase and vice versa.
- We control the $P(\text{Type I error})$, α , at a given threshold, called the level of significance, and the test is called the test of significance.

Hypothesis testing: Definitions

- $P(\text{Type I error}) = P(\text{reject } H_0 \mid H_0 \text{ is true}) = \alpha$.
- $P(\text{Type II error}) = P(\text{accept } H_0 \mid H_0 \text{ is false}) = \beta$.
- $\text{Power of test} = 1 - \beta = P(\text{we accept } H_1 \text{ when it is true})$.
- If α decreases, then β will increase and vice versa.
- We control the $P(\text{Type I error})$, α , at a given threshold, called the level of significance, and the test is called the test of significance.
- We usually choose the level of significance α that equals to 1%, 5% or 10%.

Procedure of a test

- **Step 1:** State the null hypothesis H_0 and the alternative hypothesis H_1 .

Procedure of a test

- Step 1: State the null hypothesis H_0 and the alternative hypothesis H_1 .
- Step 2: Consider the assumptions of the test.

Procedure of a test

- **Step 1:** State the null hypothesis H_0 and the alternative hypothesis H_1 .
- **Step 2:** Consider the assumptions of the test.
- **Step 3:** Compute the test - statistic Z : is a statistic that has fully deterministic distribution when the null hypothesis is true.

Procedure of a test

- **Step 1:** State the null hypothesis H_0 and the alternative hypothesis H_1 .
- **Step 2:** Consider the assumptions of the test.
- **Step 3:** Compute the test - statistic Z : is a statistic that has fully deterministic distribution when the null hypothesis is true.
- **Step 4:** Find a rejection region W_α such that $P(Z \in W_\alpha | H_0 \text{ is true}) \leq \alpha$.

Procedure of a test

- **Step 1:** State the null hypothesis H_0 and the alternative hypothesis H_1 .
- **Step 2:** Consider the assumptions of the test.
- **Step 3:** Compute the test - statistic Z : is a statistic that has fully deterministic distribution when the null hypothesis is true.
- **Step 4:** Find a rejection region W_α such that $P(Z \in W_\alpha | H_0 \text{ is true}) \leq \alpha$.
- **Step 5:** Make a decision: if the test-statistic $Z \in W_\alpha$ then we reject H_0 , otherwise we fail to reject H_0 . Give a conclusion about the research hypothesis.

Principle of small probability

- **Principle of small probability:** “Specified events of small probability do not occur by chance, when the experiment is performed once.”

Principle of small probability

- **Principle of small probability:** “Specified events of small probability do not occur by chance, when the experiment is performed once.”
- **Decision Rule:** if the test-statistic $Z \in W_\alpha$ then we reject H_0 , otherwise we fail to reject H_0

Principle of small probability

- **Principle of small probability:** "Specified events of small probability do not occur by chance, when the experiment is performed once."
- **Decision Rule:** if the test-statistic $Z \in W_\alpha$ then we reject H_0 , otherwise we fail to reject H_0 since:
 - If the test-statistic $Z \in W_\alpha$ and we accept H_0 , the the event " $Z \in W_\alpha | H_0$ is true " of small probability α has occurred by chance. This contradicts the principle of small probability.

Principle of small probability

- **Principle of small probability:** "Specified events of small probability do not occur by chance, when the experiment is performed once."
- **Decision Rule:** if the test-statistic $Z \in W_\alpha$ then we reject H_0 , otherwise we fail to reject H_0 since:
 - If the test-statistic $Z \in W_\alpha$ and we accept H_0 , the the event " $Z \in W_\alpha | H_0$ is true " of small probability α has occurred by chance. This contradicts the principle of small probability.
 - So if the test-statistic $Z \in W_\alpha$ then we reject H_0 , otherwise we fail to reject H_0 .

Four problems of testing

We study on the following four problems of testing:

- Test on a population mean μ .
- Test on a population proportion p .
- Test on the difference between 2 population means $\mu_1 - \mu_2$.
- Test on the difference between 2 population proportions $p_1 - p_2$.

Test on a population mean μ .

Problem 1: We observe a population X where $\mu = E(X)$ and $\sigma^2 = V(X)$. Consider a random sample (X_1, X_2, \dots, X_n) taken from the population X .

Test on a population mean μ .

Problem 1: We observe a population X where $\mu = E(X)$ and $\sigma^2 = V(X)$. Consider a random sample (X_1, X_2, \dots, X_n) taken from the population X .

- We test a research hypothesis on μ that $\mu > \mu_0$ or $\mu < \mu_0$ or $\mu \neq \mu_0$.

Test on a population mean μ .

Problem 1: We observe a population X where $\mu = E(X)$ and $\sigma^2 = V(X)$. Consider a random sample (X_1, X_2, \dots, X_n) taken from the population X .

- We test a research hypothesis on μ that $\mu > \mu_0$ or $\mu < \mu_0$ or $\mu \neq \mu_0$.
- It can be proved that the null hypotheses $H_0 : \mu = \mu_0$ and $H_0 : \mu \leq \mu_0$ against the alternative hypothesis $H_1 : \mu > \mu_0$ have the same rejection region W_α .

Test on a population mean μ .

Problem 1: We observe a population X where $\mu = E(X)$ and $\sigma^2 = V(X)$. Consider a random sample (X_1, X_2, \dots, X_n) taken from the population X .

- We test a research hypothesis on μ that $\mu > \mu_0$ or $\mu < \mu_0$ or $\mu \neq \mu_0$.
- It can be proved that the null hypotheses $H_0 : \mu = \mu_0$ and $H_0 : \mu \leq \mu_0$ against the alternative hypothesis $H_1 : \mu > \mu_0$ have the same rejection region W_α .
- Thus, we only consider the null hypothesis $H_0 : \mu = \mu_0$ against one of three types of alternative:
 - $H_1 : \mu > \mu_0$: right-tailed test
 - or $H_1 : \mu < \mu_0$: left-tailed test
 - or $H_1 : \mu \neq \mu_0$: two-tailed test.

Test on a population mean μ .

Case 1: The population $X \sim N(\mu; \sigma^2)$ where σ^2 is known.

Test on a population mean μ .

Case 1: The population $X \sim N(\mu; \sigma^2)$ where σ^2 is known.

Case 1.1: We consider the right-tailed test $H_1 : \mu > \mu_0$.

Test on a population mean μ .

Case 1: The population $X \sim N(\mu; \sigma^2)$ where σ^2 is known.

Case 1.1: We consider the right-tailed test $H_1 : \mu > \mu_0$.

- We choose the test statistic

$$Z = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}} \sim N(0, 1) \text{ if } H_0 \text{ is true}$$

Test on a population mean μ .

Case 1: The population $X \sim N(\mu; \sigma^2)$ where σ^2 is known.

Case 1.1: We consider the right-tailed test $H_1 : \mu > \mu_0$.

- We choose the test statistic

$$Z = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}} \sim N(0, 1) \text{ if } H_0 \text{ is true}$$

- Find the rejection region W_α : We reject H_0 and accept H_1 if $\bar{X} >> \mu_0$ or $\bar{X} - \mu_0$ is large enough $\Leftrightarrow Z = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}} > x$, for a critical value x to be found.

Test on a population mean μ .

Case 1: The population $X \sim N(\mu; \sigma^2)$ where σ^2 is known.

Case 1.1: We consider the right-tailed test $H_1 : \mu > \mu_0$.

- We choose the test statistic

$$Z = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}} \sim N(0, 1) \text{ if } H_0 \text{ is true}$$

- Find the rejection region W_α : We reject H_0 and accept H_1 if $\bar{X} >> \mu_0$ or $\bar{X} - \mu_0$ is large enough $\Leftrightarrow Z = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}} > x$, for a critical value x to be found.
- The rejection region $W_\alpha = \{Z > x\} = (x; +\infty)$ satisfies:

$$P(\text{Type I error}) = P(\text{reject } H_0 | \text{if } H_0 \text{ is true}) = P(Z \in W_\alpha | H_0 \text{ is true})$$

$$P(\text{Type I error}) = P(Z > x | H_0 \text{ is true}) \leq \alpha \Leftrightarrow x = Z_\alpha$$

Test on a population mean μ .

Case 1: The population $X \sim N(\mu; \sigma^2)$ where σ^2 is known.

Case 1.1: We consider the right-tailed test $H_1 : \mu > \mu_0$.

- We choose the test statistic

$$Z = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}} \sim N(0, 1) \text{ if } H_0 \text{ is true}$$

- The rejection region $W_\alpha = \{Z > Z_\alpha\} = (Z_\alpha; +\infty)$.

Test on a population mean μ .

Case 1: The population $X \sim N(\mu; \sigma^2)$ where σ^2 is known.

Case 1.1: We consider the right-tailed test $H_1 : \mu > \mu_0$.

- We choose the test statistic

$$Z = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}} \sim N(0, 1) \text{ if } H_0 \text{ is true}$$

- The rejection region $W_\alpha = \{Z > Z_\alpha\} = (Z_\alpha; +\infty)$.
- The decision rule: If $Z > Z_\alpha$, then we reject H_0 , otherwise we fail to reject H_0 .

Test on a population mean μ .

Case 1: The population $X \sim N(\mu; \sigma^2)$ where σ^2 is known.

Case 1.2: We consider the left-tailed test $H_1 : \mu < \mu_0$.

Test on a population mean μ .

Case 1: The population $X \sim N(\mu; \sigma^2)$ where σ^2 is known.

Case 1.2: We consider the left-tailed test $H_1 : \mu < \mu_0$.

- We choose the test statistic

$$Z = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}} \sim N(0, 1) \text{ if } H_0 \text{ is true}$$

Test on a population mean μ .

Case 1: The population $X \sim N(\mu; \sigma^2)$ where σ^2 is known.

Case 1.2: We consider the left-tailed test $H_1 : \mu < \mu_0$.

- We choose the test statistic

$$Z = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}} \sim N(0, 1) \text{ if } H_0 \text{ is true}$$

- Find the rejection region W_α : We reject H_0 and accept H_1 if $\bar{X} << \mu_0$ or $\bar{X} - \mu_0$ is small enough $\Leftrightarrow Z = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}} < x$, for a critical value x to be found.

Test on a population mean μ .

Case 1: The population $X \sim N(\mu; \sigma^2)$ where σ^2 is known.

Case 1.2: We consider the left-tailed test $H_1 : \mu < \mu_0$.

- We choose the test statistic

$$Z = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}} \sim N(0, 1) \text{ if } H_0 \text{ is true}$$

- Find the rejection region W_α : We reject H_0 and accept H_1 if $\bar{X} << \mu_0$ or $\bar{X} - \mu_0$ is small enough $\Leftrightarrow Z = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}} < x$, for a critical value x to be found.
- The rejection region $W_\alpha = \{Z < x\} = (-\infty; x)$ satisfies:

$$P(\text{Type I error}) = P(\text{reject } H_0 | \text{if } H_0 \text{ is true}) = P(Z \in W_\alpha | H_0 \text{ is true})$$

$$P(\text{Type I error}) = P(Z < x | H_0 \text{ is true}) \leq \alpha \Leftrightarrow x = -Z_\alpha$$

Test on a population mean μ .

Case 1: The population $X \sim N(\mu; \sigma^2)$ where σ^2 is known.

Case 1.2: We consider the left-tailed test $H_1 : \mu < \mu_0$.

- We choose the test statistic

$$Z = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}} \sim N(0, 1) \text{ if } H_0 \text{ is true}$$

- The rejection region $W_\alpha = \{Z < -Z_\alpha\} = (-\infty; -Z_\alpha)$.

Test on a population mean μ .

Case 1: The population $X \sim N(\mu; \sigma^2)$ where σ^2 is known.

Case 1.2: We consider the left-tailed test $H_1 : \mu < \mu_0$.

- We choose the test statistic

$$Z = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}} \sim N(0, 1) \text{ if } H_0 \text{ is true}$$

- The rejection region $W_\alpha = \{Z < -Z_\alpha\} = (-\infty; -Z_\alpha)$.
- The decision rule: If $Z < -Z_\alpha$, then we reject H_0 , otherwise we fail to reject H_0 .

Test on a population mean μ .

Case 1: The population $X \sim N(\mu; \sigma^2)$ where σ^2 is known.

Case 1.3: We consider the two-tailed test $H_1 : \mu \neq \mu_0$.

Test on a population mean μ .

Case 1: The population $X \sim N(\mu; \sigma^2)$ where σ^2 is known.

Case 1.3: We consider the two-tailed test $H_1 : \mu \neq \mu_0$.

- We choose the test statistic

$$Z = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}} \sim N(0, 1) \text{ if } H_0 \text{ is true}$$

Test on a population mean μ .

Case 1: The population $X \sim N(\mu; \sigma^2)$ where σ^2 is known.

Case 1.3: We consider the two-tailed test $H_1 : \mu \neq \mu_0$.

- We choose the test statistic

$$Z = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}} \sim N(0, 1) \text{ if } H_0 \text{ is true}$$

- Find the rejection region W_α : We reject H_0 and accept H_1 if \bar{X} is much different from μ_0 or $|\bar{X} - \mu_0|$ is large enough
 $\Leftrightarrow |Z| = \frac{|\bar{X} - \mu_0|}{\sigma/\sqrt{n}} > x$, for a critical value x to be found.

Test on a population mean μ .

Case 1: The population $X \sim N(\mu; \sigma^2)$ where σ^2 is known.

Case 1.3: We consider the two-tailed test $H_1 : \mu \neq \mu_0$.

- We choose the test statistic

$$Z = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}} \sim N(0, 1) \text{ if } H_0 \text{ is true}$$

- Find the rejection region W_α : We reject H_0 and accept H_1 if \bar{X} is much different from μ_0 or $|\bar{X} - \mu_0|$ is large enough
 $\Leftrightarrow |Z| = \frac{|\bar{X} - \mu_0|}{\sigma/\sqrt{n}} > x$, for a critical value x to be found.
- The rejection region $W_\alpha = \{|Z| > x\} = (-\infty; -x) \cup (x; +\infty)$ satisfies:

$$P(\text{Type I error}) = P(\text{reject } H_0 | \text{if } H_0 \text{ is true}) = P(Z \in W_\alpha | H_0 \text{ is true})$$

$$P(\text{Type I error}) = P(|Z| > x | H_0 \text{ is true}) \leq \alpha \Leftrightarrow x = Z_{\alpha/2}$$

Test on a population mean μ .

Case 1: The population $X \sim N(\mu; \sigma^2)$ where σ^2 is known.

Case 1.3: We consider the two-tailed test $H_1 : \mu \neq \mu_0$.

- We choose the test statistic

$$Z = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}} \sim N(0, 1) \text{ if } H_0 \text{ is true}$$

- The rejection region

$$W_\alpha = \{|Z| > Z_{\alpha/2}\} = (-\infty; -Z_{\alpha/2}) \cup (Z_{\alpha/2}; +\infty).$$

Test on a population mean μ .

Case 1: The population $X \sim N(\mu; \sigma^2)$ where σ^2 is known.

Case 1.3: We consider the two-tailed test $H_1 : \mu \neq \mu_0$.

- We choose the test statistic

$$Z = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}} \sim N(0, 1) \text{ if } H_0 \text{ is true}$$

- The rejection region

$$W_\alpha = \{|Z| > Z_{\alpha/2}\} = (-\infty; -Z_{\alpha/2}) \cup (Z_{\alpha/2}; +\infty).$$

- The decision rule: If $|Z| > Z_{\alpha/2}$, then we reject H_0 , otherwise we fail to reject H_0 .

Test on a population mean μ .

Case 2: The population $X \sim N(\mu; \sigma^2)$ where σ^2 is unknown.

Test on a population mean μ .

Case 2: The population $X \sim N(\mu; \sigma^2)$ where σ^2 is unknown.

- We choose the test statistic

$$T = \frac{\bar{X} - \mu_0}{S/\sqrt{n}} \sim t_{n-1} \text{ if } H_0 \text{ is true}$$

Test on a population mean μ .

Case 2: The population $X \sim N(\mu; \sigma^2)$ where σ^2 is unknown.

- We choose the test statistic

$$T = \frac{\bar{X} - \mu_0}{S/\sqrt{n}} \sim t_{n-1} \text{ if } H_0 \text{ is true}$$

- Similarly, the rejection region is as the following

$$W_\alpha = \begin{cases} (t_{n-1;\alpha}; +\infty), & \text{if } H_1 : \mu > \mu_0 \\ (-\infty; -t_{n-1;\alpha}), & \text{if } H_1 : \mu < \mu_0 \\ (-\infty; -t_{n-1;\alpha/2}) \cup (t_{n-1;\alpha/2}; +\infty), & \text{if } H_1 : \mu \neq \mu_0 \end{cases}$$

Test on a population mean μ .

Case 3: We do not know $X \sim N(\mu; \sigma^2)$ where σ^2 is known and n is large enough ($n > 30$).

Test on a population mean μ .

Case 3: We do not know $X \sim N(\mu; \sigma^2)$ where σ^2 is known and n is large enough ($n > 30$).

- We choose the test statistic

$$Z = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}} \approx N(0; 1) \text{ if } H_0 \text{ is true}$$

Test on a population mean μ .

Case 3: We do not know $X \sim N(\mu; \sigma^2)$ where σ^2 is known and n is large enough ($n > 30$).

- We choose the test statistic

$$Z = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}} \approx N(0; 1) \text{ if } H_0 \text{ is true}$$

- Similarly, the rejection region is as the following

$$W_\alpha = \begin{cases} (Z_\alpha; +\infty), & \text{if } H_1 : \mu > \mu_0 \\ (-\infty; -Z_\alpha), & \text{if } H_1 : \mu < \mu_0 \\ (-\infty; -Z_{\alpha/2}) \cup (Z_{\alpha/2}; +\infty), & \text{if } H_1 : \mu \neq \mu_0 \end{cases}$$

Test on a population mean μ .

Case 4: We do not know $X \sim N(\mu; \sigma^2)$ where σ^2 is unknown and n is large enough ($n > 30$).

Test on a population mean μ .

Case 4: We do not know $X \sim N(\mu; \sigma^2)$ where σ^2 is unknown and n is large enough ($n > 30$).

- We choose the test statistic

$$T = \frac{\bar{X} - \mu_0}{S/\sqrt{n}} \approx N(0; 1) \text{ if } H_0 \text{ is true}$$

Test on a population mean μ .

Case 4: We do not know $X \sim N(\mu; \sigma^2)$ where σ^2 is unknown and n is large enough ($n > 30$).

- We choose the test statistic

$$T = \frac{\bar{X} - \mu_0}{S/\sqrt{n}} \approx N(0; 1) \text{ if } H_0 \text{ is true}$$

- Similarly, the rejection region is as the following

$$W_\alpha = \begin{cases} (Z_\alpha; +\infty), & \text{if } H_1 : \mu > \mu_0 \\ (-\infty; -Z_\alpha), & \text{if } H_1 : \mu < \mu_0 \\ (-\infty; -Z_{\alpha/2}) \cup (Z_{\alpha/2}; +\infty), & \text{if } H_1 : \mu \neq \mu_0 \end{cases}$$

Test on a population mean μ .

Case 4: We do not know $X \sim N(\mu; \sigma^2)$ where σ^2 is unknown and n is large enough ($n > 30$).

- We choose the test statistic

$$T = \frac{\bar{X} - \mu_0}{S/\sqrt{n}} \approx N(0; 1) \text{ if } H_0 \text{ is true}$$

- Similarly, the rejection region is as the following

$$W_\alpha = \begin{cases} (Z_\alpha; +\infty), & \text{if } H_1 : \mu > \mu_0 \\ (-\infty; -Z_\alpha), & \text{if } H_1 : \mu < \mu_0 \\ (-\infty; -Z_{\alpha/2}) \cup (Z_{\alpha/2}; +\infty), & \text{if } H_1 : \mu \neq \mu_0 \end{cases}$$

Case 5: We do not know $X \sim N(\mu; \sigma^2)$ and n is small. We do not study this case (the nonparametric test).

Test on a population mean μ .

Example 5.1: Test the hypothesis that the average content of containers of a particular lubricant differs from 10 liters if the contents of a random sample of 10 containers are 10.2, 9.7, 10.1, 10.3, 10.1, 9.8, 9.9, 10.4, 10.3, and 9.8 liters. Use a 0.05 level of significance and assume that the distribution of contents is normal.

Test on a population mean μ .

Example 5.1: Test the hypothesis that the average content of containers of a particular lubricant differs from 10 liters if the contents of a random sample of 10 containers are 10.2, 9.7, 10.1, 10.3, 10.1, 9.8, 9.9, 10.4, 10.3, and 9.8 liters. Use a 0.05 level of significance and assume that the distribution of contents is normal.

- Let X be the content of containers of lubricant. It is supposed that $X \sim N(\mu; \sigma^2)$ where σ^2 is unknown. We test the null hypothesis $H_0 : \mu = 10$ against $H_1 : \mu \neq 10$ at $\alpha = 0.05$ level of significance.

Test on a population mean μ .

Example 5.1: Test the hypothesis that the average content of containers of a particular lubricant differs from 10 liters if the contents of a random sample of 10 containers are 10.2, 9.7, 10.1, 10.3, 10.1, 9.8, 9.9, 10.4, 10.3, and 9.8 liters. Use a 0.05 level of significance and assume that the distribution of contents is normal.

- Let X be the content of containers of lubricant. It is supposed that $X \sim N(\mu; \sigma^2)$ where σ^2 is unknown. We test the null hypothesis $H_0 : \mu = 10$ against $H_1 : \mu \neq 10$ at $\alpha = 0.05$ level of significance.
- The test statistic is

$$T = \frac{\bar{X} - \mu_0}{S/\sqrt{n}} \sim t_{n-1} \text{ if } H_0 \text{ is true}$$

Test on a population mean μ .

- For the given sample: $n = 10$; $\bar{x} = (10.2 + \dots + 9.8)/10 = 10.06$; $s^2 = (10.2^2 + \dots + 9.8^2 - 10 * 10.06^2)/9 = 0.06$; $s = 0.245$. Then

$$T_{obs} = \frac{10.06 - 10}{0.245/\sqrt{10}} = 0.774$$

Test on a population mean μ .

- For the given sample: $n = 10$; $\bar{x} = (10.2 + \dots + 9.8)/10 = 10.06$; $s^2 = (10.2^2 + \dots + 9.8^2 - 10 * 10.06^2)/9 = 0.06$; $s = 0.245$. Then

$$T_{obs} = \frac{10.06 - 10}{0.245/\sqrt{10}} = 0.774$$

- The rejection region is $W_\alpha = (-\infty; -t_{n-1;\alpha/2}) \cup (t_{n-1;\alpha/2}; +\infty) = (-\infty; -t_{9;0.025}) \cup (t_{9;0.025}; +\infty) = (-\infty; -2.26) \cup (2.26; +\infty)$.

Test on a population mean μ .

- For the given sample: $n = 10$; $\bar{x} = (10.2 + \dots + 9.8)/10 = 10.06$; $s^2 = (10.2^2 + \dots + 9.8^2 - 10 * 10.06^2)/9 = 0.06$; $s = 0.245$. Then

$$T_{obs} = \frac{10.06 - 10}{0.245/\sqrt{10}} = 0.774$$

- The rejection region is $W_\alpha = (-\infty; -t_{n-1;\alpha/2}) \cup (t_{n-1;\alpha/2}; +\infty) = (-\infty; -t_{9;0.025}) \cup (t_{9;0.025}; +\infty) = (-\infty; -2.26) \cup (2.26; +\infty)$.
- Since $T_{obs} \notin W_\alpha$ then we fail to reject H_0 .

Test on a population mean μ .

- For the given sample: $n = 10$; $\bar{x} = (10.2 + \dots + 9.8)/10 = 10.06$; $s^2 = (10.2^2 + \dots + 9.8^2 - 10 * 10.06^2)/9 = 0.06$; $s = 0.245$. Then

$$T_{obs} = \frac{10.06 - 10}{0.245/\sqrt{10}} = 0.774$$

- The rejection region is $W_\alpha = (-\infty; -t_{n-1;\alpha/2}) \cup (t_{n-1;\alpha/2}; +\infty) = (-\infty; -t_{9;0.025}) \cup (t_{9;0.025}; +\infty) = (-\infty; -2.26) \cup (2.26; +\infty)$.
- Since $T_{obs} \notin W_\alpha$ then we fail to reject H_0 . Thus, there is not enough evidence to infer that the average content of containers of a lubricant differs from 10 liters at 5% level of significance.

Introduction

Example 5.2: Past experience indicates that the mean of monthly long-distance telephone bills is $\mu_0 = 17.85$ (USD). After an advertising campaign aimed at increasing long-distance telephone usage, the manager of the telephone company took a random sample of 25 household bills and recorded the mean of their monthly usage $\bar{x} = 19.93$. It is supposed that the monthly long-distance telephone bill follows a normal distribution of a standard deviation $\sigma = 3.39$.

- Do the data allow us to infer that the campaign was successful at 5% level of significance?
- Find the probability of type II error of the test if the true mean of telephone bills after the advertising is 20 (USD).

Test on a population mean μ .

Solution of Example 5.2:

- Let X be the monthly long-distance telephone bill of households after the advertising campaign. It is supposed that $X \sim N(\mu; \sigma^2)$ where $\sigma = 3.39$. We test the null hypothesis $H_0 : \mu = 17.85$ against $H_1 : \mu > 17.85$ at $\alpha = 0.05$ level of significance.

Test on a population mean μ .

Solution of Example 5.2:

- Let X be the monthly long-distance telephone bill of households after the advertising campaign. It is supposed that $X \sim N(\mu; \sigma^2)$ where $\sigma = 3.39$. We test the null hypothesis $H_0 : \mu = 17.85$ against $H_1 : \mu > 17.85$ at $\alpha = 0.05$ level of significance.
- The test statistic is

$$Z = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}} \sim N(0; 1) \text{ if } H_0 \text{ is true}$$

Test on a population mean μ .

Solution of Example 5.2:

- Let X be the monthly long-distance telephone bill of households after the advertising campaign. It is supposed that $X \sim N(\mu; \sigma^2)$ where $\sigma = 3.39$. We test the null hypothesis $H_0 : \mu = 17.85$ against $H_1 : \mu > 17.85$ at $\alpha = 0.05$ level of significance.
- The test statistic is

$$Z = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}} \sim N(0; 1) \text{ if } H_0 \text{ is true}$$

For the given sample, we have $n = 25$; $\bar{x} = 19.93$ then

$$Z_{obs} = \frac{19.93 - 17.85}{3.39/\sqrt{25}} = 3.07$$

Test on a population mean μ .

- The rejection region is

$$W_\alpha = (Z_\alpha; +\infty) = (Z_{0.05}; +\infty) = (1.645; +\infty).$$

Test on a population mean μ .

- The rejection region is
$$W_\alpha = (Z_\alpha; +\infty) = (Z_{0.05}; +\infty) = (1.645; +\infty).$$
- Since $T_{obs} \in W_\alpha$ then we reject H_0 . Thus, there is enough evidence to infer that the campaign was successful at 5% level of significance.

Test on a population mean μ .

- The rejection region is
 $W_\alpha = (Z_\alpha; +\infty) = (Z_{0.05}; +\infty) = (1.645; +\infty)$.
- Since $T_{obs} \in W_\alpha$ then we reject H_0 . Thus, there is enough evidence to infer that the campaign was successful at 5% level of significance.
- The probability of type II error of the test is

$$\begin{aligned}
 \beta &= P(\text{type II error}) = P(\text{accept } H_0 \mid H_0 \text{ is false}) = P(Z \notin W_\alpha \mid \mu = 20) \\
 &= P(Z \leq Z_\alpha \mid \mu = 20) = P\left(\frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}} \leq Z_\alpha \mid \mu = 20\right) \\
 &= P\left(\frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \leq Z_\alpha + \frac{\mu_0 - \mu}{\sigma/\sqrt{n}} \mid \mu = 20\right)
 \end{aligned}$$

Test on a population mean μ .

- Since

$$Z = \frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \sim N(0; 1)$$

for all $\mu = E(X)$,

Test on a population mean μ .

- Since

$$Z = \frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \sim N(0; 1)$$

for all $\mu = E(X)$, then

$$\beta = \Phi\left(Z_\alpha + \frac{\mu_0 - \mu}{\sigma/\sqrt{n}}\right) = \Phi\left(1.645 + \frac{17.85 - 20}{3.39/\sqrt{25}}\right) = \Phi(-1.53) = 0.063$$

Test on a population mean μ .

- Since

$$Z = \frac{\bar{X} - \mu}{\sigma/\sqrt{n}} \sim N(0; 1)$$

for all $\mu = E(X)$, then

$$\beta = \Phi\left(Z_\alpha + \frac{\mu_0 - \mu}{\sigma/\sqrt{n}}\right) = \Phi\left(1.645 + \frac{17.85 - 20}{3.39/\sqrt{25}}\right) = \Phi(-1.53) = 0.063$$

- The power of test is $1 - \beta = 1 - 0.063 = 0.937$.

Test on a population proportion p .

Example: Suppose that the proportion of defective items in a production line is 1.2%. A technical improvement was made to reduce the rate of defective items. Test 1000 items from the production line after the improvement, 8 of them were found to be defective. Do the data allow us to infer that the improvement was successful at 10% level of significance?

Test on a population proportion p .

Example: Suppose that the proportion of defective items in a production line is 1.2%. A technical improvement was made to reduce the rate of defective items. Test 1000 items from the production line after the improvement, 8 of them were found to be defective. Do the data allow us to infer that the improvement was successful at 10% level of significance?

- Let p be the proportion of defective items in the production line after the technical improvement.

Test on a population proportion p .

Example: Suppose that the proportion of defective items in a production line is 1.2%. A technical improvement was made to reduce the rate of defective items. Test 1000 items from the production line after the improvement, 8 of them were found to be defective. Do the data allow us to infer that the improvement was successful at 10% level of significance?

- Let p be the proportion of defective items in the production line after the technical improvement.
- We test the research hypothesis that $p < 0.012$

Test on a population proportion p .

Problem 2: Let p be a population proportion (for example, the proportion of defective items in a production line).

Test on a population proportion p .

Problem 2: Let p be a population proportion (for example, the proportion of defective items in a production line).

- We test the null hypotheses $H_0 : p = p_0$ against the alternative $H_1 : p > p_0$ (or $<, \neq$) at level of significance α .

Test on a population proportion p .

Problem 2: Let p be a population proportion (for example, the proportion of defective items in a production line).

- We test the null hypotheses $H_0 : p = p_0$ against the alternative $H_1 : p > p_0$ (or $<, \neq$) at level of significance α .
- Take a random sample of size n (large enough) from the population. Let \hat{p} be the sample proportion (point estimator of p). The test statistic is

$$Z = \frac{\hat{p} - p_0}{\sqrt{p_0(1 - p_0)/n}} \sim N(0; 1) \text{ if } H_0 \text{ is true}$$

Test on a population proportion p .

Problem 2: Let p be a population proportion (for example, the proportion of defective items in a production line).

- We test the null hypotheses $H_0 : p = p_0$ against the alternative $H_1 : p > p_0$ (or $<, \neq$) at level of significance α .
- Take a random sample of size n (large enough) from the population. Let \hat{p} be the sample proportion (point estimator of p). The test statistic is

$$Z = \frac{\hat{p} - p_0}{\sqrt{p_0(1 - p_0)/n}} \sim N(0; 1) \text{ if } H_0 \text{ is true}$$

- The rejection region is as the following

$$W_\alpha = \begin{cases} (Z_\alpha; +\infty), & \text{if } H_1 : p > p_0 \\ (-\infty; -Z_\alpha), & \text{if } H_1 : p < p_0 \\ (-\infty; -Z_{\alpha/2}) \cup (Z_{\alpha/2}; +\infty), & \text{if } H_1 : p \neq p_0 \end{cases}$$

Test on a population proportion p .

Example 5.3: Suppose that the proportion of defective items in a production line is 1.2%. A technical improvement was made to reduce the rate of defective items. Test 1000 items from the production line after the improvement, 8 of them were found to be defective.

- Do the data allow us to infer that the improvement was successful at 10% level of significance?
- Find the power of test if the true proportion of defective items after the technical improvement is $p_1 = 0.5\%$.

Test on a population proportion p .

Solution:

- Let p be the proportion of defective items in the production line after the technical improvement. We test the null hypotheses $H_0 : p = 0.012$ against the alternative $H_1 : p < 0.012$ at level of significance $\alpha = 0.1$.

Test on a population proportion p .

Solution:

- Let p be the proportion of defective items in the production line after the technical improvement. We test the null hypotheses $H_0 : p = 0.012$ against the alternative $H_1 : p < 0.012$ at level of significance $\alpha = 0.1$.
- The test statistic is

$$Z = \frac{\hat{p} - p_0}{\sqrt{p_0(1 - p_0)/n}} \sim N(0; 1) \text{ if } H_0 \text{ is true}$$

Test on a population proportion p .

Solution:

- Let p be the proportion of defective items in the production line after the technical improvement. We test the null hypotheses $H_0 : p = 0.012$ against the alternative $H_1 : p < 0.012$ at level of significance $\alpha = 0.1$.
- The test statistic is

$$Z = \frac{\hat{p} - p_0}{\sqrt{p_0(1 - p_0)/n}} \sim N(0; 1) \text{ if } H_0 \text{ is true}$$

For the given sample: $n = 1000$; $\hat{p} = 8/1000 = 0.008$, $p_0 = 0.012$ then

$$Z_{obs} = \frac{0.008 - 0.012}{\sqrt{0.008 * 0.992/1000}} = -1.42$$

Test on a population proportion p .

Solution:

- Let p be the proportion of defective items in the production line after the technical improvement. We test the null hypotheses $H_0 : p = 0.012$ against the alternative $H_1 : p < 0.012$ at level of significance $\alpha = 0.1$.
- The test statistic is

$$Z = \frac{\hat{p} - p_0}{\sqrt{p_0(1 - p_0)/n}} \sim N(0; 1) \text{ if } H_0 \text{ is true}$$

For the given sample: $n = 1000$; $\hat{p} = 8/1000 = 0.008$, $p_0 = 0.012$ then

$$Z_{obs} = \frac{0.008 - 0.012}{\sqrt{0.008 * 0.992/1000}} = -1.42$$

- The rejection region is $W_\alpha = (-\infty; -Z_\alpha) = (-\infty; -1.28)$.

Test on a population proportion p .

Solution:

- Let p be the proportion of defective items in the production line after the technical improvement. We test the null hypotheses $H_0 : p = 0.012$ against the alternative $H_1 : p < 0.012$ at level of significance $\alpha = 0.1$.
- The test statistic is

$$Z = \frac{\hat{p} - p_0}{\sqrt{p_0(1 - p_0)/n}} \sim N(0; 1) \text{ if } H_0 \text{ is true}$$

For the given sample: $n = 1000$; $\hat{p} = 8/1000 = 0.008$, $p_0 = 0.012$ then

$$Z_{obs} = \frac{0.008 - 0.012}{\sqrt{0.008 * 0.992/1000}} = -1.42$$

- The rejection region is $W_\alpha = (-\infty; -Z_\alpha) = (-\infty; -1.28)$.
- Since $Z_{obs} \in W_\alpha$ then we reject H_0 . Thus, there is enough evidence to infer that the technical improvement was successful at 10% level of significance.

Test on a population proportion p .

- The power of the test is

$$\begin{aligned} 1 - \beta &= P(\text{reject } H_0 \mid H_0 \text{ is false}) = P(Z \in W_\alpha \mid p = p_1) \\ &= P(Z < -Z_\alpha \mid p = p_1) = P\left(\frac{\hat{p} - p_0}{\sqrt{p_0(1 - p_0)/n}} < -Z_\alpha \mid p = p_1\right) \end{aligned}$$

Test on a population proportion p .

- The power of the test is

$$\begin{aligned}
 1 - \beta &= P(\text{reject } H_0 \mid H_0 \text{ is false}) = P(Z \in W_\alpha \mid p = p_1) \\
 &= P(Z < -Z_\alpha \mid p = p_1) = P\left(\frac{\hat{p} - p_0}{\sqrt{p_0(1 - p_0)/n}} < -Z_\alpha \mid p = p_1\right) \\
 &= P\left(\frac{\hat{p} - p_1}{\sqrt{p_1(1 - p_1)/n}} < -Z_\alpha \sqrt{\frac{p_0(1 - p_0)}{p_1(1 - p_1)}} + \frac{p_0 - p_1}{\sqrt{p_1(1 - p_1)/n}} \mid p = p_1\right)
 \end{aligned}$$

Test on a population proportion p .

- The power of the test is

$$\begin{aligned}
 1 - \beta &= P(\text{reject } H_0 \mid H_0 \text{ is false}) = P(Z \in W_\alpha \mid p = p_1) \\
 &= P(Z < -Z_\alpha \mid p = p_1) = P\left(\frac{\hat{p} - p_0}{\sqrt{p_0(1 - p_0)/n}} < -Z_\alpha \mid p = p_1\right) \\
 &= P\left(\frac{\hat{p} - p_1}{\sqrt{p_1(1 - p_1)/n}} < -Z_\alpha \sqrt{\frac{p_0(1 - p_0)}{p_1(1 - p_1)}} + \frac{p_0 - p_1}{\sqrt{p_1(1 - p_1)/n}} \mid p = p_1\right)
 \end{aligned}$$

- Since $Z = \frac{\hat{p} - p_1}{\sqrt{p_1(1 - p_1)/n}} \sim N(0, 1)$ if $p = p_1$, thus

$$1 - \beta = \Phi\left(-Z_\alpha \sqrt{\frac{p_0(1 - p_0)}{p_1(1 - p_1)}} + \frac{p_0 - p_1}{\sqrt{p_1(1 - p_1)/n}}\right)$$

Test on a population proportion p .

- When $p_1 = 0.005$, $n = 1000$, $p_0 = 0.012$, $Z_\alpha = 1.28$, the power of the test is

$$\begin{aligned} 1 - \beta &= \Phi\left(-1.28\sqrt{\frac{0.012 * 0.988}{0.005 * 0.995}} + \frac{0.012 - 0.005}{\sqrt{0.005 * 0.995/1000}}\right) \\ &= \Phi(1.16) = 0.88 \end{aligned}$$

Test on the difference between two means $\mu_1 - \mu_2$.

Example: A high school math teacher claims that students in her class will score higher on the math portion of the ACT than students in a colleague's math class. The mean ACT math score for 49 students in her class is 22.1 and the sample standard deviation is 4.8. The mean ACT math score for 44 of the colleague's students is 19.8 and the sample standard deviation is 5.4. At $\alpha = 0.1$, can the teacher's claim be supported?

Test on the difference between two means $\mu_1 - \mu_2$.

Example: A high school math teacher claims that students in her class will score higher on the math portion of the ACT than students in a colleague's math class. The mean ACT math score for 49 students in her class is 22.1 and the sample standard deviation is 4.8. The mean ACT math score for 44 of the colleague's students is 19.8 and the sample standard deviation is 5.4. At $\alpha = 0.1$, can the teacher's claim be supported?

- Let μ_1, μ_2 be the mean ACT math scores of students in her class and in her colleague's class, respectively.
- We want to test whether $\mu_1 - \mu_2 > 0$ at $\alpha = 0.1$ level of significance.

Test on the difference between two means $\mu_1 - \mu_2$.

Problem 3: We observe 2 populations X_1, X_2 where $\mu_1 = E(X_1)$, $\mu_2 = E(X_2)$ and $\sigma_1^2 = V(X_1)$, $\sigma_2^2 = V(X_2)$.

Test on the difference between two means $\mu_1 - \mu_2$.

Problem 3: We observe 2 populations X_1, X_2 where $\mu_1 = E(X_1)$, $\mu_2 = E(X_2)$ and $\sigma_1^2 = V(X_1)$, $\sigma_2^2 = V(X_2)$.

- We test on the difference between two means $\mu_1 - \mu_2$.

Test on the difference between two means $\mu_1 - \mu_2$.

Problem 3: We observe 2 populations X_1, X_2 where $\mu_1 = E(X_1)$, $\mu_2 = E(X_2)$ and $\sigma_1^2 = V(X_1)$, $\sigma_2^2 = V(X_2)$.

- We test on the difference between two means $\mu_1 - \mu_2$.
- The null hypothesis is $H_0 : \mu_1 - \mu_2 = 0$ against the alternative $H_1 : \mu_1 - \mu_2 > 0$ (or $>, \neq$).

Test on the difference between two means $\mu_1 - \mu_2$.

Problem 3: We observe 2 populations X_1, X_2 where $\mu_1 = E(X_1)$, $\mu_2 = E(X_2)$ and $\sigma_1^2 = V(X_1)$, $\sigma_2^2 = V(X_2)$.

- We test on the difference between two means $\mu_1 - \mu_2$.
- The null hypothesis is $H_0 : \mu_1 - \mu_2 = 0$ against the alternative $H_1 : \mu_1 - \mu_2 > 0$ (or $>, \neq$).
- Consider 2 independent random sample of size n_1, n_2 from 2 populations. Let \bar{X}_1, \bar{X}_2 be the sample means and S_1^2, S_2^2 be the sample variances of 2 samples, respectively. The sampling distribution of $\bar{X}_1 - \bar{X}_2$ is the following:

$$Z = \frac{\bar{X}_1 - \bar{X}_2 - (\mu_1 - \mu_2)}{\sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}}} \sim N(0, 1)$$

Test on the difference between two means $\mu_1 - \mu_2$

Case 1: Suppose that 2 populations X_1, X_2 are normal and 2 variances σ_1^2, σ_2^2 are known.

Test on the difference between two means $\mu_1 - \mu_2$

Case 1: Suppose that 2 populations X_1, X_2 are normal and 2 variances σ_1^2, σ_2^2 are known.

- The test statistic is

$$Z = \frac{\bar{X}_1 - \bar{X}_2}{\sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}}} \sim N(0, 1) \text{ if } H_0 \text{ is true}$$

Test on the difference between two means $\mu_1 - \mu_2$

Case 1: Suppose that 2 populations X_1, X_2 are normal and 2 variances σ_1^2, σ_2^2 are known.

- The test statistic is

$$Z = \frac{\bar{X}_1 - \bar{X}_2}{\sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}}} \sim N(0, 1) \text{ if } H_0 \text{ is true}$$

- The rejection region is as the following

$$W_\alpha = \begin{cases} (Z_\alpha; +\infty), & \text{if } H_1 : \mu_1 - \mu_2 > 0 \\ (-\infty; -Z_\alpha), & \text{if } H_1 : \mu_1 - \mu_2 < 0 \\ (-\infty; -Z_{\alpha/2}) \cup (Z_{\alpha/2}; +\infty), & \text{if } H_1 : \mu_1 - \mu_2 \neq 0 \end{cases}$$

Test on the difference between two means $\mu_1 - \mu_2$

Case 2: Two populations X_1, X_2 are not supposed to be normal, 2 variances σ_1^2, σ_2^2 are unknown and the sample sizes n_1, n_2 are large enough (> 30).

Test on the difference between two means $\mu_1 - \mu_2$

Case 2: Two populations X_1, X_2 are not supposed to be normal, 2 variances σ_1^2, σ_2^2 are unknown and the sample sizes n_1, n_2 are large enough (> 30).

- The test statistic is

$$T = \frac{\bar{X}_1 - \bar{X}_2}{\sqrt{\frac{S_1^2}{n_1} + \frac{S_2^2}{n_2}}} \approx N(0, 1) \text{ if } H_0 \text{ is true}$$

Test on the difference between two means $\mu_1 - \mu_2$

Case 2: Two populations X_1, X_2 are not supposed to be normal, 2 variances σ_1^2, σ_2^2 are unknown and the sample sizes n_1, n_2 are large enough (> 30).

- The test statistic is

$$T = \frac{\bar{X}_1 - \bar{X}_2}{\sqrt{\frac{S_1^2}{n_1} + \frac{S_2^2}{n_2}}} \approx N(0, 1) \text{ if } H_0 \text{ is true}$$

- The rejection region is as the following

$$W_\alpha = \begin{cases} (Z_\alpha; +\infty), & \text{if } H_1 : \mu_1 - \mu_2 > 0 \\ (-\infty; -Z_\alpha), & \text{if } H_1 : \mu_1 - \mu_2 < 0 \\ (-\infty; -Z_{\alpha/2}) \cup (Z_{\alpha/2}; +\infty), & \text{if } H_1 : \mu_1 - \mu_2 \neq 0 \end{cases}$$

Test on the difference between two means $\mu_1 - \mu_2$

Case 3: Two populations X_1, X_2 are supposed to be normal, 2 variances σ_1^2, σ_2^2 are unknown and equal ($\sigma_1^2 = \sigma_2^2$).

Test on the difference between two means $\mu_1 - \mu_2$

Case 3: Two populations X_1, X_2 are supposed to be normal, 2 variances σ_1^2, σ_2^2 are unknown and equal ($\sigma_1^2 = \sigma_2^2$).

- The test statistic is

$$T = \frac{\bar{X}_1 - \bar{X}_2}{\sqrt{\frac{S^2}{n_1} + \frac{S^2}{n_2}}} \sim t_{d.f} \text{ if } H_0 \text{ is true,}$$

where $d.f = n_1 + n_2 - 2$ and

$$S^2 = \frac{(n_1 - 1)S_1^2 + (n_2 - 1)S_2^2}{n_1 + n_2 - 2} \text{ is called the pooled variance.}$$

Test on the difference between two means $\mu_1 - \mu_2$

Case 3: Two populations X_1, X_2 are supposed to be normal, 2 variances σ_1^2, σ_2^2 are unknown and equal ($\sigma_1^2 = \sigma_2^2$).

- The test statistic is

$$T = \frac{\bar{X}_1 - \bar{X}_2}{\sqrt{\frac{S^2}{n_1} + \frac{S^2}{n_2}}} \sim t_{d.f} \text{ if } H_0 \text{ is true,}$$

where $d.f = n_1 + n_2 - 2$ and

$$S^2 = \frac{(n_1 - 1)S_1^2 + (n_2 - 1)S_2^2}{n_1 + n_2 - 2} \text{ is called the pooled variance.}$$

- The rejection region is as the following

$$W_\alpha = \begin{cases} (t_{d.f,\alpha}; +\infty), & \text{if } H_1 : \mu_1 - \mu_2 > 0 \\ (-\infty; -t_{d.f,\alpha}), & \text{if } H_1 : \mu_1 - \mu_2 < 0 \\ (-\infty; -t_{d.f,\alpha/2}) \cup (t_{d.f,\alpha/2}; +\infty), & \text{if } H_1 : \mu_1 - \mu_2 \neq 0. \end{cases}$$

Test on the difference between two means $\mu_1 - \mu_2$.

Example 5.4: A high school math teacher claims that students in her class will score higher on the math portion of the ACT than students in a colleague's math class. The mean ACT math score for 49 students in her class is 22.1 and the sample standard deviation is 4.8. The mean ACT math score for 44 of the colleague's students is 19.8 and the sample standard deviation is 5.4. At $\alpha = 0.1$, can the teacher's claim be supported?

Test on the difference between two means $\mu_1 - \mu_2$.

Example 5.4: A high school math teacher claims that students in her class will score higher on the math portion of the ACT than students in a colleague's math class. The mean ACT math score for 49 students in her class is 22.1 and the sample standard deviation is 4.8. The mean ACT math score for 44 of the colleague's students is 19.8 and the sample standard deviation is 5.4. At $\alpha = 0.1$, can the teacher's claim be supported?

Solution:

- Let μ_1, μ_2 be the mean ACT math scores of students in her class and in her colleague's class, respectively. We test $H_0 : \mu_1 - \mu_2 = 0$ versus $H_1 : \mu_1 - \mu_2 > 0$ at $\alpha = 0.1$ level of significance.

Test on the difference between two means $\mu_1 - \mu_2$.

- It is not supposed that the populations are normal and the two population variances are unknown, but the sample sizes $n_1 = 49$, $n_2 = 44$ are large enough. Thus, the test statistic is

$$T = \frac{\bar{X}_1 - \bar{X}_2}{\sqrt{\frac{S_1^2}{n_1} + \frac{S_2^2}{n_2}}} \approx N(0, 1) \text{ if } H_0 \text{ is true}$$

Test on the difference between two means $\mu_1 - \mu_2$.

- It is not supposed that the populations are normal and the two population variances are unknown, but the sample sizes $n_1 = 49, n_2 = 44$ are large enough. Thus, the test statistic is

$$T = \frac{\bar{X}_1 - \bar{X}_2}{\sqrt{\frac{S_1^2}{n_1} + \frac{S_2^2}{n_2}}} \approx N(0, 1) \text{ if } H_0 \text{ is true}$$

For the given sample, we have: $n_1 = 49, n_2 = 44; \bar{x}_1 = 22.1, \bar{x}_2 = 19.8, s_1 = 4.8, s_2 = 5.4$, then

$$T_{obs} = \frac{22.1 - 19.8}{\sqrt{\frac{4.8^2}{49} + \frac{5.4^2}{44}}} = 2.16$$

Test on the difference between two means $\mu_1 - \mu_2$.

- The rejection region is $W_\alpha = (Z_\alpha; +\infty) = (Z_{0.1}; +\infty) = (1.28; +\infty)$.

Test on the difference between two means $\mu_1 - \mu_2$.

- The rejection region is $W_\alpha = (Z_\alpha; +\infty) = (Z_{0.1}; +\infty) = (1.28; +\infty)$.
- Since $T_{obs} = 2.16 \in W_\alpha$ then we reject H_0 . Thus, there is enough evidence at the 10% level to support the teacher's claim that her students score better on the ACT.

Test on the difference between two means $\mu_1 - \mu_2$.

Example 5.5: To find out whether a new serum will arrest leukemia, 9 mice, all with an advanced stage of the disease, are selected. Five mice receive the treatment and 4 do not. Survival times, in years, from the time the experiment commenced are as follows:

- Treatment: 2.1, 5.3, 1.4, 4.6, 0.9
- No Treatment: 1.9, 0.5, 2.8, 3.1

At the 0.05 level of significance, can the serum be said to be effective? Assume the two populations to be normally distributed with equal variances.

Test on the difference between two means $\mu_1 - \mu_2$.

Example 5.5: To find out whether a new serum will arrest leukemia, 9 mice, all with an advanced stage of the disease, are selected. Five mice receive the treatment and 4 do not. Survival times, in years, from the time the experiment commenced are as follows:

- Treatment: 2.1, 5.3, 1.4, 4.6, 0.9
- No Treatment: 1.9, 0.5, 2.8, 3.1

At the 0.05 level of significance, can the serum be said to be effective? Assume the two populations to be normally distributed with equal variances.

Solution:

- Let X_1, X_2 be the survival times, in years, of mice with and without treatment of the new serum. To test whether the serum is said to be effective, we test $H_0 : \mu_1 - \mu_2 = 0$ versus $H_1 : \mu_1 - \mu_2 > 0$ at $\alpha = 0.05$ level of significance.

Test on the difference between two means $\mu_1 - \mu_2$.

- It is supposed that $X_1 \sim N(\mu_1, \sigma_1^2)$, $X_2 \sim N(\mu_2, \sigma_2^2)$ where σ_1, σ_2 are unknown but equal. Thus, the test statistic is

$$T = \frac{\bar{X}_1 - \bar{X}_2}{\sqrt{\frac{S^2}{n_1} + \frac{S^2}{n_2}}} \sim t_{n_1+n_2-2} \text{ if } H_0 \text{ is true}$$

Test on the difference between two means $\mu_1 - \mu_2$.

- It is supposed that $X_1 \sim N(\mu_1, \sigma_1^2)$, $X_2 \sim N(\mu_2, \sigma_2^2)$ where σ_1, σ_2 are unknown but equal. Thus, the test statistic is

$$T = \frac{\bar{X}_1 - \bar{X}_2}{\sqrt{\frac{S^2}{n_1} + \frac{S^2}{n_2}}} \sim t_{n_1+n_2-2} \text{ if } H_0 \text{ is true}$$

For the given sample, we have: $n_1 = 5, n_2 = 4$,

$$\bar{x}_1 = (2.1 + 5.3 + 1.4 + 4.6 + 0.9)/5 = 2.86,$$

$$\bar{x}_2 = (1.9 + 0.5 + 2.8 + 3.1)/4 = 2.075,$$

$$s_1^2 = (2.1^2 + \dots + 0.9^2 - 5 * 2.86^2)/4 = 3.883,$$

$$s_2^2 = (1.9^2 + \dots + 3.1^2 - 4 * 2.075^2)/3 = 1.3625, \text{ then}$$

$$s^2 = \frac{(n_1 - 1)s_1^2 + (n_2 - 1)s_2^2}{n_1 + n_2 - 2} = \frac{4 * 3.883 + 3 * 1.3625}{7} = 2.803$$

Test on the difference between two means $\mu_1 - \mu_2$.

- Thus

$$T_{obs} = \frac{2.86 - 2.075}{\sqrt{\frac{2.803}{5} + \frac{2.803}{4}}} = 0.699$$

Test on the difference between two means $\mu_1 - \mu_2$.

- Thus

$$T_{obs} = \frac{2.86 - 2.075}{\sqrt{\frac{2.803}{5} + \frac{2.803}{4}}} = 0.699$$

- The rejection region is

$$W_{\alpha} = (t_{n_1+n_2-2;\alpha}; +\infty) = (t_{7;0.05}; +\infty) = (1.89; +\infty).$$

Test on the difference between two means $\mu_1 - \mu_2$.

- Thus

$$T_{obs} = \frac{2.86 - 2.075}{\sqrt{\frac{2.803}{5} + \frac{2.803}{4}}} = 0.699$$

- The rejection region is $W_\alpha = (t_{n_1+n_2-2;\alpha}; +\infty) = (t_{7;0.05}; +\infty) = (1.89; +\infty)$.
- Since $T_{obs} = 0.699 \notin W_\alpha$ then we fail to reject H_0 . Thus, there is not enough evidence at the 5% level to infer that the new serum is said to be effective.

Test on the difference between two proportions $p_1 - p_2$.

Problem 4: Let p_1, p_2 be two population proportions (for example, proportions of defective items in 2 different production lines).

Test on the difference between two proportions $p_1 - p_2$.

Problem 4: Let p_1, p_2 be two population proportions (for example, proportions of defective items in 2 different production lines). We consider the following test: $H_0 : p_1 - p_2 = 0$ against $H_1 : p_1 - p_2 > 0$ (or $>, \neq$).

Test on the difference between two proportions $p_1 - p_2$.

Problem 4: Let p_1, p_2 be two population proportions (for example, proportions of defective items in 2 different production lines). We consider the following test: $H_0 : p_1 - p_2 = 0$ against $H_1 : p_1 - p_2 > 0$ (or $>, \neq$).

- Draw 2 independent random samples of size n_1, n_2 from 2 populations.

Test on the difference between two proportions $p_1 - p_2$.

Problem 4: Let p_1, p_2 be two population proportions (for example, proportions of defective items in 2 different production lines). We consider the following test: $H_0 : p_1 - p_2 = 0$ against $H_1 : p_1 - p_2 > 0$ (or $< 0, \neq$).

- Draw 2 independent random samples of size n_1, n_2 from 2 populations.
- Let \hat{p}_1, \hat{p}_2 be two sample proportions, the point estimates of p_1, p_2 , respectively.

Test on the difference between two proportions $p_1 - p_2$.

Problem 4: Let p_1, p_2 be two population proportions (for example, proportions of defective items in 2 different production lines). We consider the following test: $H_0 : p_1 - p_2 = 0$ against $H_1 : p_1 - p_2 > 0$ (or < 0).

- Draw 2 independent random samples of size n_1, n_2 from 2 populations.
- Let \hat{p}_1, \hat{p}_2 be two sample proportions, the point estimates of p_1, p_2 , respectively.
- The test statistic is

$$Z = \frac{\hat{p}_1 - \hat{p}_2}{\sqrt{\hat{p}(1 - \hat{p})\left(\frac{1}{n_1} + \frac{1}{n_2}\right)}} \approx N(0, 1) \text{ if } H_0 \text{ is true,}$$

where \hat{p} is the weighted estimate of p_1 and p_2 :

$$\hat{p} = \frac{n_1 \hat{p}_1 + n_2 \hat{p}_2}{n_1 + n_2}$$

Test on the difference between two proportions $p_1 - p_2$

- Remark that

$$\hat{p}_1 = \frac{m_1}{n_1}; \hat{p}_2 = \frac{m_2}{n_2}; \hat{p} = \frac{m_1 + m_2}{n_1 + n_2},$$

where m_1, m_2 are the number of successes in 2 samples, respectively.

Test on the difference between two proportions $p_1 - p_2$

- Remark that

$$\hat{p}_1 = \frac{m_1}{n_1}; \hat{p}_2 = \frac{m_2}{n_2}; \hat{p} = \frac{m_1 + m_2}{n_1 + n_2},$$

where m_1, m_2 are the number of successes in 2 samples, respectively.

- The rejection region is as the following

$$W_\alpha = \begin{cases} (Z_\alpha; +\infty), & \text{if } H_1 : p_1 - p_2 > 0 \\ (-\infty; -Z_\alpha), & \text{if } H_1 : p_1 - p_2 < 0 \\ (-\infty; -Z_{\alpha/2}) \cup (Z_{\alpha/2}; +\infty), & \text{if } H_1 : p_1 - p_2 \neq 0 \end{cases}$$

Test on the difference between two proportions $p_1 - p_2$

- Remark that

$$\hat{p}_1 = \frac{m_1}{n_1}; \hat{p}_2 = \frac{m_2}{n_2}; \hat{p} = \frac{m_1 + m_2}{n_1 + n_2},$$

where m_1, m_2 are the number of successes in 2 samples, respectively.

- The rejection region is as the following

$$W_\alpha = \begin{cases} (Z_\alpha; +\infty), & \text{if } H_1 : p_1 - p_2 > 0 \\ (-\infty; -Z_\alpha), & \text{if } H_1 : p_1 - p_2 < 0 \\ (-\infty; -Z_{\alpha/2}) \cup (Z_{\alpha/2}; +\infty), & \text{if } H_1 : p_1 - p_2 \neq 0 \end{cases}$$

- The sample sizes n_1, n_2 must be large enough (we usually suppose that $n_1 p_1, n_1(1 - p_1), n_2 p_2, n_2(1 - p_2)$ are ≥ 5).

Test on the difference between two proportions $p_1 - p_2$.

Example 5.6: A recent survey stated that male college students smoke less than female college students. In a survey of 1245 male students, 361 said they smoke at least one pack of cigarettes a day. In a survey of 1065 female students, 341 said they smoke at least one pack a day. At $\alpha = 0.1$, can you support the claim that the proportion of male college students who smoke at least one pack of cigarettes a day is lower than the proportion of female college students who smoke at least one pack a day?

Test on the difference between two proportions $p_1 - p_2$.

Example 5.6: A recent survey stated that male college students smoke less than female college students. In a survey of 1245 male students, 361 said they smoke at least one pack of cigarettes a day. In a survey of 1065 female students, 341 said they smoke at least one pack a day. At $\alpha = 0.1$, can you support the claim that the proportion of male college students who smoke at least one pack of cigarettes a day is lower than the proportion of female college students who smoke at least one pack a day?

Solution:

- Let p_1, p_2 be the proportions of male and of female college students who smoke at least one pack of cigarettes a day, respectively. We test $H_0 : p_1 - p_2 = 0$ versus $H_1 : p_1 - p_2 < 0$ at $\alpha = 0.1$ level of significance.

Test on the difference between two proportions $p_1 - p_2$.

- The test statistic is

$$Z = \frac{\hat{p}_1 - \hat{p}_2}{\sqrt{\hat{p}(1 - \hat{p})\left(\frac{1}{n_1} + \frac{1}{n_2}\right)}} \approx N(0, 1) \text{ if } H_0 \text{ is true.}$$

Test on the difference between two proportions $p_1 - p_2$.

- The test statistic is

$$Z = \frac{\hat{p}_1 - \hat{p}_2}{\sqrt{\hat{p}(1 - \hat{p})\left(\frac{1}{n_1} + \frac{1}{n_2}\right)}} \approx N(0, 1) \text{ if } H_0 \text{ is true.}$$

For the given samples, we have: $n_1 = 1245, n_2 = 1065$;
 $m_1 = 361, m_2 = 341$,

Test on the difference between two proportions $p_1 - p_2$.

- The test statistic is

$$Z = \frac{\hat{p}_1 - \hat{p}_2}{\sqrt{\hat{p}(1 - \hat{p})\left(\frac{1}{n_1} + \frac{1}{n_2}\right)}} \approx N(0, 1) \text{ if } H_0 \text{ is true.}$$

For the given samples, we have: $n_1 = 1245, n_2 = 1065$;
 $m_1 = 361, m_2 = 341$, then

$$\hat{p}_1 = \frac{361}{1245} = 0.29; \hat{p}_2 = \frac{341}{1065} = 0.32; \hat{p} = \frac{361 + 341}{1245 + 1065} = 0.304$$

Test on the difference between two proportions $p_1 - p_2$.

- The test statistic is

$$Z = \frac{\hat{p}_1 - \hat{p}_2}{\sqrt{\hat{p}(1 - \hat{p})\left(\frac{1}{n_1} + \frac{1}{n_2}\right)}} \approx N(0, 1) \text{ if } H_0 \text{ is true.}$$

For the given samples, we have: $n_1 = 1245, n_2 = 1065$;
 $m_1 = 361, m_2 = 341$, then

$$\hat{p}_1 = \frac{361}{1245} = 0.29; \hat{p}_2 = \frac{341}{1065} = 0.32; \hat{p} = \frac{361 + 341}{1245 + 1065} = 0.304$$

Thus,

$$Z = \frac{0.29 - 0.32}{\sqrt{0.304 * 0.694\left(\frac{1}{1245} + \frac{1}{1065}\right)}} = -1.56$$

Test on the difference between two proportions $p_1 - p_2$.

- The rejection region is

$$W_\alpha = (-\infty; -Z_\alpha) = (-\infty; Z_{0.1}) = (-\infty; -1.28).$$

Test on the difference between two proportions $p_1 - p_2$.

- The rejection region is $W_\alpha = (-\infty; -Z_\alpha) = (-\infty; Z_{0.1}) = (-\infty; -1.28)$.
- Since $Z_{obs} = -1.56 \in W_\alpha$ then we reject H_0 . Thus, there is enough evidence at the 10% level to claim that the proportion of male college students who smoke at least one pack of cigarettes a day is lower than the proportion of female college students who smoke at least one pack a day.