

Verified Integer Rational Database

Round Three

Registry: [[HOWL-DATA-3-2026](#)]

Series Path: [[HOWL-DATA-1-2026](#)] → [[HOWL-DATA-2-2026](#)] → [[HOWL-DATA-3-2026](#)]

DOI: 10.5281/zenodo.19528532

Date: April 1 2026

Domain: Cross Domain Data

Status: Documentation

AI Usage Disclosure: Only the top metadata, figures, refs and final copyright sections were edited by the author. All paper content was LLM-generated using Anthropic's Claude Opus 4.6.

Abstract

DATA-3 is the verified successor to DATA-2. The same 123 entries (107 original Q335 conversions + 16 Koide-derived quantities) are carried forward with zero numerical changes. What DATA-3 adds is a certificate of internal consistency: 32 independent cross-checks covering mass ratios, analytical constant identities, physical relations, Koide derivations, and exact SI values. All 32 pass.

One structural finding emerged from the verification: the lattice QCD mass ratios (m_c/m_s , m_b/m_c , m_u/m_d) are independent data that cannot be derived from the individual PDG quark masses by division. The discrepancy (up to 28%) is the renormalization scale mismatch between PDG masses evaluated at different scales and lattice ratios evaluated at a common scale. These entries are annotated as independent measurements.

DATA-3 is the sole data reference for all future HOWL computation. DATA-2 is retired.

1. Relationship to DATA-1 and DATA-2

DATA-1 collected 268 precision values across 17 domains. DATA-2 converted 107 of these to the $Q335 = 2^{335}$ integer rational basis, performed multi-base scans across 19 bases with 90 control irrationals, ran continued fraction analysis, and added 16 Koide-derived entries. DATA-2 confirmed the Q335 basis and established that no measured constant has a compact rational representation in any basis tested.

DATA-3 inherits every entry and finding from DATA-2 and adds verification. No value changes. The promotion from DATA-2 to DATA-3 means: every testable internal relation has been checked and passes.

2. The Verification

2.1 Method

For each pair of entries related by a known identity, compute both sides independently from the stored Fraction values and compare. A test passes if the relative difference is below 10^{-N} where N is the number of significant digits of the least-precise input.

All computation uses Python Fraction arithmetic. Square roots use mpmath at 120-digit precision converted to Fraction. No floating point enters the comparison logic.

2.2 Results Summary

Group	Tests	Pass	Description
A: Mass ratios	9	9	Stored ratios vs ratios computed from individual masses
B: Analytical constants	8	8	Q335 numerator identities and mpmath cross-checks
C: Physical relations	4	4	R_∞ , a_0 , μ_0 from defining formulas; m D from constituents
D: Koide derived	5	5	K1-K16 recomputed from source masses
E: SI exact constants	6	6	c, h, e, k B, N A, $\Delta \nu$ Cs at exact SI 2019 values
Total	32	32	All pass

2.3 Group A: Mass Ratio Identities

Test	Relation	Digits Achieved	Digits Required
A1	m_p/m_e stored vs m_p/m_e	12.2	11
A2	m_μ/m_e stored vs m_μ/m_e	21.0	10

Test	Relation	Digits Achieved	Digits Required
A3	m_{τ}/m_e stored vs m_{τ} / m_e	6.3	6
A4	m_{τ}/m_{μ} stored vs m_{τ} / m_{μ}	5.8	5
A5	m_n/m_p stored vs m_n / m_p	20.0	11
A6	M_W/M_Z stored vs M_W / M_Z	7.2	6
A7	$m_n - m_p$ stored vs $m_n - m_p$	exact	8
A8	m_H/M_Z stored vs m_H / M_Z	5.6	5
A9	m_t/M_Z stored vs m_t / M_Z	5.9	5

Every stored ratio is consistent with the ratio computed from the individual mass entries. The margin ranges from 0.3 digits (A3: m_{τ}/m_e at 6.3 vs 6 required) to 10 digits (A5: m_n/m_p at 20.0 vs 11 required). The exact match on A7 ($m_n - m_p$) confirms that the neutron mass, proton mass, and mass difference entries are arithmetically consistent to all stored digits.

2.4 Group B: Analytical Constant Identities

Test	Relation	Digits Achieved	Digits Required
B1	$R_2 = \pi^2/4$ (Q335 vs mpmath)	102.0	100
B2	$R_4 = \pi^4/32$ (Q335 vs mpmath)	102.4	100
B3	$2\pi = 8R_2$	exact	100
B4	$\zeta(2) = \pi^2/6$	101.5	99
B5	α/π vs $1/(\alpha^{-1} \times \pi)$	13.9	12
B6	$\varphi = (1 + \sqrt{5})/2$	101.4	100
B7	π^2 stored vs $\pi \times \pi$	101.7	99
B8	2π stored vs $2 \times \pi$	exact	100

The Q335 basis is internally consistent at its full 101-digit capacity. The exact matches on B3 and B8 confirm that the numerators for 2π and π satisfy the integer identity $2p_{\pi} = p_{2\pi}$ within the Q335 representation. The 101-102 digit matches on B1, B2, B4, B6, B7 confirm that the independently computed numerators for R_2 , R_4 , $\zeta(2)$, φ , and π^2 are consistent with their defining formulas to the precision of the basis.

2.5 Group C: Physical Relations

Test	Relation	Digits Achieved	Digits Required
C1	$R_{\infty} = \alpha^2 m_e c / (2h)$	11.3	11
C2	$a_0 = \hbar / (m_e c \alpha)$	11.4	11
C3	$\mu_0 = 2\alpha h / (c e^2)$	11.9	11
C4	$m_D = m_p + m_n - E_D$	9.9	8

These are the most important tests. They verify that the measured constants stored at different precisions from different sources are mutually consistent through the exact physical relations connecting them.

C1 confirms that R_∞ (13 digits), α (12 digits), and m_e (11 digits) satisfy the Rydberg formula using the exact SI values of c and h . This is how CODATA determines m_e , so agreement is expected — but it verifies that our database entries reproduce the CODATA consistency.

C2 confirms the Bohr radius a_0 (12 digits) is consistent with $\hbar$, m_e , c , and α through $a_0 = \hbar/(m_e c \alpha)$. The 11.4-digit agreement is limited by m_e at 11 digits.

C3 confirms the vacuum permeability μ_0 (12 digits) satisfies the exact SI 2019 relation $\mu_0 = 2\alpha h/(ce^2)$. In the 2019 SI system, h , c , and e are exact, so μ_0 is determined entirely by α . The 11.9-digit agreement is consistent with α at 12 digits minus rounding at the last digit.

C4 confirms the deuteron mass m_D (12 digits) equals $m_p + m_n - E_D$ to 9.9 digits, limited by the binding energy E_D at 8 digits.

An additional soft check (not scored) compared the H 1S-2S transition frequency to $(3/4)cR_\infty$ and found 3.3-digit agreement, consistent with the known ~ 500 ppm QED, Lamb shift, and recoil corrections that separate the leading-order Bohr formula from the measured frequency.

2.6 Group D: Koide Derived Entries

Test	Relation	Digits Achieved	Digits Required
D1	$K(e, \mu, \tau)$ from masses	10.3	6
D2	$a(\text{leptons})$ from K	10.2	6
D3	$M(\text{leptons}) = (\Sigma \sqrt{m})/3$	4.6	4
D4	$K(u, c, t)$ from masses	10.5	3
D5	$K(d, s, b)$ from masses	10.4	3

All 16 Koide addendum entries (K1-K16) trace back to the source masses. The Koide ratios for all three sectors agree to 10+ digits when recomputed, far exceeding the 3-6 digit requirement set by the least-precise input masses. The scale parameter $M(\text{leptons})$ agrees to 4.6 digits against a 4-digit floor, the tightest margin in Group D.

2.7 Group E: SI Exact Constants

Test	Relation	Result
E1	$h \times \Delta \nu$ Cs product	exact
E2	$c = 299792458$	exact
E3	$h = 6.62607015 \times 10^{-34}$	exact
E4	$e = 1.602176634 \times 10^{-19}$	exact

Test	Relation	Result
E5	$k B = 1.380649 \times 10^{-23}$	exact
E6	$N A = 6.02214076 \times 10^{23}$	exact

All six SI defining constants are stored at their exact values with zero discrepancy. These are the anchors of the unit system and must be exact by construction.

2.8 Group F: Lattice Ratio Annotation

The verification process revealed that three entries cannot be cross-checked by simple division of other entries:

Entry	Lattice Value	PDG Ratio	Discrepancy
m c/m s (entry 41)	11.783	m c/m s = 13.615	15.5%
m b/m c (entry 42)	4.578	m b/m c = 3.286	28.2%
m u/m d (entry 43)	0.485	m u/m d = 0.460	5.2%

This is not a database error. The individual PDG quark masses are evaluated at different renormalization scales (m_u, m_d, m_s at 2 GeV MS-bar; m_c at m_c; m_b at m_b). The lattice ratios are evaluated at a common scale by lattice QCD collaborations. The renormalization group running between scales changes the ratio by the amounts observed. Entries 41-43 are independent data from lattice QCD and are not derivable from entries 33-37.

This annotation is the one structural finding from the DATA-3 verification. It does not affect any numerical value but clarifies the independence structure of the database: 123 entries contain information from $123 - 16$ (Koide derived) = 107 independent sources, of which the lattice ratios are independent of the individual quark masses despite appearing to be redundant.

3. Database Content

DATA-3 contains 123 entries unchanged from DATA-2:

Category	Count	Description
Type E (Exact defined)	16	SI constants, defined reference values
Type M (Measured)	61	CODATA 2022, LEP/PDG, clock frequencies
Type A (Analytical)	17	Transcendental constants at 105 digits

Category	Count	Description
Type S (Standard nominal)	13	Engineering specifications
Type K (Koide derived)	16	Koide ratios, amplitudes, phases, scales
Total	123	

Precision tiers:

Tier	Source Digits	Count	Use
1	≥ 10	41	Primary targets for precision computation
2	5-9	37	Electroweak observables, Koide
3	< 5	29	Engineering, coarse measurements
K	3-6	16	Derived from Tier 1-2 masses

4. What DATA-3 Certifies

Internal consistency. Every testable relation between entries holds to the precision of the least-precise input. No entry contradicts another.

Q335 fidelity. The integer rational basis 2^{335} faithfully represents all 107 original values. The analytical constants are consistent with their defining formulas at 101+ digits. The basis is validated (from DATA-2) against 19 alternative bases and 90 control irrationals.

Physical consistency. The CODATA web of relations ($R_\infty \leftrightarrow \alpha \leftrightarrow m_e \leftrightarrow a_0 \leftrightarrow \mu_0$) is satisfied to 11+ digits. The nuclear binding relation $m_D = m_p + m_n - E_D$ holds to 9.9 digits. The Koide-derived quantities reproduce from the source masses.

Independence structure. 107 original entries from independent sources. 16 Koide entries derived from source masses. 3 lattice ratio entries independent of the individual quark masses (annotated, not derivable by division).

5. Operational Rule

DATA-3 is the sole data reference for all future HOWL computation. DATA-2 is retired. Any script or paper that previously referenced DATA-2 values now references DATA-3. The numerical content is identical; the distinction is the verification certificate.

When new measurements become available (e.g., improved m_τ from Belle II, updated PDG values, new lattice computations), they enter as DATA-3 amendments with the same verification protocol: every affected cross-check must pass before the amendment is accepted.

DATA-3: 123 entries, 32 cross-checks, 32 pass. The foundation is verified.

=====

HOWL-DATA-3: DATABASE CONSISTENCY VERIFICATION

=====

GROUP A: MASS RATIO IDENTITIES

[PASS] A1: m_p/m_e direct vs m_p / m_e

Computed: 1.836152673431236e+03

Stored: 1.836152673430000e+03

Relative diff: 6.73e-13 (12.2 digits, need 11)

[PASS] A2: m_{μ}/m_e direct vs m_{μ} / m_e

Computed: 2.067682827084672e+02

Stored: 2.067682827084672e+02

Relative diff: 1.01e-21 (21.0 digits, need 10)

[PASS] A3: m_{τ}/m_e direct vs m_{τ} / m_e

Computed: 3.477228275323682e+03

Stored: 3.477230000000000e+03

Relative diff: 4.96e-07 (6.3 digits, need 6)

[PASS] A4: m_{τ}/m_{μ} direct vs m_{τ} / m_{μ}

Computed: 1.681702933242618e+01

Stored: 1.681700000000000e+01

Relative diff: 1.74e-06 (5.8 digits, need 5)

[PASS] A5: m_n/m_p direct vs m_n / m_p

Computed: 1.001378419463362e+00

Stored: 1.001378419463362e+00

Relative diff: 9.40e-21 (20.0 digits, need 11)

[PASS] A6: M_W/M_Z direct vs M_W / M_Z

Computed: 8.813610622496918e-01

Stored: 8.813610000000000e-01

Relative diff: 7.06e-08 (7.2 digits, need 6)

[PASS] A7: $m_n - m_p$ direct vs stored difference

Computed: 1.293332510000000e+00

Stored: 1.293332510000000e+00

Agreement: EXACT

[PASS] A8: m_H/M_Z direct vs stored ratio

Computed: 1.372993696511368e+00

Stored: 1.372990000000000e+00

Relative diff: 2.69e-06 (5.6 digits, need 5)

[PASS] A9: m_t/M_Z direct vs stored ratio

Computed: 1.892472222100373e+00

Stored: 1.892470000000000e+00

Relative diff: 1.17e-06 (5.9 digits, need 5)

GROUP B: ANALYTICAL CONSTANT IDENTITIES (Q335 basis)

```

-----

[PASS] B1:  $R_2 = \pi / 4$  (Q335 vs mpmath@120)
Q335: 7.853981633974483e-01
mpmath: 7.853981633974483e-01
Relative diff: 9.13e-103 (102.0 digits, need 100)

[PASS] B2:  $R_4 = \pi^2 / 32$  (Q335 vs mpmath@120)
Q335: 3.084251375340424e-01
mpmath: 3.084251375340424e-01
Relative diff: 3.79e-103 (102.4 digits, need 100)

[PASS] B3:  $2\pi = 8R_2$  (exact within Q335)
Computed: 6.283185307179586e+00
Stored: 6.283185307179586e+00
Agreement: EXACT

[PASS] B4:  $\zeta(2) = \pi^2 / 6$  (Q335 numerators)
Computed: 1.644934066848226e+00
Stored: 1.644934066848226e+00
Relative diff: 2.90e-102 (101.5 digits, need 99)

[PASS] B5:  $\alpha / \pi$  stored vs  $1 / (\alpha^{-1} \times \pi)$ 
Computed: 2.322819464195329e-03
Stored: 2.322819464195300e-03
Relative diff: 1.24e-14 (13.9 digits, need 12)

[PASS] B6:  $\varphi = (1 + \sqrt{5}) / 2$  (Q335)
Computed: 1.618033988749895e+00

```

Stored: 1.618033988749895e+00

Relative diff: 4.42e-102 (101.4 digits, need 100)

[PASS] B7: π^2 stored vs $\pi \times \pi$ (Q335)

Computed: 9.869604401089358e+00

Stored: 9.869604401089358e+00

Relative diff: 2.21e-102 (101.7 digits, need 99)

[PASS] B8: 2π stored vs $2 \times \pi$ (exact)

Computed: 6.283185307179586e+00

Stored: 6.283185307179586e+00

Agreement: EXACT

GROUP C: PHYSICAL RELATIONS (SI 2019)

[PASS] C1: $R_\infty = \alpha^2 m_e c / (2h)$

From α , m_e , c , h : 1.097373156809536e+07

Stored R_∞ : 1.097373156815700e+07

Relative diff: 5.62e-12 (11.3 digits, need 11)

[PASS] C2: $a_0 = \hbar / (m_e c \alpha)$

From $\hbar$, m_e , c , α : 5.291772105462769e-11

Stored a_0 : 5.291772105440000e-11

Relative diff: 4.30e-12 (11.4 digits, need 11)

[PASS] C3: $\mu_0 = 2ah / (ce^2)$

From α , h , c , e : 1.256637061268226e-06

Stored μ_0 : 1.256637061270000e-06

Relative diff: 1.41e-12 (11.9 digits, need 11)

[PASS] C4: $m_D = m_p + m_n - E_D$

From $m_p + m_n - E_D$: 1.875612945230000e+03

Stored m_D : 1.875612945000000e+03

Relative diff: 1.23e-10 (9.9 digits, need 8)

[INFO] C5: H 1S-2S leading order (soft check)

Measured: 2466061413187018 Hz

(3/4)cR ∞ : 2467381470187486 Hz

Rel diff: 5.35e-04 (3.3 digits)

Note: ~5 digit agreement expected (Lamb shift, QED, recoil)

GROUP D: KOIDE DERIVED ENTRIES (K1-K16)

[PASS] D1: Koide $K(e, \mu, \tau)$ from masses

From m_e, m_μ, m_τ : 6.666605114655210e-01

Stored K1: 6.666605115000001e-01

Relative diff: 5.17e-11 (10.3 digits, need 6)

[PASS] D2: $a(\text{leptons})$ from K

Computed: 1.414200505100000e+00

Stored K4: 1.414200505200000e+00

Relative diff: 7.07e-11 (10.2 digits, need 6)

[PASS] D3: $M(\text{leptons}) = (\Sigma\sqrt{m})/3$

Computed: 1.771556100000000e+01

Stored K14: 1.771600000000000e+01

Relative diff: 2.48e-05 (4.6 digits, need 4)

[PASS] D4: Koide $K(u,c,t)$ from masses

Computed: 8.487935475760000e-01

Stored K2: 8.487935476000000e-01

Relative diff: 2.83e-11 (10.5 digits, need 3)

[PASS] D5: Koide $K(d,s,b)$ from masses

Computed: 7.312875768270000e-01

Stored K3: 7.312875768000000e-01

Relative diff: 3.69e-11 (10.4 digits, need 3)

GROUP E: EXACT SI CONSTANTS

[PASS] E1: $h \times \Delta \nu_{\text{Cs}}$ product consistency

Computed: 6.091102297113866e-24

Stored: 6.091102297113866e-24

Agreement: EXACT

[PASS] E2: c is exact integer 299792458

Computed: 2.997924580000000e+08

Stored: 2.997924580000000e+08

Agreement: EXACT

[PASS] E3: h exact SI 2019 value

Computed: 6.626070150000000e-34

Stored: 6.626070150000000e-34

Agreement: EXACT

[PASS] E4: e exact SI 2019 value

Computed: 1.6021766340000000e-19

Stored: 1.6021766340000000e-19

Agreement: EXACT

[PASS] E5: k_B exact SI 2019 value

Computed: 1.3806490000000000e-23

Stored: 1.3806490000000000e-23

Agreement: EXACT

[PASS] E6: N_A exact SI 2019 value

Computed: 6.0221407600000000e+23

Stored: 6.0221407600000000e+23

Agreement: EXACT

GROUP F: LATTICE RATIO ANNOTATION

[NOTE] Lattice ratios (m_c/m_s , m_b/m_c , m_u/m_d) are evaluated
at a COMMON renormalization scale by lattice QCD collaborations.

The individual PDG masses (m_u , m_d , m_s at 2 GeV $\overline{\text{MS}}$ -
bar;

m_c at m_c ; m_b at m_b) are at DIFFERENT scales.

Simple division of PDG masses does NOT reproduce lattice ratios.

This is not a database error – it is a renormalization scale mismatch.

Entries 41 ($m_c/m_s = 11.783$), 42 ($m_b/m_c = 4.578$),

43 ($m_u/m_d = 0.485$) are INDEPENDENT data from lattice QCD
and are NOT derivable from entries 33-37.

Discrepancies if naively compared:

m_c/m_s : PDG gives 13.615, lattice = 11.783

m_b/m_c : PDG gives 3.286, lattice = 4.578

m_u/m_d : PDG gives 0.460, lattice = 0.485

RESULTS

ID	Test	Digits	Need	Result
A1	m_p/m_e direct vs m_p / m_e	12.2	11	PASS
A2	m_{μ}/m_e direct vs m_{μ} / m_e		21.0	10
A3	m_{τ}/m_e direct vs m_{τ} / m_e	6.3	6	PASS
A4	m_{τ}/m_{μ} direct vs m_{τ} / m_{μ}		5.8	5
A5	m_n/m_p direct vs m_n / m_p	20.0	11	PASS
A6	M_W/M_Z direct vs M_W / M_Z	7.2	6	PASS
A7	$m_n - m_p$ direct vs stored difference	exact	8	PASS
A8	m_H/M_Z direct vs stored ratio	5.6	5	PASS
A9	m_t/M_Z direct vs stored ratio	5.9	5	PASS
B1	$R_2 = \pi/4$ (Q335 vs mpmath@120)		102.0	100 PASS
B2	$R_4 = \pi^2/32$ (Q335 vs mpmath@120)		102.4	100 PAS

B3	$2\pi = 8R_2$ (exact within Q335)	exact	100	PASS
B4	$\zeta(2) = \pi^2/6$ (Q335 numerators)	101.5	99	PASS
B5	α/π stored vs $1/(\alpha^{-1} \times \pi)$			13.9
B6	$\psi = (1 + \sqrt{5})/2$ (Q335)	101.4	100	PASS
B7	π^2 stored vs $\pi \times \pi$ (Q335)			
B8	2π stored vs $2 \times \pi$ (exact)			exact
C1	$R_\infty = \alpha^2 m_{ec}/(2h)$	11.3	11	PASS
C2	$a_0 = \hbar/(m_e c \alpha)$	11.4	11	PASS
C3	$\mu_0 = 2ah/(ce^2)$	11.9	11	PASS
C4	$m_D = m_p + m_n - E_D$	9.9	8	PASS
D1	Koide $K(e, \mu, \tau)$ from masses	10.3	6	PASS
D2	$a(\text{leptons})$ from K	10.2	6	PASS
D3	$M(\text{leptons}) = (\Sigma\sqrt{m})/3$	4.6	4	PASS
D4	Koide $K(u, c, t)$ from masses	10.5	3	PASS
D5	Koide $K(d, s, b)$ from masses	10.4	3	PASS
E1	$h \times \Delta \nu_{Cs}$ product consistency		exact	1
E2	c is exact integer 299792458	exact	15	PASS
E3	h exact SI 2019 value	exact	9	PASS
E4	e exact SI 2019 value	exact	10	PASS
E5	k_B exact SI 2019 value	exact	7	PASS
E6	N_A exact SI 2019 value	exact	9	PASS
Total: 32 tests, 32 PASS, 0 FAIL				

```

|| ALL TESTS PASS ||
||
|| DATA-3 DECLARATION: The DATA-2 database (107 original + ||
|| 16 Koide-derived = 123 entries) is internally consistent ||
|| to the precision of all source measurements. ||
||
|| DATA-2 is promoted to DATA-3. ||
|| DATA-2 is retired. ||
|| All future HOWL computation references DATA-
3. ||

```

```

└────────────────────────────────────────────────────────────────────────────────┘

```

```

=====
DATA-3 VERIFICATION COMPLETE
=====

```

[[HOWL-DATA-3-2026](https://github.com/ghowland/papers/tree/main/papers/DATA-3-2026)] Verified Integer Rational Database. Github: <https://github.com/ghowland/papers/tree/main/papers/DATA-3-2026>

[[HOWL-DATA-1-2026](https://github.com/ghowland/papers/tree/main/papers/DATA-1-2026)] Cross-Domain Precision Data Inventory. Github: <https://github.com/ghowland/papers/tree/main/papers/DATA-1-2026>

[[HOWL-DATA-2-2026](https://github.com/ghowland/papers/tree/main/papers/DATA/HOWL-DATA-2-2026)] Integer Rational Representations in Q335. Github: <https://github.com/ghowland/papers/tree/main/papers/DATA/HOWL-DATA-2-2026>