

# $K$ -theoretic Catalan functions

George H. Seelinger (joint with J. Blasiak and J. Morse)

CAGE

*ghs9ae@virginia.edu*

6 February 2020

- Schubert calculus
- Catalan functions: a new approach to old problems
- $K$ -theoretic Catalan functions

# Overview of Schubert Calculus Combinatorics

## Geometric problem

Find  $c_{\lambda\mu}^\nu = \#$  of points in intersection of subvarieties in a variety  $X$ .

# Overview of Schubert Calculus Combinatorics

## Geometric problem

Find  $c_{\lambda\mu}^\nu = \#$  of points in intersection of subvarieties in a variety  $X$ .



## Cohomology

Schubert basis  $\{\sigma_\lambda\}$  for  $H^*(X)$  with property  $\sigma_\lambda \cup \sigma_\mu = \sum_\nu c_{\lambda\mu}^\nu \sigma_\nu$

# Overview of Schubert Calculus Combinatorics

## Geometric problem

Find  $c_{\lambda\mu}^\nu = \#$  of points in intersection of subvarieties in a variety  $X$ .



## Cohomology

Schubert basis  $\{\sigma_\lambda\}$  for  $H^*(X)$  with property  $\sigma_\lambda \cup \sigma_\mu = \sum_\nu c_{\lambda\mu}^\nu \sigma_\nu$



## Representatives

Special basis of polynomials  $\{f_\lambda\}$  such that  $f_\lambda \cdot f_\mu = \sum_\nu c_{\lambda\mu}^\nu f_\nu$

# Overview of Schubert Calculus Combinatorics (cont.)

Combinatorial study of  $\{f_\lambda\}$  enlightens the geometry (and cohomology).

## Goal

Identify  $\{f_\lambda\}$  in explicit (simple) terms amenable to calculation and proofs.

## Geometric problem

Find  $c_{\lambda\mu}^\nu = \#$  of points in intersection of Schubert varieties  $\{X_\lambda\}_{\lambda \subseteq (n^m)}$  in variety  $X = \text{Gr}(m, n)$ .

# Classical Schubert Calculus

## Geometric problem

Find  $c_{\lambda\mu}^\nu = \#$  of points in intersection of Schubert varieties  $\{X_\lambda\}_{\lambda \subseteq (n^m)}$  in variety  $X = \text{Gr}(m, n)$ .



## Cohomology

Schubert basis  $\{\sigma_\lambda\}_{\lambda \subseteq (n^m)}$  for  $H^*(X)$  with property  $\sigma_\lambda \cup \sigma_\mu = \sum_\nu c_{\lambda\mu}^\nu \sigma_\nu$



# Classical Schubert Calculus

## Geometric problem

Find  $c_{\lambda\mu}^\nu = \#$  of points in intersection of Schubert varieties  $\{X_\lambda\}_{\lambda \subseteq (n^m)}$  in variety  $X = \text{Gr}(m, n)$ .



## Cohomology

Schubert basis  $\{\sigma_\lambda\}_{\lambda \subseteq (n^m)}$  for  $H^*(X)$  with property  $\sigma_\lambda \cup \sigma_\mu = \sum_\nu c_{\lambda\mu}^\nu \sigma_\nu$



## Representatives

Special basis of Schur polynomials  $\{s_\lambda\}$  such that  $s_\lambda \cdot s_\mu = \sum_\nu c_{\lambda\mu}^\nu s_\nu$  for Littlewood-Richardson coefficients  $c_{\lambda\mu}^\nu$ .

## Next Step: Flag Variety

- $X = Fl_n(\mathbb{C}) = \{V_0 \subseteq V_1 \subseteq \cdots \subseteq V_n \mid \dim V_i = i\}$

## Next Step: Flag Variety

- $X = Fl_n(\mathbb{C}) = \{V_0 \subseteq V_1 \subseteq \cdots \subseteq V_n \mid \dim V_i = i\}$
- Decomposes into Schubert varieties indexed by  $w \in S_n$ .

## Next Step: Flag Variety

- $X = Fl_n(\mathbb{C}) = \{V_0 \subseteq V_1 \subseteq \cdots \subseteq V_n \mid \dim V_i = i\}$
- Decomposes into Schubert varieties indexed by  $w \in S_n$ .
- $H^*(Fl_n(\mathbb{C}))$  supported by Schubert polynomials  $\mathfrak{S}_w \in \mathbb{Z}[x_1, \dots, x_n]$   
(Not necessarily symmetric!)

## Next Step: Flag Variety

- $X = Fl_n(\mathbb{C}) = \{V_0 \subseteq V_1 \subseteq \cdots \subseteq V_n \mid \dim V_i = i\}$
- Decomposes into Schubert varieties indexed by  $w \in S_n$ .
- $H^*(Fl_n(\mathbb{C}))$  supported by Schubert polynomials  $\mathfrak{S}_w \in \mathbb{Z}[x_1, \dots, x_n]$   
(Not necessarily symmetric!)

### Open Problem

Structure constants  $\mathfrak{S}_w \mathfrak{S}_u = c_{wu}^v \mathfrak{S}_v$  are combinatorially unknown.

There are many variations on classical Schubert calculus of the Grassmannian (Type A).

# Schubert Calculus Variations

There are many variations on classical Schubert calculus of the Grassmannian (Type A).

Theory	$f_\lambda$
(Co)homology of Grassmannian	Schur functions
(Co)homology of flag variety	Schubert polynomials
Quantum cohomology of flag variety	Quantum Schuberts
(Co)homology of Types BCD Grassmannian	Schur- $P$ and $Q$ functions
(Co)homology of affine Grassmannian	(dual) $k$ -Schur functions
$K$ -theory of Grassmannian	Grothendieck polynomials
$K$ -homology of affine Grassmannian	$K$ - $k$ -Schur functions

# Schubert Calculus Variations

There are many variations on classical Schubert calculus of the Grassmannian (Type A).

Theory	$f_\lambda$
(Co)homology of Grassmannian	Schur functions
(Co)homology of flag variety	Schubert polynomials
Quantum cohomology of flag variety	Quantum Schuberts
(Co)homology of Types BCD Grassmannian	Schur- $P$ and $Q$ functions
(Co)homology of affine Grassmannian	(dual) $k$ -Schur functions
$K$ -theory of Grassmannian	Grothendieck polynomials
$K$ -homology of affine Grassmannian	$K$ - $k$ -Schur functions

And many more!



- $QH^*(Fl_{k+1})$  quantum deformation of  $H^*(Fl_{k+1})$ .

# Peterson Isomorphism

- $QH^*(Fl_{k+1})$  quantum deformation of  $H^*(Fl_{k+1})$ .
- Supported by quantum Schubert polynomials  $\mathfrak{S}_w^Q$ .

# Peterson Isomorphism

- $QH^*(Fl_{k+1})$  quantum deformation of  $H^*(Fl_{k+1})$ .
- Supported by quantum Schubert polynomials  $\mathfrak{S}_w^Q$ .
- Peterson isomorphism

$$\Phi: QH^*(Fl_{k+1}) \rightarrow H_*(Gr_{SL_{k+1}})_{loc}$$

$$\mathfrak{S}_w^Q \mapsto \frac{s_\lambda^{(k)}}{\prod_{i \in Des(w)} \tau_i}$$

where  $s_\lambda^{(k)}$  is a  $k$ -Schur function.

# Peterson Isomorphism

- $QH^*(Fl_{k+1})$  quantum deformation of  $H^*(Fl_{k+1})$ .
- Supported by quantum Schubert polynomials  $\mathfrak{S}_w^Q$ .
- Peterson isomorphism

$$\Phi: QH^*(Fl_{k+1}) \rightarrow H_*(Gr_{SL_{k+1}})_{loc}$$

$$\mathfrak{S}_w^Q \mapsto \frac{s_\lambda^{(k)}}{\prod_{i \in Des(w)} \tau_i}$$

where  $s_\lambda^{(k)}$  is a  $k$ -Schur function.

## Upshot

Computations for (quantum) Schubert polynomials can be moved into symmetric functions.

# $k$ -Schur functions

- $s_{\lambda}^{(k)}$  for  $\lambda_1 \leq k$  a basis for  $\mathbb{Z}[s_1, s_2, \dots, s_k]$  (Lapointe et al., 2003).

# $k$ -Schur functions

- $s_{\lambda}^{(k)}$  for  $\lambda_1 \leq k$  a basis for  $\mathbb{Z}[s_1, s_2, \dots, s_k]$  (Lapointe et al., 2003).
- Schubert representatives for  $H_*(Gr_{SL_{k+1}})$  (Lam, 2008).

# $k$ -Schur functions

- $s_{\lambda}^{(k)}$  for  $\lambda_1 \leq k$  a basis for  $\mathbb{Z}[s_1, s_2, \dots, s_k]$  (Lapointe et al., 2003).
- Schubert representatives for  $H_*(Gr_{SL_{k+1}})$  (Lam, 2008).
- Has a tableaux formulation and Pieri rule:  $s_1 s_{\lambda}^{(k)} = \sum_{\mu} a_{\lambda\mu} s_{\mu}^{(k)}$

# $k$ -Schur functions

- $s_{\lambda}^{(k)}$  for  $\lambda_1 \leq k$  a basis for  $\mathbb{Z}[s_1, s_2, \dots, s_k]$  (Lapointe et al., 2003).
- Schubert representatives for  $H_*(Gr_{SL_{k+1}})$  (Lam, 2008).
- Has a tableaux formulation and Pieri rule:  $s_1 s_{\lambda}^{(k)} = \sum_{\mu} a_{\lambda\mu} s_{\mu}^{(k)}$
- $s_{\lambda}^{(k)} = s_{\lambda}$  as  $k \rightarrow \infty$ .



# $k$ -Schur functions

- $s_{\lambda}^{(k)}$  for  $\lambda_1 \leq k$  a basis for  $\mathbb{Z}[s_1, s_2, \dots, s_k]$  (Lapointe et al., 2003).
- Schubert representatives for  $H_*(Gr_{SL_{k+1}})$  (Lam, 2008).
- Has a tableaux formulation and Pieri rule:  $s_1 s_{\lambda}^{(k)} = \sum_{\mu} a_{\lambda\mu} s_{\mu}^{(k)}$
- $s_{\lambda}^{(k)} = s_{\lambda}$  as  $k \rightarrow \infty$ .
- Branching with positive coefficients (Lam et al., 2010):

$$s_{\lambda}^{(2)} = s_{\lambda}^{(3)} + s_{\mu}^{(3)} + s_{\nu}^{(3)}$$

# $k$ -Schur functions

- $s_{\lambda}^{(k)}$  for  $\lambda_1 \leq k$  a basis for  $\mathbb{Z}[s_1, s_2, \dots, s_k]$  (Lapointe et al., 2003).
- Schubert representatives for  $H_*(Gr_{SL_{k+1}})$  (Lam, 2008).
- Has a tableaux formulation and Pieri rule:  $s_1 s_{\lambda}^{(k)} = \sum_{\mu} a_{\lambda\mu} s_{\mu}^{(k)}$
- $s_{\lambda}^{(k)} = s_{\lambda}$  as  $k \rightarrow \infty$ .
- Branching with positive coefficients (Lam et al., 2010):

$$s_{\lambda}^{(2)} = s_{\lambda} + s_{\lambda} + s_{\lambda}$$

The diagram illustrates the branching rule for  $k=2$ . The left side shows  $s_{\lambda}^{(2)}$  for  $\lambda = (2, 2)$ , represented by a 2x2 grid of boxes. The right side shows the sum of three terms, each representing a  $s_{\mu}^{(3)}$  for  $\mu \vdash (2, 2)$ . The first term is  $s_{(2, 2)}^{(3)}$ , represented by a 2x2 grid of boxes. The second and third terms are  $s_{(2, 1, 1)}^{(3)}$ , represented by a 2x1 grid of boxes with a single box to the right of the bottom box. The terms are grouped by curly braces and labeled  $s_{\lambda}^{(3)}$  below them.

- Has geometric interpretation.

# $k$ -Schur functions

- $s_{\lambda}^{(k)}$  for  $\lambda_1 \leq k$  a basis for  $\mathbb{Z}[s_1, s_2, \dots, s_k]$  (Lapointe et al., 2003).
- Schubert representatives for  $H_*(Gr_{SL_{k+1}})$  (Lam, 2008).
- Has a tableaux formulation and Pieri rule:  $s_1^r s_{\lambda}^{(k)} = \sum_{\mu} a_{\lambda\mu} s_{\mu}^{(k)}$
- $s_{\lambda}^{(k)} = s_{\lambda}$  as  $k \rightarrow \infty$ .
- Branching with positive coefficients (Lam et al., 2010):

$$s_{\lambda}^{(2)} = s_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}} + s_{\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \end{array}} + s_{\begin{array}{|c|c|c|c|} \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \end{array}}$$

$\underbrace{\hspace{10em}}_{s_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}}^{(3)}} \quad \underbrace{\hspace{10em}}_{s_{\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \end{array}}^{(3)}}$

- Has geometric interpretation.
- No combinatorial interpretation of branching coefficients.

# $k$ -Schur functions

- $s_{\lambda}^{(k)}$  for  $\lambda_1 \leq k$  a basis for  $\mathbb{Z}[s_1, s_2, \dots, s_k]$  (Lapointe et al., 2003).
- Schubert representatives for  $H_*(Gr_{SL_{k+1}})$  (Lam, 2008).
- Has a tableaux formulation and Pieri rule:  $s_1^r s_{\lambda}^{(k)} = \sum_{\mu} a_{\lambda\mu} s_{\mu}^{(k)}$
- $s_{\lambda}^{(k)} = s_{\lambda}$  as  $k \rightarrow \infty$ .
- Branching with positive coefficients (Lam et al., 2010):

$$s_{\lambda}^{(2)} = s_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}} + s_{\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \end{array}} + s_{\begin{array}{|c|c|c|c|} \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \end{array}}$$

$\underbrace{\hspace{10em}}_{s_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}}^{(3)}} \quad \underbrace{\hspace{10em}}_{s_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}}^{(3)}}$

- Has geometric interpretation.
- No combinatorial interpretation of branching coefficients.
- Definition with  $t$  important for Macdonald polynomials.

# $k$ -Schur functions

- $s_{\lambda}^{(k)}$  for  $\lambda_1 \leq k$  a basis for  $\mathbb{Z}[s_1, s_2, \dots, s_k]$  (Lapointe et al., 2003).
- Schubert representatives for  $H_*(Gr_{SL_{k+1}})$  (Lam, 2008).
- Has a tableaux formulation and Pieri rule:  $s_1^r s_{\lambda}^{(k)} = \sum_{\mu} a_{\lambda\mu} s_{\mu}^{(k)}$
- $s_{\lambda}^{(k)} = s_{\lambda}$  as  $k \rightarrow \infty$ .
- Branching with positive coefficients (Lam et al., 2010):

$$s_{\lambda}^{(2)} = s_{\lambda}^{(3)} + s_{\mu}^{(3)} + s_{\nu}^{(3)}$$

The diagram shows the branching of the Schur function  $s_{\lambda}^{(2)}$  into three Schur functions of degree 3. On the left,  $s_{\lambda}^{(2)}$  is represented by a 2x2 square. On the right, it is equal to the sum of three terms:  $s_{\lambda}^{(3)}$  (a 3x2 rectangle),  $s_{\mu}^{(3)}$  (a 2x2 square with an additional cell to the right), and  $s_{\nu}^{(3)}$  (a horizontal row of three cells). Brackets below the terms on the right group them under  $s_{(3)}^{(3)}$  and  $s_{(2,1)}^{(3)}$ .

- Has geometric interpretation.
- No combinatorial interpretation of branching coefficients.
- Definition with  $t$  important for Macdonald polynomials.
- Many definitions. A new one makes proofs easier!

- Schubert calculus
- **Catalan functions: a new approach to old problems**
- $K$ -theoretic Catalan functions

# Raising Operators on Symmetric Functions

- Raising operators  $R_{i,j}$  act on diagrams

$$R_{1,3} \left( \begin{array}{|c|c|c|c|} \hline \text{red} & & & \\ \hline & & & \\ \hline & & & \\ \hline \end{array} \right) = \begin{array}{|c|c|c|c|} \hline & & & \\ \hline & & & \text{red} \\ \hline & & & \\ \hline \end{array} \quad R_{2,3} \left( \begin{array}{|c|} \hline \text{red} \\ \hline \\ \hline \\ \hline \end{array} \right) = \begin{array}{|c|c|} \hline & \text{red} \\ \hline & \\ \hline \end{array}$$

# Raising Operators on Symmetric Functions

- Raising operators  $R_{i,j}$  act on diagrams

$$R_{1,3} \left( \begin{array}{|c|c|c|c|} \hline \text{red} & & & \\ \hline & & & \\ \hline & & & \\ \hline \end{array} \right) = \begin{array}{|c|c|c|c|} \hline & & & \\ \hline & & & \text{red} \\ \hline & & & \\ \hline \end{array} \quad R_{2,3} \left( \begin{array}{|c|} \hline \text{red} \\ \hline \\ \hline \\ \hline \end{array} \right) = \begin{array}{|c|c|} \hline & \text{red} \\ \hline & \\ \hline & \\ \hline \end{array}$$

- Extend action to a symmetric function  $f_\lambda$  by  $R_{i,j}(f_\lambda) = f_{\lambda + \epsilon_i - \epsilon_j}$ .



# Raising Operators on Symmetric Functions

- Raising operators  $R_{i,j}$  act on diagrams

$$R_{1,3} \left( \begin{array}{|c|c|c|} \hline \text{red} & & \\ \hline & & \\ \hline & & \\ \hline \end{array} \right) = \begin{array}{|c|c|c|c|} \hline & & & \\ \hline & & & \text{red} \\ \hline & & & \\ \hline \end{array} \quad R_{2,3} \left( \begin{array}{|c|} \hline \text{red} \\ \hline & \\ \hline & \\ \hline \end{array} \right) = \begin{array}{|c|c|} \hline & \text{red} \\ \hline & \\ \hline \end{array}$$

- Extend action to a symmetric function  $f_\lambda$  by  $R_{i,j}(f_\lambda) = f_{\lambda + \epsilon_i - \epsilon_j}$ .
- For  $h_\lambda = s_{\lambda_1} \cdots s_{\lambda_r}$ , we have the *Jacobi-Trudi identity*

$$s_\lambda = \prod_{i < j} (1 - R_{ij}) h_\lambda$$

# Raising Operators on Symmetric Functions

- Raising operators  $R_{i,j}$  act on diagrams

$$R_{1,3} \left( \begin{array}{|c|c|c|c|} \hline \text{red} & & & \\ \hline & & & \\ \hline & & & \\ \hline \end{array} \right) = \begin{array}{|c|c|c|c|} \hline & & & \\ \hline & & & \text{red} \\ \hline & & & \\ \hline \end{array} \quad R_{2,3} \left( \begin{array}{|c|} \hline \text{red} \\ \hline \\ \hline \\ \hline \end{array} \right) = \begin{array}{|c|c|} \hline & \text{red} \\ \hline & \\ \hline & \\ \hline \end{array}$$

- Extend action to a symmetric function  $f_\lambda$  by  $R_{i,j}(f_\lambda) = f_{\lambda+\epsilon_i-\epsilon_j}$ .
- For  $h_\lambda = s_{\lambda_1} \cdots s_{\lambda_r}$ , we have the *Jacobi-Trudi identity*

$$s_\lambda = \prod_{i < j} (1 - R_{ij}) h_\lambda$$

$$s_{22} = (1 - R_{12}) h_{22} = h_{22} - h_{31}$$

$$s_{211} = (1 - R_{12})(1 - R_{23})(1 - R_{13}) h_{211}$$

$$= h_{211} - h_{301} - h_{220} - \text{red } h_{310} + \text{red } h_{310} + \underbrace{h_{32-1}}_{=0} + h_{400} - \underbrace{h_{41-1}}_{=0}$$

some terms cancel

# Raising Operators on Symmetric Functions

Gives definition for Schur function indexed by any integer vector  $\alpha \in \mathbb{Z}^\ell$ .

# Raising Operators on Symmetric Functions

Gives definition for Schur function indexed by any integer vector  $\alpha \in \mathbb{Z}^\ell$ .

Straightening:

$$s_\alpha = \prod_{i < j} (1 - R_{ij}) h_\alpha = \begin{cases} \pm s_\lambda & \text{for a partition } \lambda \\ 0 & \end{cases}$$

# Raising Operators on Symmetric Functions

Gives definition for Schur function indexed by any integer vector  $\alpha \in \mathbb{Z}^\ell$ .

Straightening:

$$s_\alpha = \prod_{i < j} (1 - R_{ij}) h_\alpha = \begin{cases} \pm s_\lambda & \text{for a partition } \lambda \\ 0 & \end{cases}$$


For  $\langle s_{1^r}^\perp s_\lambda, s_\mu \rangle = \langle s_\lambda, s_{1^r} s_\mu \rangle$ ,

$$s_{1^r}^\perp s_\lambda = \sum_{S \subseteq [1, \ell], |S|=r} s_{\lambda - \epsilon_S}$$

$$s_{1^2}^\perp s_{333} = s_{322} + s_{232} + s_{223}$$

# Root Ideals

A root ideal  $\Psi$  of type  $A_{\ell-1}$  positive roots: given by Dyck path (lattice path above diagonal).




$\Psi =$  Roots above Dyck path

$\Delta_{\ell}^{+} \setminus \Psi =$  Non-roots below

# Root Ideals

A root ideal  $\Psi$  of type  $A_{\ell-1}$  positive roots: given by Dyck path (lattice path above diagonal).



$\Psi =$  Roots above Dyck path  
 $\Delta_{\ell}^{+} \setminus \Psi =$  Non-roots below


Catalan Function (Chen, 2010; Panyushev, 2010; Blasiak et al., 2019)

For  $\Psi$  and  $\gamma \in \mathbb{Z}^{\ell}$

$$H(\Psi; \gamma)(x) = \prod_{(i,j) \in \Delta_{\ell}^{+} \setminus \Psi} (1 - R_{ij}) h_{\gamma}(x)$$

# Root Ideals

A root ideal  $\Psi$  of type  $A_{\ell-1}$  positive roots: given by Dyck path (lattice path above diagonal).



$\Psi$  = Roots above Dyck path  
 $\Delta_{\ell}^{+} \setminus \Psi$  = Non-roots below

Catalan Function (Chen, 2010; Panyushev, 2010; Blasiak et al., 2019)

For  $\Psi$  and  $\gamma \in \mathbb{Z}^{\ell}$


$$H(\Psi; \gamma)(x) = \prod_{(i,j) \in \Delta_{\ell}^{+} \setminus \Psi} (1 - R_{ij}) h_{\gamma}(x)$$

- $\Psi = \emptyset \implies H(\emptyset; \gamma) = s_{\gamma}$



# Root Ideals

A root ideal  $\Psi$  of type  $A_{\ell-1}$  positive roots: given by Dyck path (lattice path above diagonal).



$\Psi$  = Roots above Dyck path  
 $\Delta_{\ell}^{+} \setminus \Psi$  = Non-roots below

Catalan Function (Chen, 2010; Panyushev, 2010; Blasiak et al., 2019)

For  $\Psi$  and  $\gamma \in \mathbb{Z}^{\ell}$

$$H(\Psi; \gamma)(x) = \prod_{(i,j) \in \Delta_{\ell}^{+} \setminus \Psi} (1 - R_{ij}) h_{\gamma}(x)$$

- $\Psi = \emptyset \implies H(\emptyset; \gamma) = s_{\gamma}$
- $\Psi = \text{all roots} \implies H(\Psi; \gamma) = h_{\gamma}$

## $k$ -Schur root ideal for $\lambda$

$$\begin{aligned}\psi = \Delta^k(\lambda) &= \{(i, j) : j > k - \lambda_i\} \\ &= \text{root ideal with } k - \lambda_i \text{ non-roots in row } i\end{aligned}$$

# Catalan functions

## $k$ -Schur root ideal for $\lambda$

$$\begin{aligned}\psi = \Delta^k(\lambda) &= \{(i, j) : j > k - \lambda_i\} \\ &= \text{root ideal with } k - \lambda_i \text{ non-roots in row } i\end{aligned}$$

$$\Delta^4(3, 3, 2, 2, 1, 1) =$$

3					
	3				
		2			
			2		
				1	
					1

$\leftarrow$  row  $i$  has  $4 - \lambda_i$  non-roots

# Catalan functions

## $k$ -Schur root ideal for $\lambda$

$$\begin{aligned}\psi = \Delta^k(\lambda) &= \{(i, j) : j > k - \lambda_i\} \\ &= \text{root ideal with } k - \lambda_i \text{ non-roots in row } i\end{aligned}$$

$$\Delta^4(3, 3, 2, 2, 1, 1) =$$

3						
	3					
		2				
			2			
				1		
					1	

$\leftarrow$  row  $i$  has  $4 - \lambda_i$  non-roots

$k$ -Schur is a Catalan function (Blasiak et al., 2019).

For partition  $\lambda$  with  $\lambda_1 \leq k$ ,

$$s_{\lambda}^{(k)} = H(\Delta^k(\lambda); \lambda).$$

# Key ingredient of branching proof

Dual vertical Pieri rule:  $s_{1^r}^\perp s_\lambda^{(k)} = \sum_\mu a_{\lambda\mu} s_\mu^{(k)}$  for  $\langle s_{1^r}^\perp f, g \rangle = \langle f, s_{1^r} g \rangle$ .

# Key ingredient of branching proof

Dual vertical Pieri rule:  $s_{1^r}^\perp s_\lambda^{(k)} = \sum_\mu a_{\lambda\mu} s_\mu^{(k)}$  for  $\langle s_{1^r}^\perp f, g \rangle = \langle f, s_{1^r} g \rangle$ .

Shift Invariance (Blasiak et al., 2019)

For partition  $\lambda$  of length  $\ell$  with  $\lambda_1 \leq k$ ,

$$s_{1^\ell}^\perp s_{\lambda+1^\ell}^{(k+1)} = s_\lambda^{(k)}.$$

# Key ingredient of branching proof

Dual vertical Pieri rule:  $s_{1^r}^\perp s_\lambda^{(k)} = \sum_\mu a_{\lambda\mu} s_\mu^{(k)}$  for  $\langle s_{1^r}^\perp f, g \rangle = \langle f, s_{1^r} g \rangle$ .

## Shift Invariance (Blasiak et al., 2019)

For partition  $\lambda$  of length  $\ell$  with  $\lambda_1 \leq k$ ,

$$s_{1^\ell}^\perp s_{\lambda+1^\ell}^{(k+1)} = s_\lambda^{(k)}.$$

Proof:  $k - \lambda_i = (k + 1) - (\lambda_i + 1)$

# Key ingredient of branching proof

Dual vertical Pieri rule:  $s_{1^r}^\perp s_\lambda^{(k)} = \sum_\mu a_{\lambda\mu} s_\mu^{(k)}$  for  $\langle s_{1^r}^\perp f, g \rangle = \langle f, s_{1^r} g \rangle$ .

## Shift Invariance (Blasiak et al., 2019)

For partition  $\lambda$  of length  $\ell$  with  $\lambda_1 \leq k$ ,

$$s_{1^\ell}^\perp s_{\lambda+1^\ell}^{(k+1)} = s_\lambda^{(k)}.$$

Proof:  $k - \lambda_i = (k + 1) - (\lambda_i + 1)$

$$\Delta^4(3, 3, 2, 2, 1, 1) =$$

3					
	3				
		2			
			2		
				1	
					1

$$\Delta^5(4, 4, 3, 3, 2, 2) =$$

4					
	4				
		3			
			3		
				2	
					2



# Key ingredient of branching proof


Dual vertical Pieri rule:  $s_{1^r}^\perp s_\lambda^{(k)} = \sum_\mu a_{\lambda\mu} s_\mu^{(k)}$  for  $\langle s_{1^r}^\perp f, g \rangle = \langle f, s_{1^r} g \rangle$ .


## Shift Invariance (Blasiak et al., 2019)

For partition  $\lambda$  of length  $\ell$  with  $\lambda_1 \leq k$ ,

$$s_{1^\ell}^\perp s_{\lambda+1^\ell}^{(k+1)} = s_\lambda^{(k)}.$$

Proof:  $k - \lambda_i = (k + 1) - (\lambda_i + 1)$

$$\Delta^4(3, 3, 2, 2, 1, 1) =$$


$$\Delta^5(4, 4, 3, 3, 2, 2) =$$


Pieri:

$$s_{1^\ell}^\perp s_{\lambda+1^\ell}^{(k+1)} = \sum_\mu a_{\lambda+1^\ell, \mu} s_\mu^{(k+1)}$$

# Key ingredient of branching proof


Dual vertical Pieri rule:  $s_{1^r}^\perp s_\lambda^{(k)} = \sum_\mu a_{\lambda\mu} s_\mu^{(k)}$  for  $\langle s_{1^r}^\perp f, g \rangle = \langle f, s_{1^r} g \rangle$ .


## Shift Invariance (Blasiak et al., 2019)

For partition  $\lambda$  of length  $\ell$  with  $\lambda_1 \leq k$ ,

$$s_{1^\ell}^\perp s_{\lambda+1^\ell}^{(k+1)} = s_\lambda^{(k)}.$$

Proof:  $k - \lambda_i = (k + 1) - (\lambda_i + 1)$

$$\Delta^4(3, 3, 2, 2, 1, 1) =$$


$$\Delta^5(4, 4, 3, 3, 2, 2) =$$


Branching is a special case of Pieri:

$$s_\lambda^{(k)} = s_{1^\ell}^\perp s_{\lambda+1^\ell}^{(k+1)} = \sum_\mu a_{\lambda+1^\ell, \mu} s_\mu^{(k+1)}$$

- Schubert calculus
- Catalan functions: a new approach to old problems
- ***K*-theoretic Catalan functions**

# Dual Grothendieck polynomials

- Inhomogeneous basis:  $g_\lambda = s_\lambda + \text{lower degree terms}$ .

# Dual Grothendieck polynomials

- Inhomogeneous basis:  $g_\lambda = s_\lambda +$  lower degree terms.
- Satisfies Pieri rule on “set-valued strips”

# Dual Grothendieck polynomials

- Inhomogeneous basis:  $g_\lambda = s_\lambda + \text{lower degree terms}$ .
- Satisfies Pieri rule on “set-valued strips”

$$g_{1^2}g_{3,2} = g_{43} + g_{421} + g_{331} - g_{42} - g_{33} - 2g_{321} + g_{31}$$



# Dual Grothendieck polynomials

- Inhomogeneous basis:  $g_\lambda = s_\lambda + \text{lower degree terms}$ .
- Satisfies Pieri rule on “set-valued strips”

$$g_{1^2}g_{3,2} = g_{43} + g_{421} + g_{331} - g_{42} - g_{33} - 2g_{321} + g_{31}$$



- $g_\lambda = \prod_{i < j} (1 - R_{ij}) k_\lambda$  for  $k_\lambda$  and inhomogeneous analogue of  $h_\lambda$ .

# Dual Grothendieck polynomials

- Inhomogeneous basis:  $g_\lambda = s_\lambda + \text{lower degree terms}$ .
- Satisfies Pieri rule on “set-valued strips”

$$g_{1^2}g_{3,2} = g_{43} + g_{421} + g_{331} - g_{42} - g_{33} - 2g_{321} + g_{31}$$



- $g_\lambda = \prod_{i < j} (1 - R_{ij}) k_\lambda$  for  $k_\lambda$  and inhomogeneous analogue of  $h_\lambda$ .
- Dual to Grothendieck polynomials  $G_\lambda$ : Schubert representatives for  $K^*(Gr(m, n))$



# $K$ - $k$ -Schur functions

- Inhomogeneous basis:  $g_{\lambda}^{(k)} = s_{\lambda}^{(k)} + \text{lower degree terms}$

# $K$ - $k$ -Schur functions

- Inhomogeneous basis:  $g_{\lambda}^{(k)} = s_{\lambda}^{(k)} + \text{lower degree terms}$
- Satisfies Pieri rule on “affine set-valued strips”

# $K$ - $k$ -Schur functions

- Inhomogeneous basis:  $g_{\lambda}^{(k)} = s_{\lambda}^{(k)} + \text{lower degree terms}$
- Satisfies Pieri rule on “affine set-valued strips”

$$g_1 g_{211}^{(2)} = g_{2111}^{(2)} - 2g_{211}^{(2)}$$

2-bounded partitions  $\leftrightarrow$  3-cores

The diagram illustrates the Pieri rule for  $K$ -Schur functions. It shows the product of a single box (1) and a 3-core (211) resulting in the difference of two 3-cores (2111 and 211). The boxes are colored red, blue, and black, and the diagram shows the addition and subtraction of these boxes to form the result.

# $K$ - $k$ -Schur functions

- Inhomogeneous basis:  $g_{\lambda}^{(k)} = s_{\lambda}^{(k)} + \text{lower degree terms}$
- Satisfies Pieri rule on “affine set-valued strips”

$$g_1 g_{211}^{(2)} = g_{2111}^{(2)} - 2g_{211}^{(2)}$$

2-bounded partitions  $\leftrightarrow$  3-cores

- Conjecture:  $g_{\lambda}^{(k)}$  have positive branching into  $g_{\mu}^{(k+1)}$  (Lam et al., 2010; Morse, 2011).

# $K$ - $k$ -Schur functions

- Inhomogeneous basis:  $g_{\lambda}^{(k)} = s_{\lambda}^{(k)} + \text{lower degree terms}$
- Satisfies Pieri rule on “affine set-valued strips”

$$g_1 g_{211}^{(2)} = g_{2111}^{(2)} - 2g_{211}^{(2)}$$

2-bounded partitions  $\leftrightarrow$  3-cores

The diagram illustrates the Pieri rule for  $K$ -Schur functions. It shows the product of  $g_1$  and  $g_{211}^{(2)}$  as a difference of two 2-bounded partitions (3-cores). The first partition is  $g_{2111}^{(2)}$  and the second is  $2g_{211}^{(2)}$ . The diagram uses colored dots (red, blue, black) to represent the partitions and boxes to highlight the relevant cells.

- Conjecture:  $g_{\lambda}^{(k)}$  have positive branching into  $g_{\mu}^{(k+1)}$  (Lam et al., 2010; Morse, 2011).

## Problem

No direct formula for  $g_{\lambda}^{(k)}$

# An Extra Ingredient: Lowering Operators

Lowering Operators  $L_j(f_\lambda) = f_{\lambda - \epsilon_j}$

$$L_3 \left( \begin{array}{|c|c|c|c|} \hline \text{red} & & & \\ \hline & & & \\ \hline & & & \\ \hline \end{array} \right) = \begin{array}{|c|c|c|} \hline & & \\ \hline & & \\ \hline \end{array}, \quad L_1 \left( \begin{array}{|c|c|c|c|} \hline & & & \\ \hline & & & \\ \hline & & & \\ \hline & & \text{red} & \\ \hline \end{array} \right) = \begin{array}{|c|c|} \hline & \\ \hline & \\ \hline & \\ \hline \end{array}$$

## $K$ -theoretic Catalan function

Let  $\Psi, \mathcal{L} \subseteq \Delta_\ell^+$  be order ideals of positive roots and  $\gamma \in \mathbb{Z}^\ell$ , then

$$K(\Psi; \mathcal{L}; \gamma) := \prod_{(i,j) \in \mathcal{L}} (1 - L_j) \prod_{(i,j) \in \Delta_\ell^+ \setminus \Psi} (1 - R_{ij}) k_\gamma$$

# Affine $K$ -Theory Representatives with Raising Operators

## $K$ -theoretic Catalan function

Let  $\Psi, \mathcal{L} \subseteq \Delta_\ell^+$  be order ideals of positive roots and  $\gamma \in \mathbb{Z}^\ell$ , then

$$K(\Psi; \mathcal{L}; \gamma) := \prod_{(i,j) \in \mathcal{L}} (1 - L_j) \prod_{(i,j) \in \Delta_\ell^+ \setminus \Psi} (1 - R_{ij}) k_\gamma$$

## Example

non-roots of  $\Psi$ , roots of  $\mathcal{L}$

	(12)	(13)	(14)	(15)
		(23)	(24)	(25)
			(34)	(35)
				(45)

$$\begin{aligned} K(\Psi; \mathcal{L}; 54332) \\ = (1 - L_4)^2 (1 - L_5)^2 (1 - R_{12}) (1 - R_{34}) (1 - R_{45}) k_{54332} \end{aligned}$$



Answer (Blasiak-Morse-S., 2020)

## Answer (Blasiak-Morse-S., 2020)


For  $K$ -homology of affine Grassmannian,  $g_{\lambda}^{(k)} = K(\Delta^k(\lambda); \Delta^{k+1}(\lambda); \lambda)$  since this family satisfies the Pieri rule.

# Affine $K$ -Theory Representatives with Raising Operators

Answer (Blasiak-Morse-S., 2020)

For  $K$ -homology of affine Grassmannian,  $g_{\lambda}^{(k)} = K(\Delta^k(\lambda); \Delta^{k+1}(\lambda); \lambda)$  since this family satisfies the Pieri rule.

Example



$$\Delta_9^+ / \Delta^4(332111111), \Delta^5(332111111)$$

## Theorem (Blasiak-Morse-S., 2020)

## Theorem (Blasiak-Morse-S., 2020)

The  $g_{\lambda}^{(k)}$  are “shift invariant”, i.e. for  $\ell = \ell(\lambda)$

$$G_{1^{\ell}}^{\perp} g_{\lambda+1^{\ell}}^{(k+1)} = g_{\lambda}^{(k)}$$

## Theorem (Blasiak-Morse-S., 2020)

The  $g_\lambda^{(k)}$  are “shift invariant”, i.e. for  $\ell = \ell(\lambda)$

$$G_{1^\ell}^\perp g_{\lambda+1^\ell}^{(k+1)} = g_\lambda^{(k)}$$

## Theorem (Blasiak-Morse-S., 2020)

The branching coefficients in

$$g_\lambda^{(k)} = \sum_{\mu} a_{\lambda\mu} g_\mu^{(k+1)}$$

satisfy  $(-1)^{|\lambda|-|\mu|} a_{\lambda\mu} \in \mathbb{Z}_{\geq 0}$ .

# $K$ -theoretic Peterson isomorphism

$$\Phi: QK^*(Fl_{k+1}) \rightarrow K_*(Gr_{SL_{k+1}})_{loc}$$

# $K$ -theoretic Peterson isomorphism

$$\Phi: QK^*(Fl_{k+1}) \rightarrow K_*(Gr_{SL_{k+1}})_{loc}$$

Conjecture (Ikeda et al., 2018)

For  $w \in S_{k+1}$  and  $\mathfrak{G}_w^Q$  a “quantum Grothendieck polynomial”,

$$\Phi(\mathfrak{G}_w^Q) = \frac{\tilde{g}_w}{\prod_{i \in Des(w)} \tau_i}$$



# $K$ -theoretic Peterson isomorphism

$$\Phi: QK^*(Fl_{k+1}) \rightarrow K_*(Gr_{SL_{k+1}})_{loc}$$

Conjecture (Ikeda et al., 2018)

For  $w \in S_{k+1}$  and  $\mathfrak{G}_w^Q$  a “quantum Grothendieck polynomial”,

$$\Phi(\mathfrak{G}_w^Q) = \frac{\tilde{g}_w}{\prod_{i \in Des(w)} \tau_i}$$

satisfies  $\tilde{g}_w = g_\lambda^{(k)} + \sum_\mu a_{\lambda\mu} g_\mu^{(k)}$  such that  $(-1)^{|\lambda|-|\mu|} a_{\lambda\mu} \in \mathbb{Z}_{\geq 0}$ .

# $K$ -theoretic Peterson isomorphism

$$\Phi: QK^*(Fl_{k+1}) \rightarrow K_*(Gr_{SL_{k+1}})_{loc}$$

## Conjecture (Ikeda et al., 2018)

For  $w \in S_{k+1}$  and  $\mathfrak{G}_w^Q$  a “quantum Grothendieck polynomial”,

$$\Phi(\mathfrak{G}_w^Q) = \frac{\tilde{g}_w}{\prod_{i \in Des(w)} \tau_i}$$

satisfies  $\tilde{g}_w = g_\lambda^{(k)} + \sum_\mu a_{\lambda\mu} g_\mu^{(k)}$  such that  $(-1)^{|\lambda|-|\mu|} a_{\lambda\mu} \in \mathbb{Z}_{\geq 0}$ .

## Theorem (Blasiak-Morse-S., 2020)

*If  $\lambda \subseteq (d^{k+1-d})$  for some  $1 \leq d \leq k$ , then  $g_\lambda^{(k)} = g_\lambda$ . Thus, conjecture is true for  $w$  a Grassmannian permutation (i.e.  $w$  has only one descent).*

# $K$ -theoretic Peterson isomorphism

$$\Phi: QK^*(Fl_{k+1}) \rightarrow K_*(Gr_{SL_{k+1}})_{loc}$$

## Conjecture (Ikeda et al., 2018)

For  $w \in S_{k+1}$  and  $\mathfrak{G}_w^Q$  a “quantum Grothendieck polynomial”,

$$\Phi(\mathfrak{G}_w^Q) = \frac{\tilde{g}_w}{\prod_{i \in Des(w)} \tau_i}$$

satisfies  $\tilde{g}_w = g_\lambda^{(k)} + \sum_\mu a_{\lambda\mu} g_\mu^{(k)}$  such that  $(-1)^{|\lambda|-|\mu|} a_{\lambda\mu} \in \mathbb{Z}_{\geq 0}$ .

## Theorem (Blasiak-Morse-S., 2020)

*If  $\lambda \subseteq (d^{k+1-d})$  for some  $1 \leq d \leq k$ , then  $g_\lambda^{(k)} = g_\lambda$ . Thus, conjecture is true for  $w$  a Grassmannian permutation (i.e.  $w$  has only one descent).*

## Conjecture (Blasiak-Morse-S., 2020)

$$\tilde{g}_w = K(\Delta^k(\lambda); \Delta^k(\lambda); \lambda)$$

For  $G_{\lambda}^{(k)}$  an affine Grothendieck polynomial (dual to  $g_{\lambda}^{(k)}$ ),

For  $G_\lambda^{(k)}$  an affine Grothendieck polynomial (dual to  $g_\lambda^{(k)}$ ),

- 1 Combinatorially describe dual Pieri rule:

$$G_{1^r}^\perp g_\lambda^{(k)} = \sum_\mu ?? g_\mu^{(k)} \iff G_{1^r} G_\mu^{(k)} = \sum_\lambda ?? G_\lambda^{(k)}, \quad 1 \leq r \leq k.$$

For  $G_\lambda^{(k)}$  an affine Grothendieck polynomial (dual to  $g_\lambda^{(k)}$ ),

- 1 Combinatorially describe dual Pieri rule:

$$G_{1^r}^\perp g_\lambda^{(k)} = \sum_\mu g_\mu^{(k)} \iff G_{1^r} G_\mu^{(k)} = \sum_\lambda G_\lambda^{(k)}, \quad 1 \leq r \leq k.$$

- 2 Combinatorially describe branching coefficients:  $g_\lambda^{(k)} = \sum_\mu g_\mu^{(k+1)}$ .

For  $G_\lambda^{(k)}$  an affine Grothendieck polynomial (dual to  $g_\lambda^{(k)}$ ),

- 1 Combinatorially describe dual Pieri rule:

$$G_{1^r}^\perp g_\lambda^{(k)} = \sum_\mu g_\mu^{(k)} \iff G_{1^r} G_\mu^{(k)} = \sum_\lambda G_\lambda^{(k)}, \quad 1 \leq r \leq k.$$

- 2 Combinatorially describe branching coefficients:  $g_\lambda^{(k)} = \sum_\mu g_\mu^{(k+1)}$ .

- 3 Combinatorially describe  $g_\lambda^{(k)} = \sum_\mu s_\mu^{(k)}$ .

For  $G_\lambda^{(k)}$  an affine Grothendieck polynomial (dual to  $g_\lambda^{(k)}$ ),

- 1 Combinatorially describe dual Pieri rule:

$$G_{1^r}^\perp g_\lambda^{(k)} = \sum_\mu g_\mu^{(k)} \iff G_{1^r} G_\mu^{(k)} = \sum_\lambda G_\lambda^{(k)}, \quad 1 \leq r \leq k.$$

- 2 Combinatorially describe branching coefficients:  $g_\lambda^{(k)} = \sum_\mu g_\mu^{(k+1)}$ .

- 3 Combinatorially describe  $g_\lambda^{(k)} = \sum_\mu s_\mu^{(k)}$ .

- 4 Prove the image of  $\mathfrak{G}_w^Q$  under  $K$ -theoretic Peterson isomorphism for all  $w \in S_{k+1}$ .



## Thank you!

- Anderson, David, Linda Chen, and Hsian-Hua Tseng. 2017. *On the quantum  $K$ -ring of the flag manifold*, preprint. arXiv: 1711.08414.
- Blasiak, Jonah, Jennifer Morse, Anna Pun, and Daniel Summers. 2019. *Catalan Functions and  $k$ -Schur Positivity*, J. Amer. Math. Soc. **32**, no. 4, 921–963.
- Chen, Li-Chung. 2010. *Skew-linked partitions and a representation theoretic model for  $k$ -Schur functions*, Ph.D. thesis.
- Ikedo, Takeshi, Shinsuke Iwao, and Toshiaki Maeno. 2018. *Peterson Isomorphism in  $K$ -theory and Relativistic Toda Lattice*, preprint. arXiv: 1703.08664.
- Lam, Thomas. 2008. *Schubert polynomials for the affine Grassmannian*, J. Amer. Math. Soc. **21**, no. 1, 259–281.
- Lam, Thomas, Luc Lapointe, Jennifer Morse, and Mark Shimozono. 2010. *Affine insertion and Pieri rules for the affine Grassmannian*, Mem. Amer. Math. Soc. **208**, no. 977.
- Lam, Thomas, Anne Schilling, and Mark Shimozono. 2010.  *$K$ -theory Schubert calculus of the affine Grassmannian*, Compositio Math. **146**, 811–852.
- Lapointe, Luc, Alain Lascoux, and Jennifer Morse. 2003. *Tableau atoms and a new Macdonald positivity conjecture*, Duke Mathematical Journal **116**, no. 1, 103–146.
- Morse, Jennifer. 2011. *Combinatorics of the  $K$ -theory of affine Grassmannians*, Advances in Mathematics.
- Panyushev, Dmitri I. 2010. *Generalised Kostka-Foulkes polynomials and cohomology of line bundles on homogeneous vector bundles*, Selecta Math. (N.S.) **16**, no. 2, 315–342.