

Catalanimals: a walk through the zoo of shuffle theorems

George H. Seelinger
(University of Michigan)

ghseeli@umich.edu

joint with Jonah Blasiak, Mark Haiman, Jennifer Morse, and Anna Pun


University of Pennsylvania Mathematics Colloquium

December 10th, 2024


Fundamental Object: Symmetric polynomials

Symmetric
polynomials


Fundamental Object: Symmetric polynomials



Fundamental Object: Symmetric polynomials



Fundamental Object: Symmetric polynomials



Symmetric polynomials

Symmetric polynomials

$f(z_1, \dots, z_n)$ symmetric if swapping variables does not change f .

Symmetric polynomials

$f(z_1, \dots, z_n)$ symmetric if swapping variables does not change f .

- $f(z_1, z_2, z_3) = z_1 + z_2 + z_3$

Symmetric polynomials

$f(z_1, \dots, z_n)$ symmetric if swapping variables does not change f .

- $f(z_1, z_2, z_3) = z_1 + z_2 + z_3$
- $f(z_1, z_2, z_3) = z_1z_2 + z_1z_3 + z_2z_3$

Symmetric polynomials

$f(z_1, \dots, z_n)$ symmetric if swapping variables does not change f .

- $f(z_1, z_2, z_3) = z_1 + z_2 + z_3$
- $f(z_1, z_2, z_3) = z_1 z_2 + z_1 z_3 + z_2 z_3$
- $f(z_1, z_2, z_3) = z_1 z_2 z_3$

Symmetric polynomials

$f(z_1, \dots, z_n)$ symmetric if swapping variables does not change f .

- $f(z_1, z_2, z_3) = z_1 + z_2 + z_3$
- $f(z_1, z_2, z_3) = z_1z_2 + z_1z_3 + z_2z_3$
- $f(z_1, z_2, z_3) = z_1z_2z_3$

Facts

- e_d = sum of degree d square-free monomials.

Symmetric polynomials

$f(z_1, \dots, z_n)$ symmetric if swapping variables does not change f .

- $f(z_1, z_2, z_3) = z_1 + z_2 + z_3 = e_1$
- $f(z_1, z_2, z_3) = z_1z_2 + z_1z_3 + z_2z_3 = e_2$
- $f(z_1, z_2, z_3) = z_1z_2z_3 = e_3$

Facts

- $e_d =$ sum of degree d square-free monomials.

Symmetric polynomials

$f(z_1, \dots, z_n)$ symmetric if swapping variables does not change f .

- $f(z_1, z_2, z_3) = z_1 + z_2 + z_3 = e_1$
- $f(z_1, z_2, z_3) = z_1z_2 + z_1z_3 + z_2z_3 = e_2$
- $f(z_1, z_2, z_3) = z_1z_2z_3 = e_3$

Facts

- e_d = sum of degree d square-free monomials.
- Symmetric polynomials are an algebra generated by e_d 's.

Symmetric polynomials

$f(z_1, \dots, z_n)$ symmetric if swapping variables does not change f .

- $f(z_1, z_2, z_3) = z_1 + z_2 + z_3 = e_1$
- $f(z_1, z_2, z_3) = z_1z_2 + z_1z_3 + z_2z_3 = e_2$
- $f(z_1, z_2, z_3) = z_1z_2z_3 = e_3$

Facts

- e_d = sum of degree d square-free monomials.
- Symmetric polynomials are an algebra generated by e_d 's.
- A basis: products $e_{\lambda_1} \cdots e_{\lambda_n}$ for $\lambda_1 \geq \cdots \geq \lambda_n \geq 0$.

Symmetric polynomials

$f(z_1, \dots, z_n)$ symmetric if swapping variables does not change f .

- $f(z_1, z_2, z_3) = z_1 + z_2 + z_3 = e_1$
- $f(z_1, z_2, z_3) = z_1 z_2 + z_1 z_3 + z_2 z_3 = e_2$
- $f(z_1, z_2, z_3) = z_1 z_2 z_3 = e_3$

Facts

- e_d = sum of degree d square-free monomials.
- Symmetric polynomials are an algebra generated by e_d 's.
- A basis: products $e_{\lambda_1} \cdots e_{\lambda_n}$ for $\lambda_1 \geq \cdots \geq \lambda_n \geq 0$.

Natural occurrence

$$x^n + a_{n-1}x^{n-1} + \cdots + a_0 = \prod_{i=1}^n (x - \lambda_i)$$

Symmetric polynomials

$f(z_1, \dots, z_n)$ symmetric if swapping variables does not change f .

- $f(z_1, z_2, z_3) = z_1 + z_2 + z_3 = e_1$
- $f(z_1, z_2, z_3) = z_1 z_2 + z_1 z_3 + z_2 z_3 = e_2$
- $f(z_1, z_2, z_3) = z_1 z_2 z_3 = e_3$

Facts

- e_d = sum of degree d square-free monomials.
- Symmetric polynomials are an algebra generated by e_d 's.
- A basis: products $e_{\lambda_1} \cdots e_{\lambda_n}$ for $\lambda_1 \geq \cdots \geq \lambda_n \geq 0$.

Natural occurrence

$$x^n + a_{n-1}x^{n-1} + \cdots + a_0 = \prod_{i=1}^n (x - \lambda_i)$$

a_i = symmetric polynomial in $\lambda_1, \dots, \lambda_n$

Symmetric polynomials

$f(z_1, \dots, z_n)$ symmetric if swapping variables does not change f .

- $f(z_1, z_2, z_3) = z_1 + z_2 + z_3 = e_1$
- $f(z_1, z_2, z_3) = z_1 z_2 + z_1 z_3 + z_2 z_3 = e_2$
- $f(z_1, z_2, z_3) = z_1 z_2 z_3 = e_3$

Facts

- e_d = sum of degree d square-free monomials.
- Symmetric polynomials are an algebra generated by e_d 's.
- A basis: products $e_{\lambda_1} \cdots e_{\lambda_n}$ for $\lambda_1 \geq \cdots \geq \lambda_n \geq 0$.

Natural occurrence

$$x^n + a_{n-1}x^{n-1} + \cdots + a_0 = \prod_{i=1}^n (x - \lambda_i)$$

a_i = symmetric polynomial in $\lambda_1, \dots, \lambda_n = \pm e_{n-i}(\lambda_1, \dots, \lambda_n)$.

Symmetric polynomials

Linear algebra



Symmetric polynomials

Some representation theory

Setup

Group $G \curvearrowright V \cong \mathbb{C}^n$.

Some representation theory

Setup

Group $G \curvearrowright V \cong \mathbb{C}^n$. What are the G -invariant subspaces?

Some representation theory

Setup

Group $G \curvearrowright V \cong \mathbb{C}^n$. What are the G -invariant subspaces?

Traces

- $\text{tr}(G \curvearrowright V)$

Some representation theory

Setup

Group $G \curvearrowright V \cong \mathbb{C}^n$. What are the G -invariant subspaces?

Traces

- $\text{tr}(G \curvearrowright V)$
- $\text{tr}(G \curvearrowright U \oplus W) = \text{tr}(G \curvearrowright U) + \text{tr}(G \curvearrowright W).$

Some representation theory

Setup

Group $G \curvearrowright V \cong \mathbb{C}^n$. What are the G -invariant subspaces?

Traces

- $\text{tr}(G \curvearrowright V)$
- $\text{tr}(G \curvearrowright U \oplus W) = \text{tr}(G \curvearrowright U) + \text{tr}(G \curvearrowright W).$

Over \mathbb{C} , $G \curvearrowright V \cong \bigoplus \text{Irreducibles}$

$$\iff \text{tr}(G \curvearrowright V) = \sum \text{tr}(G \curvearrowright \text{Irreducible})$$

The miraculous basis of Schur polynomials



The miraculous basis of Schur polynomials

$$GL_2(\mathbb{C}) \curvearrowright \mathrm{Sym}^2(\mathbb{C}^2) \cong \mathbb{C}^3$$

The miraculous basis of Schur polynomials

$$GL_2(\mathbb{C}) \curvearrowright \text{Sym}^2(\mathbb{C}^2) \cong \mathbb{C}^3$$

Restrict to $\begin{pmatrix} z_1 & 0 \\ 0 & z_2 \end{pmatrix} \cdot v$

The miraculous basis of Schur polynomials

$$GL_2(\mathbb{C}) \curvearrowright \text{Sym}^2(\mathbb{C}^2) \cong \mathbb{C}^3$$

$$\text{Restrict to } \begin{pmatrix} z_1 & 0 \\ 0 & z_2 \end{pmatrix} \cdot v = \begin{pmatrix} z_1^2 & 0 & 0 \\ 0 & z_1 z_2 & 0 \\ 0 & 0 & z_2^2 \end{pmatrix} v$$

The miraculous basis of Schur polynomials

$$GL_2(\mathbb{C}) \curvearrowright \text{Sym}^2(\mathbb{C}^2) \cong \mathbb{C}^3$$

$$\text{Restrict to } \begin{pmatrix} z_1 & 0 \\ 0 & z_2 \end{pmatrix} \cdot v = \begin{pmatrix} z_1^2 & 0 & 0 \\ 0 & z_1 z_2 & 0 \\ 0 & 0 & z_2^2 \end{pmatrix} v$$

Trace = $z_1^2 + z_1 z_2 + z_2^2$ is symmetric!

The miraculous basis of Schur polynomials

$$GL_2(\mathbb{C}) \curvearrowright \text{Sym}^2(\mathbb{C}^2) \cong \mathbb{C}^3$$

$$\text{Restrict to } \begin{pmatrix} z_1 & 0 \\ 0 & z_2 \end{pmatrix} \cdot v = \begin{pmatrix} z_1^2 & 0 & 0 \\ 0 & z_1 z_2 & 0 \\ 0 & 0 & z_2^2 \end{pmatrix} v$$

Trace = $z_1^2 + z_1 z_2 + z_2^2$ is symmetric!

Integer partitions

$$\mu = (\mu_1 \geq \dots \geq \mu_n)$$

$$\mu_i \in \mathbb{Z}_{\geq 0}$$

Irreducible (polynomial) $GL_n(\mathbb{C})$
representations V^μ

The miraculous basis of Schur polynomials

$$GL_2(\mathbb{C}) \curvearrowright \text{Sym}^2(\mathbb{C}^2) \cong \mathbb{C}^3$$

$$\text{Restrict to } \begin{pmatrix} z_1 & 0 \\ 0 & z_2 \end{pmatrix} \cdot v = \begin{pmatrix} z_1^2 & 0 & 0 \\ 0 & z_1 z_2 & 0 \\ 0 & 0 & z_2^2 \end{pmatrix} v$$

Trace = $z_1^2 + z_1 z_2 + z_2^2$ is symmetric!

Integer partitions

$$\mu = (\mu_1 \geq \dots \geq \mu_n)$$

$$\mu_i \in \mathbb{Z}_{\geq 0}$$



Irreducible (polynomial) $GL_n(\mathbb{C})$
representations V^μ

Trace



Symmetric "Schur polynomial" s_μ

The miraculous basis of Schur polynomials

$$GL_2(\mathbb{C}) \curvearrowright \text{Sym}^2(\mathbb{C}^2) \cong \mathbb{C}^3$$

$$\text{Restrict to } \begin{pmatrix} z_1 & 0 \\ 0 & z_2 \end{pmatrix} \cdot v = \begin{pmatrix} z_1^2 & 0 & 0 \\ 0 & z_1 z_2 & 0 \\ 0 & 0 & z_2^2 \end{pmatrix} v$$

Trace = $z_1^2 + z_1 z_2 + z_2^2$ is symmetric!

Integer partitions

$$\mu = (\mu_1 \geq \dots \geq \mu_n) \\ \mu_i \in \mathbb{Z}_{\geq 0}$$




Irreducible (polynomial) $GL_n(\mathbb{C})$
representations V^μ

Trace


Symmetric "Schur polynomial" s_μ

$$s_{(2,0)}(z_1, z_2) = z_1^2 + z_1 z_2 + z_2^2.$$

The miraculous basis of Schur polynomials




The miraculous basis of Schur polynomials



Example

$f = z_1^2 + z_2^2$ the trace of some $GL_2(\mathbb{C})$ representation?

The miraculous basis of Schur polynomials




Example

$f = z_1^2 + z_2^2$ the trace of some $GL_2(\mathbb{C})$ representation?

Approach

$\{s_\mu\}$ is a basis of symmetric polynomials.

The miraculous basis of Schur polynomials



Example


$f = z_1^2 + z_2^2$ the trace of some $GL_2(\mathbb{C})$ representation?

Approach

$\{s_\mu\}$ is a basis of symmetric polynomials.

$$f = s_{(2,0)} - s_{(1,1)}$$

The miraculous basis of Schur polynomials



Example

$f = z_1^2 + z_2^2$ the trace of some $GL_2(\mathbb{C})$ representation?


Approach

$\{s_\mu\}$ is a basis of symmetric polynomials.

$$f = s_{(2,0)} - s_{(1,1)}$$

f not "Schur-positive" \implies not the trace of a GL_2 -representation!

The miraculous basis of Schur polynomials



Example

$f = z_1^2 + z_2^2$ the trace of some $GL_2(\mathbb{C})$ representation?

Approach

$\{s_\mu\}$ is a basis of symmetric polynomials.

$$f = s_{(2,0)} - s_{(1,1)}$$

f not "Schur-positive" \implies not the trace of a GL_2 -representation!

Guiding principle

Positive integer sums of Schurs \implies representation-theoretically meaningful.

The miraculous basis of Schur polynomials

$$GL_n(\mathbb{C}) \curvearrowright V^\mu$$

Trace
↓

Schur polynomial s_μ

The miraculous basis of Schur polynomials

$$GL_n(\mathbb{C}) \curvearrowright V^\mu$$

Trace

Schur polynomial s_μ

Semistandard Young Tableaux $\text{SSYT}(\mu)$


The miraculous basis of Schur polynomials

$$GL_n(\mathbb{C}) \curvearrowright V^\mu$$

Trace

Schur polynomial s_μ

Semistandard Young Tableaux $\text{SSYT}(\mu)$



The miraculous basis of Schur polynomials

$$GL_n(\mathbb{C}) \curvearrowright V^\mu$$

Trace

Schur polynomial s_μ

Semistandard Young Tableaux $\text{SSYT}(\mu)$

$$\mu = (4, 3, 1, 1) \rightarrow \begin{array}{|c|c|c|c|} \hline \square & & & \\ \hline \square & & & \\ \hline \square & \square & \square & \\ \hline \square & \square & \square & \square \\ \hline \end{array} \rightarrow T =$$



The miraculous basis of Schur polynomials

$$GL_n(\mathbb{C}) \curvearrowright V^\mu$$

Trace

Schur polynomial s_μ

Semistandard Young Tableaux $\text{SSYT}(\mu)$



The miraculous basis of Schur polynomials

$$GL_n(\mathbb{C}) \curvearrowright V^\mu$$

Trace

Schur polynomial s_μ

Semistandard Young Tableaux $SSYT(\mu)$

$$\mu = (4, 3, 1, 1) \rightarrow \begin{array}{|c|c|c|c|} \hline & & & \\ \hline & & & \\ \hline & & & \\ \hline & & & \\ \hline \end{array} \rightarrow T = \begin{array}{|c|c|c|c|} \hline z_5 & & & \\ \hline z_3 & & & \\ \hline z_2 & z_3 & z_5 & \\ \hline z_1 & z_1 & z_2 & z_3 \\ \hline \end{array}$$

The miraculous basis of Schur polynomials

$$GL_n(\mathbb{C}) \curvearrowright V^\mu$$

Trace

Schur polynomial s_μ

$$s_\mu = \sum_{T \in \text{SSYT}(\mu)} \prod_{z_i \in T} z_i$$

Sum of tableaux weights

Semistandard Young Tableaux $\text{SSYT}(\mu)$

$$\mu = (4, 3, 1, 1) \rightarrow$$

$$\rightarrow T =$$

z_5			
z_3			
z_2	z_3	z_5	
z_1	z_1	z_2	z_3

The miraculous basis of Schur polynomials

$$GL_n(\mathbb{C}) \curvearrowright V^\mu$$

Trace

Schur polynomial s_μ

$$s_\mu = \sum_{T \in \text{SSYT}(\mu)} \prod_{z_i \in T} z_i$$

Sum of tableaux weights

Semistandard Young Tableaux $\text{SSYT}(\mu)$

$$\mu = (4, 3, 1, 1) \rightarrow \begin{array}{|c|c|c|c|} \hline \square & & & \\ \hline \square & & & \\ \hline \square & \square & \square & \\ \hline \square & \square & \square & \square \\ \hline \end{array} \rightarrow T = \begin{array}{|c|c|c|c|} \hline z_5 & & & \\ \hline z_3 & & & \\ \hline z_2 & z_3 & z_5 & \\ \hline z_1 & z_1 & z_2 & z_3 \\ \hline \end{array}$$

A combinatorial interpretation

$\dim V^\mu = \text{number of semistandard Young tableaux of } \mu.$

The miraculous basis of Schur polynomials

- $S_n \curvearrowright V$

The miraculous basis of Schur polynomials

- $S_n \curvearrowright V$
- Integer partition of $n \leftrightarrow$ irreducible S_n representation V_μ

The miraculous basis of Schur polynomials

- $S_n \curvearrowright V$
- Integer partition of $n \leftrightarrow$ irreducible S_n representation V_μ

$$S_n \curvearrowright V_\mu$$

The miraculous basis of Schur polynomials

- $S_n \curvearrowright V$
- Integer partition of $n \leftrightarrow$ irreducible S_n representation V_μ

$$S_n \curvearrowright V_\mu$$

Frob

Schur polynomial s_μ

The miraculous basis of Schur polynomials

- $S_n \curvearrowright V$
- Integer partition of $n \leftrightarrow$ irreducible S_n representation V_μ

$$S_n \curvearrowright V_\mu$$

Frob

$$GL_n(\mathbb{C}) \curvearrowright V^\mu$$

Trace

Schur polynomial s_μ


Sum of weights

Semistandard Young Tableaux


z_5			
z_3			
z_2	z_3	z_5	
z_1	z_1	z_2	z_3

The miraculous basis of Schur polynomials


- $S_n \curvearrowright V$
- Integer partition of $n \leftrightarrow$ irreducible S_n representation V_μ




Upgrading with gradings




Upgrading with gradings




Upgrading with gradings



Upgrading with gradings




Upgrading with gradings



Shuffle Theorem Idea

Bigraded S_n representation \leftrightarrow Bigraded combinatorial formula

Upgrading with gradings



Shuffle Theorem Idea

Bigraded S_n representation \leftrightarrow Bigraded combinatorial formula

Generalized Shuffle Theorem Idea

(Blasiak, Haiman, Morse, Pun, S., 2023)

General family of symmetric polynomials with $\mathbb{Q}(q, t)$ coefficients
 \leftrightarrow Bigraded combinatorial formula

S_n -coinvariant representation

Symmetric group $S_n \curvearrowright \mathbb{C}[x_1, \dots, x_n]$

S_n -coinvariant representation

Symmetric group $S_n \curvearrowright \mathbb{C}[x_1, \dots, x_n]$



$$S_n \curvearrowright \mathcal{R}_n = \mathbb{C}[x_1, \dots, x_n]/I$$

$I =$ all positive degree symmetric polynomials

S_n -coinvariant representation

Symmetric group $S_n \curvearrowright \mathbb{C}[x_1, \dots, x_n]$



$S_n \curvearrowright \mathcal{R}_n = \mathbb{C}[x_1, \dots, x_n]/I$
 $I =$ all positive degree symmetric polynomials

\cong

$H^*(Fl_n(\mathbb{C}))$

S_n -coinvariant representation

Symmetric group $S_n \curvearrowright \mathbb{C}[x_1, \dots, x_n]$



$S_n \curvearrowright \mathcal{R}_n = \mathbb{C}[x_1, \dots, x_n]/I$
 $I = \text{all positive degree symmetric polynomials}$

\cong

$H^*(Fl_n(\mathbb{C}))$

$$\mathcal{R}_2 = \mathbb{C}[x_1, x_2]/\langle x_1 + x_2, x_1 x_2 \rangle$$

S_n -coinvariant representation

Symmetric group $S_n \curvearrowright \mathbb{C}[x_1, \dots, x_n]$



$S_n \curvearrowright \mathcal{R}_n = \mathbb{C}[x_1, \dots, x_n]/I$
 $I =$ all positive degree symmetric polynomials

\cong

$H^*(Fl_n(\mathbb{C}))$

$$\mathcal{R}_2 = \mathbb{C}[x_1, x_2]/\langle x_1 + x_2, x_1 x_2 \rangle \implies x_1 = -x_2 \text{ and } x_1^2 = x_2^2 = 0.$$

S_n -coinvariant representation

Symmetric group $S_n \curvearrowright \mathbb{C}[x_1, \dots, x_n]$



$S_n \curvearrowright \mathcal{R}_n = \mathbb{C}[x_1, \dots, x_n]/I$
 $I =$ all positive degree symmetric polynomials

\cong

$H^*(Fl_n(\mathbb{C}))$

$\mathcal{R}_2 = \mathbb{C}[x_1, x_2]/\langle x_1 + x_2, x_1 x_2 \rangle \implies x_1 = -x_2$ and $x_1^2 = x_2^2 = 0$.
For $S_2 \curvearrowright \mathcal{R}_2$, $\sigma = (1 \leftrightarrow 2)$,

$$\sigma.x_1 = x_2 = -x_1, \quad \sigma.1 = 1$$

S_n -coinvariant representation

Symmetric group $S_n \curvearrowright \mathbb{C}[x_1, \dots, x_n]$



$S_n \curvearrowright \mathcal{R}_n = \mathbb{C}[x_1, \dots, x_n]/I$
 $I =$ all positive degree symmetric polynomials

\cong

$H^*(Fl_n(\mathbb{C}))$

$\mathcal{R}_2 = \mathbb{C}[x_1, x_2]/\langle x_1 + x_2, x_1 x_2 \rangle \implies x_1 = -x_2$ and $x_1^2 = x_2^2 = 0$.
For $S_2 \curvearrowright \mathcal{R}_2$, $\sigma = (1 \leftrightarrow 2)$,

$$\sigma.x_1 = x_2 = -x_1, \quad \sigma.1 = 1$$

$$\implies \mathcal{R}_2 = \text{span}\{x_1\} \oplus \text{span}\{1\}.$$

S_n -coinvariant representation

Symmetric group $S_n \curvearrowright \mathbb{C}[x_1, \dots, x_n]$



$$S_n \curvearrowright \mathcal{R}_n = \mathbb{C}[x_1, \dots, x_n]/I \quad \xrightarrow{\cong} \quad H^*(Fl_n(\mathbb{C}))$$

$I =$ all positive degree symmetric polynomials

$\mathcal{R}_2 = \mathbb{C}[x_1, x_2]/\langle x_1 + x_2, x_1 x_2 \rangle \implies x_1 = -x_2$ and $x_1^2 = x_2^2 = 0$.
For $S_2 \curvearrowright \mathcal{R}_2$, $\sigma = (1 \leftrightarrow 2)$,

$$\sigma.x_1 = x_2 = -x_1, \quad \sigma.1 = 1$$

$$\implies \mathcal{R}_2 = \text{span}\{x_1\} \oplus \text{span}\{1\}.$$

Question

What are the irreducible (S_n -invariant) pieces of \mathcal{R}_n ?

S_n coinvariant representation

$$\mathcal{R}_2 \cong \text{span}\{x_1\} \oplus \text{span}\{1\}$$

S_n coinvariant representation

$$\begin{aligned}\mathcal{R}_2 &\cong \text{span}\{x_1\} \oplus \text{span}\{1\} \\ &\cong V_{(1,1)} \oplus V_{(2)}\end{aligned}$$

S_n coinvariant representation

$$\begin{aligned}\mathcal{R}_2 &\cong \text{span}\{x_1\} \oplus \text{span}\{1\} \\ &\cong V_{(1,1)} \oplus V_{(2)} \\ \implies \text{Frob}(\mathcal{R}_2) &= s_{(1,1)} + s_{(2)}\end{aligned}$$

S_n coinvariant representation

$$\begin{aligned}\mathcal{R}_2 &\cong \text{span}\{x_1\} \oplus \text{span}\{1\} \\ &\cong V_{(1,1)} \oplus V_{(2)} \\ \implies \text{Frob}(\mathcal{R}_2) &= s_{(1,1)} + s_{(2)}\end{aligned}$$

Identification ignores grading

S_n coinvariant representation

$$\begin{aligned}\mathcal{R}_2 &\cong \text{span}\{x_1\} \oplus \text{span}\{1\} \\ &\cong V_{(1,1)} \oplus V_{(2)} \\ \implies \text{Frob}(\mathcal{R}_2) &= s_{(1,1)} + s_{(2)}\end{aligned}$$

Identification ignores grading \implies add coefficient parameter q .

S_n coinvariant representation

$$\begin{aligned}\mathcal{R}_2 &\cong \text{span}\{x_1\} \oplus \text{span}\{1\} \\ &\cong V_{(1,1)} \oplus V_{(2)} \\ \implies \text{Frob}(\mathcal{R}_2) &= s_{(1,1)} + s_{(2)}\end{aligned}$$

Identification ignores grading \implies add coefficient parameter q .

$$\mathcal{R}_2 \cong \text{span}\{x_1\} \oplus \text{span}\{1\}$$

S_n coinvariant representation

$$\begin{aligned}\mathcal{R}_2 &\cong \text{span}\{x_1\} \oplus \text{span}\{1\} \\ &\cong V_{(1,1)} \oplus V_{(2)} \\ \implies \text{Frob}(\mathcal{R}_2) &= s_{(1,1)} + s_{(2)}\end{aligned}$$

Identification ignores grading \implies add coefficient parameter q .

$$\begin{aligned}\mathcal{R}_2 &\cong \text{span}\{x_1\} \oplus \text{span}\{1\} \\ \text{Frob}_q(\mathcal{R}_2) &= qs_{(1,1)} + s_{(2)}\end{aligned}$$

S_n coinvariant representation

$$\begin{aligned}\mathcal{R}_2 &\cong \text{span}\{x_1\} \oplus \text{span}\{1\} \\ &\cong V_{(1,1)} \oplus V_{(2)} \\ \implies \text{Frob}(\mathcal{R}_2) &= s_{(1,1)} + s_{(2)}\end{aligned}$$

Identification ignores grading \implies add coefficient parameter q .

$$\begin{aligned}\mathcal{R}_2 &\cong \text{span}\{x_1\} \oplus \text{span}\{1\} \\ \text{Frob}_q(\mathcal{R}_2) &= qs_{(1,1)} + s_{(2)}\end{aligned}$$

$$\text{Frob}(\mathcal{R}_3) = s_{(1,1,1)} + 2s_{(2,1)} + s_{(3)}$$

S_n coinvariant representation

$$\begin{aligned}\mathcal{R}_2 &\cong \text{span}\{x_1\} \oplus \text{span}\{1\} \\ &\cong V_{(1,1)} \oplus V_{(2)} \\ \implies \text{Frob}(\mathcal{R}_2) &= s_{(1,1)} + s_{(2)}\end{aligned}$$

Identification ignores grading \implies add coefficient parameter q .

$$\begin{aligned}\mathcal{R}_2 &\cong \text{span}\{x_1\} \oplus \text{span}\{1\} \\ \text{Frob}_q(\mathcal{R}_2) &= qs_{(1,1)} + s_{(2)}\end{aligned}$$

$$\begin{aligned}\text{Frob}(\mathcal{R}_3) &= s_{(1,1,1)} + 2s_{(2,1)} + s_{(3)} \\ \text{Frob}_q(\mathcal{R}_3) &= q^3 s_{(1,1,1)} + (q^2 + q)s_{(2,1)} + s_{(3)}\end{aligned}$$

S_n coinvariant representation

$$\begin{aligned}\mathcal{R}_2 &\cong \text{span}\{x_1\} \oplus \text{span}\{1\} \\ &\cong V_{(1,1)} \oplus V_{(2)} \\ \implies \text{Frob}(\mathcal{R}_2) &= s_{(1,1)} + s_{(2)}\end{aligned}$$

Identification ignores grading \implies add coefficient parameter q .

$$\begin{aligned}\mathcal{R}_2 &\cong \text{span}\{x_1\} \oplus \text{span}\{1\} \\ \text{Frob}_q(\mathcal{R}_2) &= qs_{(1,1)} + s_{(2)}\end{aligned}$$

$$\begin{aligned}\text{Frob}(\mathcal{R}_3) &= s_{(1,1,1)} + 2s_{(2,1)} + s_{(3)} \\ \text{Frob}_q(\mathcal{R}_3) &= q^3 s_{(1,1,1)} + (q^2 + q)s_{(2,1)} + s_{(3)}\end{aligned}$$

Combinatorial Interpretation

- What is $\text{Frob}_q(\mathcal{R}_n)$?

S_n coinvariant representation

$$\begin{aligned}\mathcal{R}_2 &\cong \text{span}\{x_1\} \oplus \text{span}\{1\} \\ &\cong V_{(1,1)} \oplus V_{(2)} \\ \implies \text{Frob}(\mathcal{R}_2) &= s_{(1,1)} + s_{(2)}\end{aligned}$$

Identification ignores grading \implies add coefficient parameter q .

$$\begin{aligned}\mathcal{R}_2 &\cong \text{span}\{x_1\} \oplus \text{span}\{1\} \\ \text{Frob}_q(\mathcal{R}_2) &= qs_{(1,1)} + s_{(2)}\end{aligned}$$

$$\begin{aligned}\text{Frob}(\mathcal{R}_3) &= s_{(1,1,1)} + 2s_{(2,1)} + s_{(3)} \\ \text{Frob}_q(\mathcal{R}_3) &= q^3 s_{(1,1,1)} + (q^2 + q)s_{(2,1)} + s_{(3)}\end{aligned}$$

Combinatorial Interpretation

- What is $\text{Frob}_q(\mathcal{R}_n)$? A “Hall-Littlewood polynomial.”

S_n coinvariant representation

$$\begin{aligned}\mathcal{R}_2 &\cong \text{span}\{x_1\} \oplus \text{span}\{1\} \\ &\cong V_{(1,1)} \oplus V_{(2)} \\ \implies \text{Frob}(\mathcal{R}_2) &= s_{(1,1)} + s_{(2)}\end{aligned}$$

Identification ignores grading \implies add coefficient parameter q .

$$\begin{aligned}\mathcal{R}_2 &\cong \text{span}\{x_1\} \oplus \text{span}\{1\} \\ \text{Frob}_q(\mathcal{R}_2) &= qs_{(1,1)} + s_{(2)}\end{aligned}$$

$$\begin{aligned}\text{Frob}(\mathcal{R}_3) &= s_{(1,1,1)} + 2s_{(2,1)} + s_{(3)} \\ \text{Frob}_q(\mathcal{R}_3) &= q^3 s_{(1,1,1)} + (q^2 + q)s_{(2,1)} + s_{(3)}\end{aligned}$$

Combinatorial Interpretation

- What is $\text{Frob}_q(\mathcal{R}_n)$? A “Hall-Littlewood polynomial.”
- Combinatorics for coefficients?

S_n coinvariant representation

$$\begin{aligned}\mathcal{R}_2 &\cong \text{span}\{x_1\} \oplus \text{span}\{1\} \\ &\cong V_{(1,1)} \oplus V_{(2)} \\ \implies \text{Frob}(\mathcal{R}_2) &= s_{(1,1)} + s_{(2)}\end{aligned}$$

Identification ignores grading \implies add coefficient parameter q .

$$\begin{aligned}\mathcal{R}_2 &\cong \text{span}\{x_1\} \oplus \text{span}\{1\} \\ \text{Frob}_q(\mathcal{R}_2) &= qs_{(1,1)} + s_{(2)}\end{aligned}$$

$$\begin{aligned}\text{Frob}(\mathcal{R}_3) &= s_{(1,1,1)} + 2s_{(2,1)} + s_{(3)} \\ \text{Frob}_q(\mathcal{R}_3) &= q^3 s_{(1,1,1)} + (q^2 + q)s_{(2,1)} + s_{(3)}\end{aligned}$$

Combinatorial Interpretation

- What is $\text{Frob}_q(\mathcal{R}_n)$? A “Hall-Littlewood polynomial.”
- Combinatorics for coefficients? Lascoux-Schützenberger, 1985

Natural generalization: Diagonal coinvariants

Symmetric group $S_n \curvearrowright \mathbb{C}[x_1, \dots, x_n, y_1, \dots, y_n]$

Natural generalization: Diagonal coinvariants

Symmetric group $S_n \curvearrowright \mathbb{C}[x_1, \dots, x_n, y_1, \dots, y_n]$



$$S_n \curvearrowright D\mathcal{R}_n = \mathbb{C}[x_1, \dots, x_n, y_1, \dots, y_n]/I$$

I = all positive degree polynomials symmetric in x_i 's and in y_i 's

Natural generalization: Diagonal coinvariants

Symmetric group $S_n \curvearrowright \mathbb{C}[x_1, \dots, x_n, y_1, \dots, y_n]$



$$S_n \curvearrowright D\mathcal{R}_n = \mathbb{C}[x_1, \dots, x_n, y_1, \dots, y_n]/I$$

I = all positive degree polynomials symmetric in x_i 's and in y_i 's

$$D\mathcal{R}_2 = \text{span}\{x_1\} \oplus \text{span}\{y_1\} \oplus \text{span}\{1\}$$

$$\cong V_{(1,1)} \oplus V_{(1,1)} \oplus V_{(2)}$$

Natural generalization: Diagonal coinvariants

Symmetric group $S_n \curvearrowright \mathbb{C}[x_1, \dots, x_n, y_1, \dots, y_n]$



$$S_n \curvearrowright D\mathcal{R}_n = \mathbb{C}[x_1, \dots, x_n, y_1, \dots, y_n]/I$$

I = all positive degree polynomials symmetric in x_i 's and in y_i 's

$$D\mathcal{R}_2 = \text{span}\{x_1\} \oplus \text{span}\{y_1\} \oplus \text{span}\{1\}$$

$$\cong V_{(1,1)} \oplus V_{(1,1)} \oplus V_{(2)}$$

Identification ignores bigrading \implies add coefficient parameters q, t .

Natural generalization: Diagonal coinvariants

Symmetric group $S_n \curvearrowright \mathbb{C}[x_1, \dots, x_n, y_1, \dots, y_n]$



$$S_n \curvearrowright D\mathcal{R}_n = \mathbb{C}[x_1, \dots, x_n, y_1, \dots, y_n]/I$$

I = all positive degree polynomials symmetric in x_i 's and in y_i 's

$$D\mathcal{R}_2 = \text{span}\{x_1\} \oplus \text{span}\{y_1\} \oplus \text{span}\{1\}$$

$$\cong V_{(1,1)} \oplus V_{(1,1)} \oplus V_{(2)}$$

Identification ignores bigrading \implies add coefficient parameters q, t .

$$D\mathcal{R}_2 = \text{span}\{x_1\} \oplus \text{span}\{y_1\} \oplus \text{span}\{1\}$$

Natural generalization: Diagonal coinvariants

Symmetric group $S_n \curvearrowright \mathbb{C}[x_1, \dots, x_n, y_1, \dots, y_n]$



$$S_n \curvearrowright D\mathcal{R}_n = \mathbb{C}[x_1, \dots, x_n, y_1, \dots, y_n]/I$$

I = all positive degree polynomials symmetric in x_i 's and in y_i 's

$$D\mathcal{R}_2 = \text{span}\{x_1\} \oplus \text{span}\{y_1\} \oplus \text{span}\{1\}$$

$$\cong V_{(1,1)} \oplus V_{(1,1)} \oplus V_{(2)}$$

Identification ignores bigrading \implies add coefficient parameters q, t .

$$D\mathcal{R}_2 = \text{span}\{x_1\} \oplus \text{span}\{y_1\} \oplus \text{span}\{1\}$$

$$\text{Frob}_{q,t}(D\mathcal{R}_2) = qs_{(1,1)} + ts_{(1,1)} + s_{(2)}$$

Natural generalization: Diagonal coinvariants

Symmetric group $S_n \curvearrowright \mathbb{C}[x_1, \dots, x_n, y_1, \dots, y_n]$



$$S_n \curvearrowright D\mathcal{R}_n = \mathbb{C}[x_1, \dots, x_n, y_1, \dots, y_n] / I$$

I = all positive degree polynomials symmetric in x_i 's and in y_i 's

$$D\mathcal{R}_2 = \text{span}\{x_1\} \oplus \text{span}\{y_1\} \oplus \text{span}\{1\}$$

$$\cong V_{(1,1)} \oplus V_{(1,1)} \oplus V_{(2)}$$

Identification ignores bigrading \implies add coefficient parameters q, t .

$$D\mathcal{R}_2 = \text{span}\{x_1\} \oplus \text{span}\{y_1\} \oplus \text{span}\{1\}$$

$$\text{Frob}_{q,t}(D\mathcal{R}_2) = qs_{(1,1)} + ts_{(1,1)} + s_{(2)}$$

Foundational unsolved question

What are the combinatorics for this expression?

Natural generalization: Diagonal coinvariants

Symmetric group $S_n \curvearrowright \mathbb{C}[x_1, \dots, x_n, y_1, \dots, y_n]$



$$S_n \curvearrowright D\mathcal{R}_n = \mathbb{C}[x_1, \dots, x_n, y_1, \dots, y_n] / I$$

I = all positive degree polynomials symmetric in x_i 's and in y_i 's

$$D\mathcal{R}_2 = \text{span}\{x_1\} \oplus \text{span}\{y_1\} \oplus \text{span}\{1\}$$

$$\cong V_{(1,1)} \oplus V_{(1,1)} \oplus V_{(2)}$$

Identification ignores bigrading \implies add coefficient parameters q, t .

$$D\mathcal{R}_2 = \text{span}\{x_1\} \oplus \text{span}\{y_1\} \oplus \text{span}\{1\}$$

$$\text{Frob}_{q,t}(D\mathcal{R}_2) = qs_{(1,1)} + ts_{(1,1)} + s_{(2)}$$

Foundational unsolved question

What are the combinatorics for this expression?

$$\text{Frob}_{q,t}(D\mathcal{R}_3) = (q^3 + q^2t + qt^2 + t^3 + qt)s_{111} + (q^2 + qt + t^2 + q + t)s_{21} + s_3$$


The fluttering of a combinatorialist's heart

What do the following objects have in common?

The fluttering of a combinatorialist's heart

What do the following objects have in common?


- Dyck paths in an $n \times n$ grid.



The fluttering of a combinatorialist's heart

What do the following objects have in common?

- Dyck paths in an $n \times n$ grid.




- Triangulations of an $(n + 2)$ -gon



The fluttering of a combinatorialist's heart

What do the following objects have in common?

- Dyck paths in an $n \times n$ grid.



- Triangulations of an $(n + 2)$ -gon



- The number of $2 \times n$ rectangle standard Young tableaux

4	5	6
1	2	3

3	4	6
1	2	5

3	5	6
1	2	4


2	5	6
1	3	4

2	4	6
1	3	5

The fluttering of a combinatorialist's heart

What do the following objects have in common?

- Dyck paths in an $n \times n$ grid.



- Triangulations of an $(n+2)$ -gon



- The number of $2 \times n$ rectangle standard Young tableaux

4	5	6
1	2	3

3	4	6
1	2	5

3	5	6
1	2	4

2	5	6
1	3	4

2	4	6
1	3	5


- Number of terms in coefficient of $s_{(1,\dots,1)}$ in $\text{Frob}_{q,t}(\mathcal{DR}_n)$?

$$(q^3 + q^2t + qt^2 + t^3 + qt)s_{111} + \cdots$$

The fluttering of a combinatorialist's heart

What do the following objects have in common?

- Dyck paths in an $n \times n$ grid.



- Triangulations of an $(n+2)$ -gon



- The number of $2 \times n$ rectangle standard Young tableaux

4	5	6
1	2	3

3	4	6
1	2	5

3	5	6
1	2	4

2	5	6
1	3	4

2	4	6
1	3	5

- Number of terms in coefficient of $s_{(1,\dots,1)}$ in $\text{Frob}_{q,t}(\mathcal{DR}_n)$?

$$(q^3 + q^2t + qt^2 + t^3 + qt)s_{111} + \cdots$$

Catalan numbers!

Catalan numbers C_n

Catalan numbers C_n

$$C_0 = 1 \text{ and } C_n = \sum_{i=1}^n C_{i-1} C_{n-i}$$

Catalan numbers C_n

$$C_0 = 1 \text{ and } C_n = \sum_{i=1}^n C_{i-1} C_{n-i}$$

1, 1, 2, 5, 14, 42, 132, ...

Catalan numbers C_n

$$C_0 = 1 \text{ and } C_n = \sum_{i=1}^n C_{i-1} C_{n-i}$$

1, 1, 2, 5, 14, 42, 132, ...



甲乙與丙庚爲第一率
二位二率爲四率
三位三率爲四率
四位一率除

一	二	三	四	五	六	七	八	九	十	十一	十二	十三	十四	十五	十六	十七	十八	十九	二十
1	1	2	5	14	42	132	429	1430	4862	16796	58786	208012	742900	2673165	9694845	35357670	129644790	477735720	1771329920



Catalan numbers C_n




$$C_0 = 1 \text{ and } C_n = \sum_{i=1}^n C_{i-1} C_{n-i}$$

$$1, 1, 2, 5, 14, 42, 132, \dots$$


Catalan numbers C_n

$$C_0 = 1 \text{ and } C_n = \sum_{i=1}^n C_{i-1} C_{n-i}$$

1, 1, 2, 5, 14, 42, 132, ...






q, t Catalan “numbers”

Write $\text{Frob}_{q,t}(DR_n) = C_n(q, t)s_{(1,\dots,1)} + \dots$

Catalan numbers C_n

$$C_0 = 1 \text{ and } C_n = \sum_{i=1}^n C_{i-1} C_{n-i}$$

1, 1, 2, 5, 14, 42, 132, ...






q, t Catalan “numbers”

Write $\text{Frob}_{q,t}(DR_n) = C_n(q, t)s_{(1, \dots, 1)} + \dots$
 $C_n(q, t) \in \mathbb{N}[q, t]$ and $C_n(1, 1) = C_n$.

Catalan numbers C_n

$$C_0 = 1 \text{ and } C_n = \sum_{i=1}^n C_{i-1} C_{n-i}$$

1, 1, 2, 5, 14, 42, 132, ...



q, t Catalan “numbers”

Write $\text{Frob}_{q,t}(D\mathcal{R}_n) = C_n(q, t)s_{(1,\dots,1)} + \dots$

$C_n(q, t) \in \mathbb{N}[q, t]$ and $C_n(1, 1) = C_n$.

$\text{Frob}_{q,t}(D\mathcal{R}_n) =$ Sum controlled by Catalan-many objects?

Recall: $e_i =$ sum of degree i square-free monomials.

Recall: e_i = sum of degree i square-free monomials.

Special case

$$\text{Frob}_{q,t}(D\mathcal{R}_n)|_{q=1} = \sum_{\lambda \in DP_n} (t \text{ monomial})(\text{product of } e_i\text{'s})$$

Recall: e_i = sum of degree i square-free monomials.

Special case

$$\text{Frob}_{q,t}(D\mathcal{R}_n)|_{q=1} = \sum_{\lambda \in DP_n} (t \text{ monomial})(\text{product of } e_i\text{'s})$$

λ $(t \text{ monomial})(\text{product of } e_i\text{'s})$



Recall: e_i = sum of degree i square-free monomials.

Special case

$$\text{Frob}_{q,t}(D\mathcal{R}_n)|_{q=1} = \sum_{\lambda \in DP_n} (t \text{ monomial})(\text{product of } e_i\text{'s})$$

λ $(t \text{ monomial})(\text{product of } e_i\text{'s})$



t^1



Recall: e_i = sum of degree i square-free monomials.

Special case

$$\text{Frob}_{q,t}(D\mathcal{R}_n)|_{q=1} = \sum_{\lambda \in DP_n} (t \text{ monomial})(\text{product of } e_i\text{'s})$$

λ $(t \text{ monomial})(\text{product of } e_i\text{'s})$



$$t^1 e_2 e_1$$



Recall: e_i = sum of degree i square-free monomials.

Special case

$$\text{Frob}_{q,t}(D\mathcal{R}_n)|_{q=1} = \sum_{\lambda \in DP_n} (t \text{ monomial})(\text{product of } e_i\text{'s})$$

λ $(t \text{ monomial})(\text{product of } e_i\text{'s})$



$$e_1 e_1 e_1$$



$$t^1 e_2 e_1$$



$$t^1 e_2 e_1$$



$$t^2 e_2 e_1$$



$$t^3 e_3$$

Recall: e_i = sum of degree i square-free monomials.

Special case

$$\text{Frob}_{q,t}(D\mathcal{R}_n)|_{q=1} = \sum_{\lambda \in DP_n} (t \text{ monomial})(\text{product of } e_i\text{'s})$$

λ $(t \text{ monomial})(\text{product of } e_i\text{'s})$



$$e_1 e_1 e_1$$



$$t^1 e_2 e_1$$



$$t^1 e_2 e_1$$



$$t^2 e_2 e_1$$



$$t^3 e_3$$

$$\implies \text{Frob}_{q,t}(D\mathcal{R}_n)|_{q=1} = e_1^3 + (2t + t^2)e_2 e_1 + t^3 e_3$$

The Shuffle Theorem

The Shuffle Theorem (Carlsson-Mellit, 2018)

$$\mathrm{Frob}_{q,t}(D\mathcal{R}_n) =$$

The Shuffle Theorem

The Shuffle Theorem (Carlsson-Mellit, 2018)

$$\text{Frob}_{q,t}(D\mathcal{R}_n) = \sum_{\lambda \in DP_n} (q, t \text{ monomial})(\text{LLT polynomial})$$

The Shuffle Theorem

The Shuffle Theorem (Carlsson-Mellit, 2018)

$$\text{Frob}_{q,t}(D\mathcal{R}_n) = \sum_{\lambda \in DP_n} (q, t \text{ monomial})(\text{LLT polynomial})$$

- LLT polynomial = Symmetric polynomial with $\mathbb{N}[q]$ coefficients (Lascoux, Leclerc, Thibon, 1997).

The Shuffle Theorem

The Shuffle Theorem (Carlsson-Mellit, 2018)

$$\text{Frob}_{q,t}(D\mathcal{R}_n) = \sum_{\lambda \in DP_n} (q, t \text{ monomial})(\text{LLT polynomial})$$

- LLT polynomial = Symmetric polynomial with $\mathbb{N}[q]$ coefficients (Lascoux, Leclerc, Thibon, 1997).
- Conjectured by (Haglund, Haiman, Loehr, Remmel, Ulyanov, 2005).

The Shuffle Theorem

The Shuffle Theorem (Carlsson-Mellit, 2018)

$$\text{Frob}_{q,t}(D\mathcal{R}_n) = \sum_{\lambda \in DP_n} (q, t \text{ monomial})(\text{LLT polynomial})$$

- LLT polynomial = Symmetric polynomial with $\mathbb{N}[q]$ coefficients (Lascoux, Leclerc, Thibon, 1997).
- Conjectured by (Haglund, Haiman, Loehr, Remmel, Ulyanov, 2005).
- LLT polynomial Schur expansion coefficients in $\mathbb{N}[q]$ (Grojnowski, Haiman, 2007).

The Shuffle Theorem

The Shuffle Theorem (Carlsson-Mellit, 2018)

$$\text{Frob}_{q,t}(D\mathcal{R}_n) = \sum_{\lambda \in DP_n} (q, t \text{ monomial})(\text{LLT polynomial})$$

- LLT polynomial = Symmetric polynomial with $\mathbb{N}[q]$ coefficients (Lascoux, Leclerc, Thibon, 1997).
- Conjectured by (Haglund, Haiman, Loehr, Remmel, Ulyanov, 2005).
- LLT polynomial Schur expansion coefficients in $\mathbb{N}[q]$ (Grojnowski, Haiman, 2007).

Special Case (Garsia, Haglund, 2001+2002)

$$C_n(q, t) = \sum_{\lambda \in DP_n} (q, t \text{ monomial})$$

Generalizing

LHS of Shuffle Theorem: $\text{Frob}_{q,t}(D\mathcal{R}_k)$.

Generalizing

LHS of Shuffle Theorem: $\text{Frob}_{q,t}(D\mathcal{R}_k)$.

Theorem (Haiman, 2002)

For operator $e_k^{(1,1)}$ on symmetric polynomials,
 $e_k^{(1,1)}(1) = \text{Frob}_{q,t}(D\mathcal{R}_k)$.


Generalizing

LHS of Shuffle Theorem: $\text{Frob}_{q,t}(D\mathcal{R}_k)$.

Theorem (Haiman, 2002)

For operator $e_k^{(1,1)}$ on symmetric polynomials,
 $e_k^{(1,1)}(1) = \text{Frob}_{q,t}(D\mathcal{R}_k)$.

$e_k^{(1,1)}$



Shuffle
Theorem



Generalizing

LHS of Shuffle Theorem: $\text{Frob}_{q,t}(D\mathcal{R}_k)$.

Theorem (Haiman, 2002)

For operator $e_k^{(1,1)}$ on symmetric polynomials,
 $e_k^{(1,1)}(1) = \text{Frob}_{q,t}(D\mathcal{R}_k)$.

$e_k^{(1,1)}$



Shuffle
Theorem




Generalizing

LHS of Shuffle Theorem: $\text{Frob}_{q,t}(D\mathcal{R}_k)$.

Theorem (Haiman, 2002)

For operator $e_k^{(1,1)}$ on symmetric polynomials,
 $e_k^{(1,1)}(1) = \text{Frob}_{q,t}(D\mathcal{R}_k)$.

$e_k^{(1,1)}$



Shuffle
Theorem





Generalizing

LHS of Shuffle Theorem: $\text{Frob}_{q,t}(D\mathcal{R}_k)$.

Theorem (Haiman, 2002)

For operator $e_k^{(1,1)}$ on symmetric polynomials,
 $e_k^{(1,1)}(1) = \text{Frob}_{q,t}(D\mathcal{R}_k)$.

$e_k^{(1,1)}$



Shuffle
Theorem

Generalizing

LHS of Shuffle Theorem: $\text{Frob}_{q,t}(D\mathcal{R}_k)$.

Theorem (Haiman, 2002)




For operator $e_k^{(1,1)}$ on symmetric polynomials,
 $e_k^{(1,1)}(1) = \text{Frob}_{q,t}(D\mathcal{R}_k)$.

$e_k^{(1,1)}$

operator?

operator?

operator?



Shuffle
Theorem

Generalizing with the Elliptic Hall Algebra

Algebra $\mathcal{E} \curvearrowright \Lambda =$ symmetric polynomials

Generalizing with the Elliptic Hall Algebra

Algebra $\mathcal{E} \curvearrowright \Lambda = \text{symmetric polynomials}$

$\mathcal{E} =$ Hall algebra of Coherent sheaves on
an elliptic curve over a finite field

Generalizing with the Elliptic Hall Algebra

Algebra $\mathcal{E} \curvearrowright \Lambda =$ symmetric polynomials

$\mathcal{E} =$ Hall algebra of Coherent sheaves on
an elliptic curve over a finite field


$$\mathcal{E} \cong \text{central subalgebra} \oplus \bigoplus_{m,n \text{ coprime}} \Lambda^{(m,n)}$$

Generalizing with the Elliptic Hall Algebra

Algebra $\mathcal{E} \curvearrowright \Lambda = \text{symmetric polynomials}$

$\mathcal{E} =$ Hall algebra of Coherent sheaves on
an elliptic curve over a finite field

$$\mathcal{E} \cong \text{central subalgebra} \oplus \bigoplus_{m,n \text{ coprime}} \Lambda^{(m,n)}$$




Generalizing with the Elliptic Hall Algebra

Algebra $\mathcal{E} \curvearrowright \Lambda =$ symmetric polynomials

$\mathcal{E} =$ Hall algebra of Coherent sheaves on
an elliptic curve over a finite field

$$\mathcal{E} \cong \text{central subalgebra} \oplus \bigoplus_{m,n \text{ coprime}} \Lambda^{(m,n)}$$




Generalizing with the Elliptic Hall Algebra

Algebra $\mathcal{E} \hookrightarrow \Lambda =$ symmetric polynomials

$\mathcal{E} =$ Hall algebra of Coherent sheaves on an elliptic curve over a finite field

$$\mathcal{E} \cong \text{central subalgebra} \oplus \bigoplus_{m,n \text{ coprime}} \Lambda^{(m,n)}$$




LHS of Shuffle Theorem $= e_k^{(1,1)} \in \Lambda^{(1,1)}$ acting on $1 \in \Lambda$.

Generalizing with the Elliptic Hall Algebra

Algebra $\mathcal{E} \curvearrowright \Lambda =$ symmetric polynomials

$\mathcal{E} =$ Hall algebra of Coherent sheaves on an elliptic curve over a finite field

$$\mathcal{E} \cong \text{central subalgebra} \oplus \bigoplus_{m,n \text{ coprime}} \bigoplus \Lambda^{(m,n)}$$



LHS of Shuffle Theorem $= e_k^{(1,1)} \in \Lambda^{(1,1)}$ acting on $1 \in \Lambda$.

Question

What can we say about $e_k^{(m,n)} \in \Lambda^{(m,n)}$ acting on $1 \in \Lambda$?

Rational Shuffle Theorem

Rational Shuffle Theorem (Mellit, 2021)


$$e_k^{(m,n)} \cdot 1 = \sum (q, t \text{ monomial})(\text{LLT polynomial})$$

Rational Shuffle Theorem

Rational Shuffle Theorem (Mellit, 2021)

$$e_k^{(m,n)} \cdot 1 = \sum (q, t \text{ monomial})(\text{LLT polynomial})$$

- Sum over $km \times kn$ Dyck paths




Rational Shuffle Theorem

Rational Shuffle Theorem (Mellit, 2021)

$$e_k^{(m,n)} \cdot 1 = \sum (q, t \text{ monomial})(\text{LLT polynomial})$$

- Sum over $km \times kn$ Dyck paths
- Conjectured by (F. Bergeron, Garsia, Sergel Leven, Xin, 2016)




Rational Shuffle Theorem

Rational Shuffle Theorem (Mellit, 2021)

$$e_k^{(m,n)} \cdot 1 = \sum (q, t \text{ monomial})(\text{LLT polynomial})$$

- Sum over $km \times kn$ Dyck paths
- Conjectured by (F. Bergeron, Garsia, Sergel Leven, Xin, 2016)





Rational Shuffle Theorem

Rational Shuffle Theorem (Mellit, 2021)

$$e_k^{(m,n)} \cdot 1 = \sum (q, t \text{ monomial})(\text{LLT polynomial})$$

- Sum over $km \times kn$ Dyck paths
- Conjectured by (F. Bergeron, Garsia, Sergel Leven, Xin, 2016)



Generalized Shuffle Theorem

Generalized Shuffle Theorem

Definition (Neguț, 2014)

$r, s \in \mathbb{R}_{>0} \rightarrow D^{(r,s)} \in \mathcal{E}$ such that $D^{(km, kn)} = e_k^{(m,n)}$.

Generalized Shuffle Theorem

Definition (Neguț, 2014)

$r, s \in \mathbb{R}_{>0} \rightarrow D^{(r,s)} \in \mathcal{E}$ such that $D^{(km, kn)} = e_k^{(m,n)}$.

Theorem (Blasiak, Haiman, Morse, Pun, S., 2023)

$D^{(r,s)} \cdot 1 = \sum (q, t \text{ monomial})(\text{LLT polynomial})$

Generalized Shuffle Theorem


Definition (Neguț, 2014)

$r, s \in \mathbb{R}_{>0} \rightarrow D^{(r,s)} \in \mathcal{E}$ such that $D^{(km,kn)} = e_k^{(m,n)}$.

Theorem (Blasiak, Haiman, Morse, Pun, S., 2023)

$D^{(r,s)} \cdot 1 = \sum (q, t \text{ monomial})(\text{LLT polynomial})$

Sum over all Dyck paths lying below the line $y = -\frac{s}{r}x + s$.



Generalized Shuffle Theorem

Definition (Neguț, 2014)


$r, s \in \mathbb{R}_{>0} \rightarrow D^{(r,s)} \in \mathcal{E}$ such that $D^{(km,kn)} = e_k^{(m,n)}$.

Theorem (Blasiak, Haiman, Morse, Pun, S., 2023)

$$D^{(r,s)} \cdot 1 = \sum (q, t \text{ monomial})(\text{LLT polynomial})$$

Sum over all Dyck paths lying below the line $y = -\frac{s}{r}x + s$.

$D^{(r,s)}$



Generalized Shuffle Theorem

Definition (Neguț, 2014)


$r, s \in \mathbb{R}_{>0} \rightarrow D^{(r,s)} \in \mathcal{E}$ such that $D^{(km,kn)} = e_k^{(m,n)}$.

Theorem (Blasiak, Haiman, Morse, Pun, S., 2023)


$$D^{(r,s)} \cdot 1 = \sum (q, t \text{ monomial})(\text{LLT polynomial})$$

Sum over all Dyck paths lying below the line $y = -\frac{s}{r}x + s$.

$e_k^{(m,n)}$



$D^{(r,s)}$



Generalized Shuffle Theorem

Theorem (Blasiak, Haiman, Morse, Pun, S., 2023)

$$D^{(r,s)} \cdot 1 = \sum_{\lambda} (q, t \text{ monomial})(\text{LLT polynomial})$$

Sum over all Dyck paths lying below the line $y = -\frac{s}{r}x + s$.

Generalized Shuffle Theorem

Theorem (Blasiak, Haiman, Morse, Pun, S., 2023)

$$D^{(r,s)} \cdot 1 = \sum_{\lambda} (q, t \text{ monomial})(\text{LLT polynomial})$$

Sum over all Dyck paths lying below the line $y = -\frac{s}{r}x + s$.

Consequences

- Second uniform proof of shuffle and rational shuffle theorems.

Generalized Shuffle Theorem

Theorem (Blasiak, Haiman, Morse, Pun, S., 2023)

$$D^{(r,s)} \cdot 1 = \sum_{\lambda} (q, t \text{ monomial})(\text{LLT polynomial})$$

Sum over all Dyck paths lying below the line $y = -\frac{s}{r}x + s$.

Consequences

- Second uniform proof of shuffle and rational shuffle theorems.
- Quantity is q, t symmetric, but not clear combinatorially.

Generalized Shuffle Theorem

Theorem (Blasiak, Haiman, Morse, Pun, S., 2023)

$$D^{(r,s)} \cdot 1 = \sum_{\lambda} (q, t \text{ monomial})(\text{LLT polynomial})$$

Sum over all Dyck paths lying below the line $y = -\frac{s}{r}x + s$.

Consequences

- Second uniform proof of shuffle and rational shuffle theorems.
- Quantity is q, t symmetric, but not clear combinatorially.
- Schur expansion coefficients in $\mathbb{N}[q, t]$ (i.e., “Schur-positive”).

Generalized Shuffle Theorem

Theorem (Blasiak, Haiman, Morse, Pun, S., 2023)

$$D^{(r,s)} \cdot 1 = \sum_{\lambda} (q, t \text{ monomial})(\text{LLT polynomial})$$

Sum over all Dyck paths lying below the line $y = -\frac{s}{r}x + s$.

Consequences

- Second uniform proof of shuffle and rational shuffle theorems.
- Quantity is q, t symmetric, but not clear combinatorially.
- Schur expansion coefficients in $\mathbb{N}[q, t]$ (i.e., “Schur-positive”).

Technique

Theory of “Catalanimals” bridging algebra and combinatorics.

Some applications

Some applications

Generalized q, t Catalan numbers

Some applications

Generalized q, t Catalan numbers

- Generalized Shuffle Theorem proof gives single generating function for all “Generalized q, t Catalan numbers” of (Gorsky, Hawkes, Schilling, Rainbolt, 2020)

Some applications

Generalized q, t Catalan numbers

- Generalized Shuffle Theorem proof gives single generating function for all “Generalized q, t Catalan numbers” of (Gorsky, Hawkes, Schilling, Rainbolt, 2020)
- $D^{(r,s)} \cdot 1 = C^{(r,s)}(q, t)s_{(1,\dots,1)} + \dots$

Some applications

Generalized q, t Catalan numbers

- Generalized Shuffle Theorem proof gives single generating function for all “Generalized q, t Catalan numbers” of (Gorsky, Hawkes, Schilling, Rainbolt, 2020)
- $D^{(r,s)} \cdot 1 = C^{(r,s)}(q, t)s_{(1,\dots,1)} + \dots$ ($C^{(k,k)}(q, t) = C_k(q, t)$)

Some applications

Generalized q, t Catalan numbers

- Generalized Shuffle Theorem proof gives single generating function for all “Generalized q, t Catalan numbers” of (Gorsky, Hawkes, Schilling, Rainbolt, 2020)
- $D^{(r,s)} \cdot 1 = C^{(r,s)}(q, t)s_{(1,\dots,1)} + \dots$ ($C^{(k,k)}(q, t) = C_k(q, t)$)

KR-homology (generalizes Jones and Alexander polynomial)

Some applications

Generalized q, t Catalan numbers

- Generalized Shuffle Theorem proof gives single generating function for all “Generalized q, t Catalan numbers” of (Gorsky, Hawkes, Schilling, Rainbolt, 2020)
- $D^{(r,s)} \cdot 1 = C^{(r,s)}(q, t)s_{(1,\dots,1)} + \dots$ ($C^{(k,k)}(q, t) = C_k(q, t)$)


KR-homology (generalizes Jones and Alexander polynomial)

(Capran, González, Hogancamp, Mazin, 2024+) use Generalized Shuffle Theorem formula to give KR homology of family of “Coxeter knots.”

A Mystery: Convex Curve Conjecture

A Mystery: Convex Curve Conjecture

$D^{(C)} \in \mathcal{E}$ for convex curve C between $(0, s)$ and $(r, 0)$.




A Mystery: Convex Curve Conjecture

$D^{(C)} \in \mathcal{E}$ for convex curve C between $(0, s)$ and $(r, 0)$.

Conjecture (Blasiak, Haiman, Morse, Pun, S., 2023)

$D^{(C)} \cdot 1$ is Schur-positive (coefficients in $\mathbb{N}[q, t]$).





A Mystery: Convex Curve Conjecture

$D^{(C)} \in \mathcal{E}$ for convex curve C between $(0, s)$ and $(r, 0)$.

Conjecture (Blasiak, Haiman, Morse, Pun, S., 2023)

$D^{(C)} \cdot 1$ is Schur-positive (coefficients in $\mathbb{N}[q, t]$).





A Mystery: Convex Curve Conjecture

$D^{(C)} \in \mathcal{E}$ for convex curve C between $(0, s)$ and $(r, 0)$.

Conjecture (Blasiak, Haiman, Morse, Pun, S., 2023)

$D^{(C)} \cdot 1$ is Schur-positive (coefficients in $\mathbb{N}[q, t]$).




- S_n -module?

A Mystery: Convex Curve Conjecture

$D^{(C)} \in \mathcal{E}$ for convex curve C between $(0, s)$ and $(r, 0)$.

Conjecture (Blasiak, Haiman, Morse, Pun, S., 2023)

$D^{(C)} \cdot 1$ is Schur-positive (coefficients in $\mathbb{N}[q, t]$).



- S_n -module?
- $s_{(1, \dots, 1)}$ coefficient appears in context of “positroid varieties” (Galashin, Lam, 2024).


Overview

Common theme: $\mathcal{E} \curvearrowright \Lambda = \text{Combinatorial sum}$

Overview

Common theme: $\mathcal{E} \curvearrowright \Lambda = \text{Combinatorial sum}$

$e_k^{(1,1)}$




Shuffle
Theorem
(2018)

Overview


Common theme: $\mathcal{E} \curvearrowright \Lambda = \text{Combinatorial sum}$

$$e_k^{(1,1)}$$



Shuffle
Theorem
(2018)

$$e_k^{(m,n)}$$




Rational
Shuffle Theorem
(2021)

Overview


Common theme: $\mathcal{E} \curvearrowright \Lambda = \text{Combinatorial sum}$

$$e_k^{(1,1)}$$




Shuffle
Theorem
(2018)

$$e_k^{(m,n)}$$



Rational
Shuffle Theorem
(2021)

$$D(r,s)$$




Generalized
Shuffle Theorem
(2023)

Overview


Common theme: $\mathcal{E} \curvearrowright \Lambda = \text{Combinatorial sum}$

$$e_k^{(1,1)}$$




Shuffle
Theorem
(2018)

$$e_k^{(m,n)}$$




Rational
Shuffle Theorem
(2021)

$$D(r,s)$$



Generalized
Shuffle Theorem
(2023)

$$D(C)$$




Convex Curve
Conjecture
(Open)

Overview


Common theme: $\mathcal{E} \curvearrowright \Lambda = \text{Combinatorial sum}$

$$e_k^{(1,1)}$$




Shuffle
Theorem
(2018)

$$e_k^{(m,n)}$$




Rational
Shuffle Theorem
(2021)

$$D(r,s)$$



Generalized
Shuffle Theorem
(2023)

$$D(C)$$



Convex Curve
Conjecture
(Open)

Question

What about $s_\mu^{(m,n)} \in \Lambda^{(m,n)} \subseteq \mathcal{E}$?

The Loehr-Warrington Conjecture

Theorem (BHMPS, 2021+)

$$s_{\mu}^{(m,n)} \cdot 1 = \pm \sum (q, t \text{ monomial})(\text{LLT polynomial})$$

The Loehr-Warrington Conjecture

Theorem (BHMPS, 2021+)

$$s_{\mu}^{(m,n)} \cdot 1 = \pm \sum (q, t \text{ monomial})(\text{LLT polynomial})$$


Sum over “dens” in a “nest.”

The Loehr-Warrington Conjecture

Theorem (BHMPs, 2021+)

$$s_{\mu}^{(m,n)} \cdot 1 = \pm \sum (q, t \text{ monomial})(\text{LLT polynomial})$$

Sum over “dens” in a “nest.”




The Loehr-Warrington Conjecture

Theorem (BHMPs, 2021+)

$$s_{\mu}^{(m,n)} \cdot 1 = \pm \sum (q, t \text{ monomial})(\text{LLT polynomial})$$

Sum over “dens” in a “nest.”



Loehr-Warrington Conjecture (2008)

$$s_{\mu}^{(m,1)} \cdot 1 = \pm \sum (q, t \text{ monomial})(\text{LLT polynomial})$$


Sum over “nested Dyck paths”

Catalanimals

Common theme: $\mathcal{E} \leadsto \Lambda = \text{Combinatorial sum}$


Catalanimals

Common theme: $\mathcal{E} \curvearrowright \Lambda = \text{Combinatorial sum}$

Bridge:  Catalanimals

Catalanimals


Common theme: $\mathcal{E} \curvearrowright \Lambda = \text{Combinatorial sum}$

Bridge:  Catalanimals

- Catalanimal = symmetric multivariate Laurent series.
- Governed by 3 sets of pairs $(i < j) \in \mathbb{N}_+^2$ and $\gamma \in \mathbb{Z}^n$.

Catalanimals

Common theme: $\mathcal{E} \curvearrowright \Lambda = \text{Combinatorial sum}$


Bridge:  Catalanimals

- Catalanimal = symmetric multivariate Laurent series.
- Governed by 3 sets of pairs $(i < j) \in \mathbb{N}_+^2$ and $\gamma \in \mathbb{Z}^n$.

Symmetrization of $\left(\frac{z_1 z_2 z_3 (1 - q t z_1 / z_3)}{\prod_{1 \leq i < j \leq 3} (1 - q z_i / z_j) (1 - t z_i / z_j)} \right)$

Catalanimals

Common theme: $\mathcal{E} \curvearrowright \Lambda = \text{Combinatorial sum}$


Bridge:  Catalanimals

- Catalanimal = symmetric multivariate Laurent series.
- Governed by 3 sets of pairs $(i < j) \in \mathbb{N}_+^2$ and $\gamma \in \mathbb{Z}^n$.

$$\begin{aligned} & \text{Symmetrization of } \left(\frac{z_1 z_2 z_3 (1 - q t z_1 / z_3)}{\prod_{1 \leq i < j \leq 3} (1 - q z_i / z_j) (1 - t z_i / z_j)} \right) \\ &= s_{111} + (q + t + q^2 + q t + t^2) s_{21} + (q t + q^3 + q^2 t + q t^2 + t^3) s_3 \\ & \quad + \dots \end{aligned}$$

Catalanimals

Common theme: $\mathcal{E} \curvearrowright \Lambda = \text{Combinatorial sum}$

Bridge:  Catalanimals

- Catalanimal = symmetric multivariate Laurent series.
- Governed by 3 sets of pairs $(i < j) \in \mathbb{N}_+^2$ and $\gamma \in \mathbb{Z}^n$.

$$\begin{aligned} & \text{Symmetrization of } \left(\frac{z_1 z_2 z_3 (1 - q t z_1 / z_3)}{\prod_{1 \leq i < j \leq 3} (1 - q z_i / z_j) (1 - t z_i / z_j)} \right) \\ &= s_{111} + (q + t + q^2 + q t + t^2) s_{21} + (q t + q^3 + q^2 t + q t^2 + t^3) s_3 \\ & \quad + \dots \end{aligned}$$

Thank you!