

Catalanimals: a walk through the zoo of shuffle theorems

George H. Seelinger

ghseeli@umich.edu

joint with Jonah Blasiak, Mark Haiman, Jennifer Morse, and Anna Pun


University of Pennsylvania Mathematics Colloquium

10 December 2024

Introduction

- 1 Fundamental object: multivariate symmetric polynomials
- 2 Guiding conjecture: The shuffle conjecture
- 3 Recent breakthroughs

Fundamental Object: Symmetric polynomials



Symmetric polynomials

$$x^2 + ax + b = (x - \lambda_1)(x - \lambda_2)$$

Symmetric polynomials

$$\begin{aligned}x^2 + ax + b &= (x - \lambda_1)(x - \lambda_2) \\ a &= -(\lambda_1 + \lambda_2), \quad b = \lambda_1\lambda_2\end{aligned}$$

Symmetric polynomials

$$x^2 + ax + b = (x - \lambda_1)(x - \lambda_2)$$

$$a = -(\lambda_1 + \lambda_2), \quad b = \lambda_1\lambda_2$$

$$\implies \lambda_1^2 + \lambda_2^2$$

Symmetric polynomials

$$x^2 + ax + b = (x - \lambda_1)(x - \lambda_2)$$

$$a = -(\lambda_1 + \lambda_2), \quad b = \lambda_1\lambda_2$$

$$\implies \lambda_1^2 + \lambda_2^2 = (\lambda_1 + \lambda_2)^2 - 2\lambda_1\lambda_2$$

Symmetric polynomials

$$x^2 + ax + b = (x - \lambda_1)(x - \lambda_2)$$

$$a = -(\lambda_1 + \lambda_2), \quad b = \lambda_1\lambda_2$$

$$\implies \lambda_1^2 + \lambda_2^2 = (\lambda_1 + \lambda_2)^2 - 2\lambda_1\lambda_2 = a^2 - 2b$$

Symmetric polynomials

$$x^2 + ax + b = (x - \lambda_1)(x - \lambda_2)$$

$$a = -(\lambda_1 + \lambda_2), \quad b = \lambda_1\lambda_2$$

$$\implies \lambda_1^2 + \lambda_2^2 = (\lambda_1 + \lambda_2)^2 - 2\lambda_1\lambda_2 = a^2 - 2b$$

$x^3 + ax^2 + bx + c$ with roots $\lambda_1, \lambda_2, \lambda_3$.

Symmetric polynomials

$$x^2 + ax + b = (x - \lambda_1)(x - \lambda_2)$$

$$a = -(\lambda_1 + \lambda_2), \quad b = \lambda_1\lambda_2$$

$$\implies \lambda_1^2 + \lambda_2^2 = (\lambda_1 + \lambda_2)^2 - 2\lambda_1\lambda_2 = a^2 - 2b$$

$x^3 + ax^2 + bx + c$ with roots $\lambda_1, \lambda_2, \lambda_3$.

$$\lambda_1^2 + \lambda_2^2 + \lambda_3^2$$

Symmetric polynomials

$$x^2 + ax + b = (x - \lambda_1)(x - \lambda_2)$$

$$a = -(\lambda_1 + \lambda_2), \quad b = \lambda_1\lambda_2$$

$$\implies \lambda_1^2 + \lambda_2^2 = (\lambda_1 + \lambda_2)^2 - 2\lambda_1\lambda_2 = a^2 - 2b$$

$x^3 + ax^2 + bx + c$ with roots $\lambda_1, \lambda_2, \lambda_3$.

$$\lambda_1^2 + \lambda_2^2 + \lambda_3^2 = (\lambda_1 + \lambda_2 + \lambda_3)^2 - 2(\lambda_1\lambda_2 + \lambda_1\lambda_3 + \lambda_2\lambda_3)$$

Symmetric polynomials

$$x^2 + ax + b = (x - \lambda_1)(x - \lambda_2)$$

$$a = -(\lambda_1 + \lambda_2), \quad b = \lambda_1\lambda_2$$

$$\implies \lambda_1^2 + \lambda_2^2 = (\lambda_1 + \lambda_2)^2 - 2\lambda_1\lambda_2 = a^2 - 2b$$

$x^3 + ax^2 + bx + c$ with roots $\lambda_1, \lambda_2, \lambda_3$.

$$\begin{aligned}\lambda_1^2 + \lambda_2^2 + \lambda_3^2 &= (\lambda_1 + \lambda_2 + \lambda_3)^2 - 2(\lambda_1\lambda_2 + \lambda_1\lambda_3 + \lambda_2\lambda_3) \\ &= a^2 - 2b\end{aligned}$$

Symmetric polynomials

$$\begin{aligned}x^2 + ax + b &= (x - \lambda_1)(x - \lambda_2) \\ a &= -(\lambda_1 + \lambda_2), \quad b = \lambda_1\lambda_2 \\ \implies \lambda_1^2 + \lambda_2^2 &= (\lambda_1 + \lambda_2)^2 - 2\lambda_1\lambda_2 = a^2 - 2b\end{aligned}$$

$x^3 + ax^2 + bx + c$ with roots $\lambda_1, \lambda_2, \lambda_3$.

$$\begin{aligned}\lambda_1^2 + \lambda_2^2 + \lambda_3^2 &= (\lambda_1 + \lambda_2 + \lambda_3)^2 - 2(\lambda_1\lambda_2 + \lambda_1\lambda_3 + \lambda_2\lambda_3) \\ &= a^2 - 2b\end{aligned}$$

Observation by Girard and Newton in 1600's

Consider relationships between symmetric polynomials in unspecified (sufficiently large) number of variables.

Symmetric polynomials

$$\begin{aligned}x^2 + ax + b &= (x - \lambda_1)(x - \lambda_2) \\ a &= -(\lambda_1 + \lambda_2), \quad b = \lambda_1\lambda_2 \\ \implies \lambda_1^2 + \lambda_2^2 &= (\lambda_1 + \lambda_2)^2 - 2\lambda_1\lambda_2 = a^2 - 2b\end{aligned}$$

$x^3 + ax^2 + bx + c$ with roots $\lambda_1, \lambda_2, \lambda_3$.

$$\begin{aligned}\lambda_1^2 + \lambda_2^2 + \lambda_3^2 &= (\lambda_1 + \lambda_2 + \lambda_3)^2 - 2(\lambda_1\lambda_2 + \lambda_1\lambda_3 + \lambda_2\lambda_3) \\ &= a^2 - 2b\end{aligned}$$

Observation by Girard and Newton in 1600's

Consider relationships between symmetric polynomials in unspecified (sufficiently large) number of variables.

- e_d = sum of all degree d square-free monomials.

Symmetric polynomials

$$\begin{aligned}x^2 + ax + b &= (x - \lambda_1)(x - \lambda_2) \\ a &= -(\lambda_1 + \lambda_2), \quad b = \lambda_1\lambda_2 \\ \implies \lambda_1^2 + \lambda_2^2 &= (\lambda_1 + \lambda_2)^2 - 2\lambda_1\lambda_2 = a^2 - 2b\end{aligned}$$

$x^3 + ax^2 + bx + c$ with roots $\lambda_1, \lambda_2, \lambda_3$.

$$\begin{aligned}\lambda_1^2 + \lambda_2^2 + \lambda_3^2 &= (\lambda_1 + \lambda_2 + \lambda_3)^2 - 2(\lambda_1\lambda_2 + \lambda_1\lambda_3 + \lambda_2\lambda_3) \\ &= a^2 - 2b\end{aligned}$$

Observation by Girard and Newton in 1600's

Consider relationships between symmetric polynomials in unspecified (sufficiently large) number of variables.

- e_d = sum of all degree d square-free monomials.
- $e_2 = z_1z_2 + z_1z_3 + z_2z_3 + \cdots$

Symmetric polynomials

Linear algebra



Symmetric polynomials

The miraculous basis of Schur polynomials



The miraculous basis of Schur polynomials

$$GL_2(\mathbb{C}) \curvearrowright \mathrm{Sym}^2(\mathbb{C}^2)$$

$$\begin{pmatrix} z_1 & 0 \\ 0 & z_2 \end{pmatrix} \cdot \begin{pmatrix} v_1 \\ v_2 \\ v_3 \end{pmatrix} = \begin{pmatrix} z_1^2 & 0 & 0 \\ 0 & z_1 z_2 & 0 \\ 0 & 0 & z_1^2 \end{pmatrix} \begin{pmatrix} v_1 \\ v_2 \\ v_3 \end{pmatrix}$$

The miraculous basis of Schur polynomials

$$GL_2(\mathbb{C}) \curvearrowright \text{Sym}^2(\mathbb{C}^2)$$

$$\begin{pmatrix} z_1 & 0 \\ 0 & z_2 \end{pmatrix} \cdot \begin{pmatrix} v_1 \\ v_2 \\ v_3 \end{pmatrix} = \begin{pmatrix} z_1^2 & 0 & 0 \\ 0 & z_1 z_2 & 0 \\ 0 & 0 & z_1^2 \end{pmatrix} \begin{pmatrix} v_1 \\ v_2 \\ v_3 \end{pmatrix}$$

Trace = $z_1^2 + z_1 z_2 + z_2^2$ is symmetric!

The miraculous basis of Schur polynomials

$$GL_2(\mathbb{C}) \curvearrowright \text{Sym}^2(\mathbb{C}^2)$$

$$\begin{pmatrix} z_1 & 0 \\ 0 & z_2 \end{pmatrix} \cdot \begin{pmatrix} v_1 \\ v_2 \\ v_3 \end{pmatrix} = \begin{pmatrix} z_1^2 & 0 & 0 \\ 0 & z_1 z_2 & 0 \\ 0 & 0 & z_1^2 \end{pmatrix} \begin{pmatrix} v_1 \\ v_2 \\ v_3 \end{pmatrix}$$

Trace = $z_1^2 + z_1 z_2 + z_2^2$ is symmetric!

Integer partitions

$$\mu = (\mu_1 \geq \dots \geq \mu_n)$$

$$\mu_i \in \mathbb{Z}_{\geq 0}$$



Irreducible (polynomial) $GL_n(\mathbb{C})$
representations V^μ

The miraculous basis of Schur polynomials

$$GL_2(\mathbb{C}) \curvearrowright \text{Sym}^2(\mathbb{C}^2)$$

$$\begin{pmatrix} z_1 & 0 \\ 0 & z_2 \end{pmatrix} \cdot \begin{pmatrix} v_1 \\ v_2 \\ v_3 \end{pmatrix} = \begin{pmatrix} z_1^2 & 0 & 0 \\ 0 & z_1 z_2 & 0 \\ 0 & 0 & z_1^2 \end{pmatrix} \begin{pmatrix} v_1 \\ v_2 \\ v_3 \end{pmatrix}$$

Trace = $z_1^2 + z_1 z_2 + z_2^2$ is symmetric!

Integer partitions

$$\mu = (\mu_1 \geq \dots \geq \mu_n)$$

$$\mu_i \in \mathbb{Z}_{\geq 0}$$



Irreducible (polynomial) $GL_n(\mathbb{C})$
representations V^μ

Trace

Symmetric "Schur polynomial" s_μ

The miraculous basis of Schur polynomials

$$GL_2(\mathbb{C}) \curvearrowright \text{Sym}^2(\mathbb{C}^2)$$

$$\begin{pmatrix} z_1 & 0 \\ 0 & z_2 \end{pmatrix} \cdot \begin{pmatrix} v_1 \\ v_2 \\ v_3 \end{pmatrix} = \begin{pmatrix} z_1^2 & 0 & 0 \\ 0 & z_1 z_2 & 0 \\ 0 & 0 & z_1^2 \end{pmatrix} \begin{pmatrix} v_1 \\ v_2 \\ v_3 \end{pmatrix}$$

Trace = $z_1^2 + z_1 z_2 + z_2^2$ is symmetric!

Integer partitions

$$\mu = (\mu_1 \geq \dots \geq \mu_n)$$

$$\mu_i \in \mathbb{Z}_{\geq 0}$$




Irreducible (polynomial) $GL_n(\mathbb{C})$
representations V^μ

Trace


Symmetric "Schur polynomial" s_μ

$$s_{(2,0)}(z_1, z_2) = z_1^2 + z_1 z_2 + z_2^2.$$

The miraculous basis of Schur polynomials




The miraculous basis of Schur polynomials



Question

$f = z_1^2 + z_2^2$ the trace of some $GL_2(\mathbb{C})$ representation?

The miraculous basis of Schur polynomials




Question

$f = z_1^2 + z_2^2$ the trace of some $GL_2(\mathbb{C})$ representation?

Solution

$\{s_\mu\}_\mu$ is a basis of symmetric polynomials.

The miraculous basis of Schur polynomials



Question


$f = z_1^2 + z_2^2$ the trace of some $GL_2(\mathbb{C})$ representation?

Solution

$\{s_\mu\}_\mu$ is a basis of symmetric polynomials.

$$f = s_{(2,0)} - s_{(1,1)}$$

The miraculous basis of Schur polynomials



Question

$f = z_1^2 + z_2^2$ the trace of some $GL_2(\mathbb{C})$ representation?


Solution

$\{s_\mu\}_\mu$ is a basis of symmetric polynomials.

$$f = s_{(2,0)} - s_{(1,1)}$$

f not "Schur-positive" \implies not the trace of a GL_2 -representation!

The miraculous basis of Schur polynomials



Question

$f = z_1^2 + z_2^2$ the trace of some $GL_2(\mathbb{C})$ representation?

Solution

$\{s_\mu\}_\mu$ is a basis of symmetric polynomials.

$$f = s_{(2,0)} - s_{(1,1)}$$

f not "Schur-positive" \implies not the trace of a GL_2 -representation!

Guiding principle

Positive integer sums of Schurs \implies representation-theoretically meaningful.

The miraculous basis of Schur polynomials

$$GL_n(\mathbb{C}) \curvearrowright V^\mu$$

Trace
↓

Symmetric “Schur polynomial” s_μ

$$s_\mu = \sum x^{\text{wt}(T)}$$

The miraculous basis of Schur polynomials

$$GL_n(\mathbb{C}) \curvearrowright V^\mu$$

Trace


Symmetric "Schur polynomial" s_μ

$$s_\mu = \sum x^{\text{wt}(T)}$$

Sum of weights

Semistandard Young Tableaux

$$\mu = (4, 3, 1, 1) \rightarrow$$



$$\rightarrow T =$$

| | | | |
|---|---|---|---|
| 5 | | | |
| 3 | | | |
| 2 | 3 | 5 | |
| 1 | 1 | 2 | 3 |

The miraculous basis of Schur polynomials

$$GL_n(\mathbb{C}) \curvearrowright V^\mu$$

Trace

Symmetric "Schur polynomial" s_μ

$$s_\mu = \sum x^{\text{wt}(T)}$$

Sum of weights

Semistandard Young Tableaux

$$\mu = (4, 3, 1, 1) \rightarrow \begin{array}{|c|c|c|c|} \hline & & & \\ \hline & & & \\ \hline & & & \\ \hline & & & \\ \hline \end{array} \rightarrow T = \begin{array}{|c|c|c|c|} \hline 5 & & & \\ \hline 3 & & & \\ \hline 2 & 3 & 5 & \\ \hline 1 & 1 & 2 & 3 \\ \hline \end{array}$$

Consequence for irreducible V^μ

$\dim V^\mu = \text{number of semistandard Young tableaux of } \mu.$

S_n -modules

Symmetric group $S_n \curvearrowright \mathbb{C}[x_1, \dots, x_n]$

S_n -modules

Symmetric group $S_n \curvearrowright \mathbb{C}[x_1, \dots, x_n]$



$$S_n \curvearrowright M = \mathbb{C}[x_1, \dots, x_n]/I$$

I = all positive degree symmetric polynomials

S_n -modules

Symmetric group $S_n \curvearrowright \mathbb{C}[x_1, \dots, x_n]$



$S_n \curvearrowright M = \mathbb{C}[x_1, \dots, x_n]/I$
 $I = \text{all positive degree symmetric polynomials}$

\cong

$H^*(Fl_n(\mathbb{C}))$

S_n -modules

Symmetric group $S_n \curvearrowright \mathbb{C}[x_1, \dots, x_n]$



$S_n \curvearrowright M = \mathbb{C}[x_1, \dots, x_n]/I$
 $I =$ all positive degree symmetric polynomials

\cong

$H^*(Fl_n(\mathbb{C}))$



Harmonic polynomials \mathcal{H}_n

S_n -modules

Symmetric group $S_n \curvearrowright \mathbb{C}[x_1, \dots, x_n]$



$S_n \curvearrowright M = \mathbb{C}[x_1, \dots, x_n]/I$
 $I =$ all positive degree symmetric polynomials

\cong

$H^*(Fl_n(\mathbb{C}))$



Harmonic polynomials \mathcal{H}_n

$$f(x_1, x_2) \in \mathcal{H}_2 \iff \begin{cases} (\partial_{x_1} + \partial_{x_2})f = 0 \\ \partial_{x_1} \partial_{x_2} f = 0 \end{cases} \implies \mathcal{H}_2 = \langle x_1 - x_2 \rangle \oplus \langle 1 \rangle$$

S_n -modules

Symmetric group $S_n \curvearrowright \mathbb{C}[x_1, \dots, x_n]$

$S_n \curvearrowright M = \mathbb{C}[x_1, \dots, x_n]/I$
 $I = \text{all positive degree symmetric polynomials}$

\cong

$H^*(Fl_n(\mathbb{C}))$

\cong

Harmonic polynomials \mathcal{H}_n

$$f(x_1, x_2) \in \mathcal{H}_2 \iff \begin{cases} (\partial_{x_1} + \partial_{x_2})f = 0 \\ \partial_{x_1}\partial_{x_2}f = 0 \end{cases} \implies \mathcal{H}_2 = \langle x_1 - x_2 \rangle \oplus \langle 1 \rangle$$

Question

What are the irreducible (S_n -invariant) pieces of \mathcal{H}_n ?

S_n -modules

Symmetric group $S_n \curvearrowright \mathbb{C}[x_1, \dots, x_n]$

$S_n \curvearrowright M = \mathbb{C}[x_1, \dots, x_n]/I$
 $I = \text{all positive degree symmetric polynomials}$

\cong

$H^*(Fl_n(\mathbb{C}))$

\cong

Harmonic polynomials \mathcal{H}_n

$$f(x_1, x_2) \in \mathcal{H}_2 \iff \begin{cases} (\partial_{x_1} + \partial_{x_2})f = 0 \\ \partial_{x_1}\partial_{x_2}f = 0 \end{cases} \implies \mathcal{H}_2 = \langle x_1 - x_2 \rangle \oplus \langle 1 \rangle$$

Question

What are the irreducible (S_n -invariant) pieces of \mathcal{H}_n ?

Frobenius characteristic

S_n -representation $V \mapsto \text{symmetric polynomial } \text{Frob}(V)$.

The miraculous basis of Schur polynomials

Integer partition of $n \leftrightarrow$ irreducible S_n representation V_μ

The miraculous basis of Schur polynomials

Integer partition of $n \leftrightarrow$ irreducible S_n representation V_μ

$$S_n \curvearrowright V_\mu$$

The miraculous basis of Schur polynomials

Integer partition of $n \leftrightarrow$ irreducible S_n representation V_μ

$$S_n \curvearrowright V_\mu$$

Frob

Schur polynomial s_μ

The miraculous basis of Schur polynomials

Integer partition of $n \leftrightarrow$ irreducible S_n representation V_μ

$$S_n \curvearrowright V_\mu$$

Frob

$$GL_n(\mathbb{C}) \curvearrowright V^\mu$$

Trace

Schur polynomial s_μ

Sum of weights

Semistandard Young Tableaux

| | | | | |
|---|---|---|---|--|
| 5 | | | | |
| 3 | | | | |
| 2 | 3 | 5 | | |
| 1 | 1 | 2 | 3 | |

S_n -module of Harmonic polynomials

$$\mathcal{H}_2 \cong \langle x_1 - x_2 \rangle \oplus \langle 1 \rangle$$

$$\text{Frob}(\mathcal{H}_2) = s_{(1,1)} + s_{(2)}$$

S_n -module of Harmonic polynomials

$$\mathcal{H}_2 \cong \langle x_1 - x_2 \rangle \oplus \langle 1 \rangle$$
$$\text{Frob}(\mathcal{H}_2) = s_{(1,1)} + s_{(2)}$$

Identification ignores polynomial degrees \implies add parameter q .

S_n -module of Harmonic polynomials

$$\mathcal{H}_2 \cong \langle x_1 - x_2 \rangle \oplus \langle 1 \rangle$$
$$\text{Frob}(\mathcal{H}_2) = s_{(1,1)} + s_{(2)}$$

Identification ignores polynomial degrees \implies add parameter q .

$$\mathcal{H}_2 \cong \langle x_1 - x_2 \rangle \oplus \langle 1 \rangle$$
$$\text{Frob}_q(\mathcal{H}_2) = qs_{(1,1)} + s_{(2)}$$

S_n -module of Harmonic polynomials

$$\mathcal{H}_2 \cong \langle x_1 - x_2 \rangle \oplus \langle 1 \rangle$$
$$\text{Frob}(\mathcal{H}_2) = s_{(1,1)} + s_{(2)}$$

Identification ignores polynomial degrees \implies add parameter q .

$$\mathcal{H}_2 \cong \langle x_1 - x_2 \rangle \oplus \langle 1 \rangle$$
$$\text{Frob}_q(\mathcal{H}_2) = qs_{(1,1)} + s_{(2)}$$

$$\text{Frob}(\mathcal{H}_3) = s_{(1,1,1)} + 2s_{(2,1)} + s_{(3)}$$
$$\text{Frob}_q(\mathcal{H}_3) = q^3s_{(1,1,1)} + (q^2 + q)s_{(2,1)} + s_{(3)}$$

S_n -module of Harmonic polynomials

$$\mathcal{H}_2 \cong \langle x_1 - x_2 \rangle \oplus \langle 1 \rangle$$
$$\text{Frob}(\mathcal{H}_2) = s_{(1,1)} + s_{(2)}$$

Identification ignores polynomial degrees \implies add parameter q .

$$\mathcal{H}_2 \cong \langle x_1 - x_2 \rangle \oplus \langle 1 \rangle$$
$$\text{Frob}_q(\mathcal{H}_2) = qs_{(1,1)} + s_{(2)}$$

$$\text{Frob}(\mathcal{H}_3) = s_{(1,1,1)} + 2s_{(2,1)} + s_{(3)}$$
$$\text{Frob}_q(\mathcal{H}_3) = q^3 s_{(1,1,1)} + (q^2 + q)s_{(2,1)} + s_{(3)}$$

Problems

- What is $\text{Frob}_q(\mathcal{H}_n)$?

S_n -module of Harmonic polynomials

$$\mathcal{H}_2 \cong \langle x_1 - x_2 \rangle \oplus \langle 1 \rangle$$
$$\text{Frob}(\mathcal{H}_2) = s_{(1,1)} + s_{(2)}$$

Identification ignores polynomial degrees \implies add parameter q .

$$\mathcal{H}_2 \cong \langle x_1 - x_2 \rangle \oplus \langle 1 \rangle$$
$$\text{Frob}_q(\mathcal{H}_2) = qs_{(1,1)} + s_{(2)}$$

$$\text{Frob}(\mathcal{H}_3) = s_{(1,1,1)} + 2s_{(2,1)} + s_{(3)}$$
$$\text{Frob}_q(\mathcal{H}_3) = q^3 s_{(1,1,1)} + (q^2 + q)s_{(2,1)} + s_{(3)}$$

Problems

- What is $\text{Frob}_q(\mathcal{H}_n)$? A “Hall-Littlewood polynomial.”

S_n -module of Harmonic polynomials

$$\mathcal{H}_2 \cong \langle x_1 - x_2 \rangle \oplus \langle 1 \rangle$$
$$\text{Frob}(\mathcal{H}_2) = s_{(1,1)} + s_{(2)}$$

Identification ignores polynomial degrees \implies add parameter q .

$$\mathcal{H}_2 \cong \langle x_1 - x_2 \rangle \oplus \langle 1 \rangle$$
$$\text{Frob}_q(\mathcal{H}_2) = qs_{(1,1)} + s_{(2)}$$

$$\text{Frob}(\mathcal{H}_3) = s_{(1,1,1)} + 2s_{(2,1)} + s_{(3)}$$
$$\text{Frob}_q(\mathcal{H}_3) = q^3 s_{(1,1,1)} + (q^2 + q)s_{(2,1)} + s_{(3)}$$

Problems

- What is $\text{Frob}_q(\mathcal{H}_n)$? A “Hall-Littlewood polynomial.”
- Combinatorics for coefficients?

S_n -module of Harmonic polynomials

$$\mathcal{H}_2 \cong \langle x_1 - x_2 \rangle \oplus \langle 1 \rangle$$
$$\text{Frob}(\mathcal{H}_2) = s_{(1,1)} + s_{(2)}$$

Identification ignores polynomial degrees \implies add parameter q .

$$\mathcal{H}_2 \cong \langle x_1 - x_2 \rangle \oplus \langle 1 \rangle$$
$$\text{Frob}_q(\mathcal{H}_2) = qs_{(1,1)} + s_{(2)}$$

$$\text{Frob}(\mathcal{H}_3) = s_{(1,1,1)} + 2s_{(2,1)} + s_{(3)}$$
$$\text{Frob}_q(\mathcal{H}_3) = q^3 s_{(1,1,1)} + (q^2 + q)s_{(2,1)} + s_{(3)}$$

Problems

- What is $\text{Frob}_q(\mathcal{H}_n)$? A “Hall-Littlewood polynomial.”
- Combinatorics for coefficients? Lascoux-Schutzenberger

Natural generalization: Diagonal Harmonics

Symmetric group $S_n \curvearrowright \mathbb{C}[x_1, \dots, x_n, y_1, \dots, y_n]$

Natural generalization: Diagonal Harmonics

Symmetric group $S_n \curvearrowright \mathbb{C}[x_1, \dots, x_n, y_1, \dots, y_n]$



$$S_n \curvearrowright M = \mathbb{C}[x_1, \dots, x_n, y_1, \dots, y_n] / I$$

I = all positive degree polynomials symmetric in x_i 's and in y_i 's

Natural generalization: Diagonal Harmonics

Symmetric group $S_n \curvearrowright \mathbb{C}[x_1, \dots, x_n, y_1, \dots, y_n]$



$$S_n \curvearrowright M = \mathbb{C}[x_1, \dots, x_n, y_1, \dots, y_n]/I$$

$I =$ all positive degree polynomials symmetric in x_i 's and in y_i 's



Diagonal harmonics \mathcal{H}_n

Natural generalization: Diagonal Harmonics

Symmetric group $S_n \curvearrowright \mathbb{C}[x_1, \dots, x_n, y_1, \dots, y_n]$



$$S_n \curvearrowright M = \mathbb{C}[x_1, \dots, x_n, y_1, \dots, y_n]/I$$

I = all positive degree polynomials symmetric in x_i 's and in y_i 's



Diagonal harmonics \mathcal{H}_n

$$D\mathcal{H}_2 = \langle x_1 - x_2 \rangle \oplus \langle y_1 - y_2 \rangle \oplus \langle 1 \rangle$$

$$\text{Frob}_{q,t}(D\mathcal{H}_2) = qs_{(1,1)} + ts_{(1,1)} + s_{(2)}$$

Natural generalization: Diagonal Harmonics

Symmetric group $S_n \curvearrowright \mathbb{C}[x_1, \dots, x_n, y_1, \dots, y_n]$



$$S_n \curvearrowright M = \mathbb{C}[x_1, \dots, x_n, y_1, \dots, y_n] / I$$

$I =$ all positive degree polynomials symmetric in x_i 's and in y_i 's



Diagonal harmonics \mathcal{H}_n

$$D\mathcal{H}_2 = \langle x_1 - x_2 \rangle \oplus \langle y_1 - y_2 \rangle \oplus \langle 1 \rangle$$

$$\text{Frob}_{q,t}(D\mathcal{H}_2) = qs_{(1,1)} + ts_{(1,1)} + s_{(2)}$$

Foundational unsolved question

What are the combinatorics for this expression?

Natural generalization: Diagonal Harmonics

Symmetric group $S_n \curvearrowright \mathbb{C}[x_1, \dots, x_n, y_1, \dots, y_n]$



$$S_n \curvearrowright M = \mathbb{C}[x_1, \dots, x_n, y_1, \dots, y_n] / I$$

$I =$ all positive degree polynomials symmetric in x_i 's and in y_i 's



Diagonal harmonics \mathcal{H}_n

$$D\mathcal{H}_2 = \langle x_1 - x_2 \rangle \oplus \langle y_1 - y_2 \rangle \oplus \langle 1 \rangle$$

$$\text{Frob}_{q,t}(D\mathcal{H}_2) = qs_{(1,1)} + ts_{(1,1)} + s_{(2)}$$

Foundational unsolved question

What are the combinatorics for this expression?

$$\text{Frob}_{q,t}(D\mathcal{H}_3) = (q^3 + q^2t + qt^2 + t^3 + qt)s_{111} + (q^2 + qt + t^2 + q + t)s_{21} + s_3$$


The fluttering of a combinatorialist's heart

What do the following objects have in common?

The fluttering of a combinatorialist's heart

What do the following objects have in common?


- Dyck paths



The fluttering of a combinatorialist's heart

What do the following objects have in common?

- Dyck paths




- Polygon triangulations



The fluttering of a combinatorialist's heart

What do the following objects have in common?

- Dyck paths



- Polygon triangulations



- The number of $2 \times n$ rectangle standard Young tableaux

| | | |
|---|---|---|
| 4 | 5 | 6 |
| 1 | 2 | 3 |

| | | |
|---|---|---|
| 3 | 4 | 6 |
| 1 | 2 | 5 |

| | | |
|---|---|---|
| 3 | 5 | 6 |
| 1 | 2 | 4 |


| | | |
|---|---|---|
| 2 | 5 | 6 |
| 1 | 3 | 4 |

| | | |
|---|---|---|
| 2 | 4 | 6 |
| 1 | 3 | 5 |

The fluttering of a combinatorialist's heart

What do the following objects have in common?

- Dyck paths



- Polygon triangulations



- The number of $2 \times n$ rectangle standard Young tableaux

| | | |
|---|---|---|
| 4 | 5 | 6 |
| 1 | 2 | 3 |

| | | |
|---|---|---|
| 3 | 4 | 6 |
| 1 | 2 | 5 |

| | | |
|---|---|---|
| 3 | 5 | 6 |
| 1 | 2 | 4 |

| | | |
|---|---|---|
| 2 | 5 | 6 |
| 1 | 3 | 4 |

| | | |
|---|---|---|
| 2 | 4 | 6 |
| 1 | 3 | 5 |


- The number of independent sign subrepresentations of $D\mathcal{H}_n$?

$$(q^3 + q^2t + qt^2 + t^3 + qt)s_{111} + \cdots$$

The fluttering of a combinatorialist's heart

What do the following objects have in common?

- Dyck paths



- Polygon triangulations



- The number of $2 \times n$ rectangle standard Young tableaux

| | | |
|---|---|---|
| 4 | 5 | 6 |
| 1 | 2 | 3 |

| | | |
|---|---|---|
| 3 | 4 | 6 |
| 1 | 2 | 5 |

| | | |
|---|---|---|
| 3 | 5 | 6 |
| 1 | 2 | 4 |

| | | |
|---|---|---|
| 2 | 5 | 6 |
| 1 | 3 | 4 |

| | | |
|---|---|---|
| 2 | 4 | 6 |
| 1 | 3 | 5 |

- The number of independent sign subrepresentations of $D\mathcal{H}_n$?


$$(q^3 + q^2t + qt^2 + t^3 + qt)s_{111} + \cdots$$

Catalan numbers!

The fluttering of a combinatorialist's heart

What do the following objects have in common?

- Dyck paths



- Polygon triangulations



- The number of $2 \times n$ rectangle standard Young tableaux

| | | |
|---|---|---|
| 4 | 5 | 6 |
| 1 | 2 | 3 |

| | | |
|---|---|---|
| 3 | 4 | 6 |
| 1 | 2 | 5 |

| | | |
|---|---|---|
| 3 | 5 | 6 |
| 1 | 2 | 4 |

| | | |
|---|---|---|
| 2 | 5 | 6 |
| 1 | 3 | 4 |

| | | |
|---|---|---|
| 2 | 4 | 6 |
| 1 | 3 | 5 |

- The number of independent sign subrepresentations of $D\mathcal{H}_n$?

$$(q^3 + q^2t + qt^2 + t^3 + qt)s_{111} + \cdots$$

Catalan numbers! 1, 2, 5, 14, ...

Special case

$$\text{Frob}_{q,t}(D\mathcal{H}_n)|_{q=1} = \sum_{\lambda \in DP_n} (t \text{ monomial})(\text{product of } e_i\text{'s})$$

Special case

$$\text{Frob}_{q,t}(D\mathcal{H}_n)|_{q=1} = \sum_{\lambda \in DP_n} (t \text{ monomial})(\text{product of } e_i\text{'s})$$

$$\lambda \quad (t \text{ monomial})(\text{product of } e_i\text{'s})$$

Special case

$$\text{Frob}_{q,t}(D\mathcal{H}_n)|_{q=1} = \sum_{\lambda \in DP_n} (t \text{ monomial})(\text{product of } e_i\text{'s})$$

λ $(t \text{ monomial})(\text{product of } e_i\text{'s})$



Special case

$$\text{Frob}_{q,t}(D\mathcal{H}_n)|_{q=1} = \sum_{\lambda \in DP_n} (t \text{ monomial})(\text{product of } e_i\text{'s})$$

λ $(t \text{ monomial})(\text{product of } e_i\text{'s})$



$$e_1^3$$



$$te_2e_1$$



$$te_2e_1$$



$$t^2e_2e_1$$



$$t^3e_3$$

Special case

$$\text{Frob}_{q,t}(D\mathcal{H}_n)|_{q=1} = \sum_{\lambda \in DP_n} (t \text{ monomial})(\text{product of } e_i\text{'s})$$

λ $(t \text{ monomial})(\text{product of } e_i\text{'s})$



$$e_1^3$$



$$te_2e_1$$



$$te_2e_1$$



$$t^2e_2e_1$$



$$t^3e_3$$

$$\implies \text{Frob}_{q,t}(D\mathcal{H}_n)|_{q=1} = e_1^3 + (t^2 + 2t)e_2e_1 + t^3e_3$$

The Shuffle Theorem

The Shuffle Theorem (Carlsson-Mellit, 2018)

$$\mathrm{Frob}_{q,t}(D\mathcal{H}_n) =$$

The Shuffle Theorem

The Shuffle Theorem (Carlsson-Mellit, 2018)

$$\text{Frob}_{q,t}(D\mathcal{H}_n) = \sum_{\lambda \in DP_n} (q, t \text{ monomial})(\text{symmetric polynomial})$$

The Shuffle Theorem

The Shuffle Theorem (Carlsson-Mellit, 2018)

$$\text{Frob}_{q,t}(D\mathcal{H}_n) = \sum_{\lambda \in DP_n} (q, t \text{ monomial})(\text{symmetric polynomial})$$

- Conjectured by (Haglund-Haiman-Loehr-Remmel-Ulyanov, 2005)

The Shuffle Theorem

The Shuffle Theorem (Carlsson-Mellit, 2018)

$$\text{Frob}_{q,t}(D\mathcal{H}_n) = \sum_{\lambda \in DP_n} (q, t \text{ monomial})(\text{symmetric polynomial})$$

- Conjectured by (Haglund-Haiman-Loehr-Remmel-Ulyanov, 2005)
- Symmetric polynomial = “LLT polynomial” (Lascoux, Leclerc, Thibon, 1997)

The Shuffle Theorem

The Shuffle Theorem (Carlsson-Mellit, 2018)

$$\text{Frob}_{q,t}(D\mathcal{H}_n) = \sum_{\lambda \in DP_n} (q, t \text{ monomial})(\text{symmetric polynomial})$$

- Conjectured by (Haglund-Haiman-Loehr-Remmel-Ulyanov, 2005)
- Symmetric polynomial = “LLT polynomial” (Lascoux, Leclerc, Thibon, 1997)
- Note LLT polynomial = $\sum c_\mu(q) s_\mu$ has $c_\mu(q) \in \mathbb{N}[q]$ (Grojnowski, Haiman, 2007).

The Shuffle Theorem

The Shuffle Theorem (Carlsson-Mellit, 2018)

$$\text{Frob}_{q,t}(D\mathcal{H}_n) = \sum_{\lambda \in DP_n} (q, t \text{ monomial})(\text{symmetric polynomial})$$

- Conjectured by (Haglund-Haiman-Loehr-Remmel-Ulyanov, 2005)
- Symmetric polynomial = “LLT polynomial” (Lascoux, Leclerc, Thibon, 1997)
- Note LLT polynomial = $\sum c_\mu(q) s_\mu$ has $c_\mu(q) \in \mathbb{N}[q]$ (Grojnowski, Haiman, 2007).

q, t Catalan numbers

$$C_n(q, t) = \sum_{\lambda \in DP_n} (q, t \text{ monomial})$$

Generalizing with the Elliptic Hall Algebra

Algebra $\mathcal{E} \curvearrowright \Lambda =$ symmetric polynomials

Generalizing with the Elliptic Hall Algebra

Algebra $\mathcal{E} \curvearrowright \Lambda =$ symmetric polynomials

$\mathcal{E} =$ Hall algebra of Coherent sheaves
on an elliptic curve

Generalizing with the Elliptic Hall Algebra

Algebra $\mathcal{E} \curvearrowright \Lambda =$ symmetric polynomials

$\mathcal{E} =$ Hall algebra of Coherent sheaves
on an elliptic curve

$$\mathcal{E} \cong \text{central subalgebra} \bigoplus_{m,n \text{ coprime}} \bigoplus \Lambda^{(m,n)} \quad \Lambda^{(m,n)} \cong \text{symmetric polynomials}$$

Generalizing with the Elliptic Hall Algebra

Algebra $\mathcal{E} \curvearrowright \Lambda =$ symmetric polynomials

$\mathcal{E} =$ Hall algebra of Coherent sheaves
on an elliptic curve

$$\mathcal{E} \cong \text{central subalgebra} \oplus \bigoplus_{m,n \text{ coprime}} \bigoplus \Lambda^{(m,n)} \quad \Lambda^{(m,n)} \cong \text{symmetric polynomials}$$

Theorem (Haiman, 2002)

$e_k^{(1,1)} \in \Lambda^{(1,1)}$ acts by $e_k^{(1,1)} \cdot 1 = \text{Frob}_{q,t}(D\mathcal{H}_k)$ via connections to $\text{Hilb}^n(\mathbb{C}^2)$.

Generalizing with the Elliptic Hall Algebra

Algebra $\mathcal{E} \curvearrowright \Lambda =$ symmetric polynomials

$\mathcal{E} =$ Hall algebra of Coherent sheaves
on an elliptic curve

$$\mathcal{E} \cong \text{central subalgebra} \bigoplus_{m,n \text{ coprime}} \bigoplus \Lambda^{(m,n)} \quad \Lambda^{(m,n)} \cong \text{symmetric polynomials}$$

Theorem (Haiman, 2002)

$e_k^{(1,1)} \in \Lambda^{(1,1)}$ acts by $e_k^{(1,1)} \cdot 1 = \text{Frob}_{q,t}(D\mathcal{H}_k)$ via connections to $\text{Hilb}^n(\mathbb{C}^2)$.

Question

What can we say about $e_k^{(m,n)} \cdot 1$?

Rational Shuffle Theorem

Rational Shuffle Theorem (Mellit, 2021)

$$e_k^{(m,n)} \cdot 1 = \sum (q, t \text{ monomial})(\text{symmetric LLT polynomial})$$

Rational Shuffle Theorem

Rational Shuffle Theorem (Mellit, 2021)

$$e_k^{(m,n)} \cdot 1 = \sum (q, t \text{ monomial})(\text{symmetric LLT polynomial})$$

- Conjectured by (F. Bergeron, Garsia, Sergel Leven, Xin, 2016)

Rational Shuffle Theorem

Rational Shuffle Theorem (Mellit, 2021)

$$e_k^{(m,n)} \cdot 1 = \sum (q, t \text{ monomial})(\text{symmetric LLT polynomial})$$

- Conjectured by (F. Bergeron, Garsia, Sergel Leven, Xin, 2016)
- Sum over $km \times kn$ Dyck paths

Rational Shuffle Theorem

Rational Shuffle Theorem (Mellit, 2021)

$$e_k^{(m,n)} \cdot 1 = \sum (q, t \text{ monomial})(\text{symmetric LLT polynomial})$$


- Conjectured by (F. Bergeron, Garsia, Sergel Leven, Xin, 2016)
- Sum over $km \times kn$ Dyck paths
- Coefficient of $s_{(1,\dots,1)} \implies$ Rational q, t Catalan numbers.

Rational Shuffle Theorem

Rational Shuffle Theorem (Mellit, 2021)

$$e_k^{(m,n)} \cdot 1 = \sum (q, t \text{ monomial})(\text{symmetric LLT polynomial})$$

- Conjectured by (F. Bergeron, Garsia, Sergel Leven, Xin, 2016)
- Sum over $km \times kn$ Dyck paths
- Coefficient of $s_{(1,\dots,1)} \implies$ Rational q, t Catalan numbers.



Generalized Shuffle Theorem

Generalized Shuffle Theorem

Negut

$r, s \in \mathbb{R}_{>0} \rightarrow D^{(r,s)} \in \mathcal{E}$ such that $D^{(km, kn)} = e_k^{(m,n)}$.

Generalized Shuffle Theorem

Negut

$r, s \in \mathbb{R}_{>0} \rightarrow D^{(r,s)} \in \mathcal{E}$ such that $D^{(km, kn)} = e_k^{(m,n)}$.

Theorem (Blasiak, Haiman, Morse, Pun, S., 2023)

$D^{(r,s)} \cdot 1 = \sum (q, t \text{ monomial})(\text{symmetric LLT polynomial})$

Generalized Shuffle Theorem


Negut

$r, s \in \mathbb{R}_{>0} \rightarrow D^{(r,s)} \in \mathcal{E}$ such that $D^{(km, kn)} = e_k^{(m,n)}$.

Theorem (Blasiak, Haiman, Morse, Pun, S., 2023)

$D^{(r,s)} \cdot 1 = \sum (q, t \text{ monomial})(\text{symmetric LLT polynomial})$

Sum over all SE lattice paths lying below the line $y = -\frac{s}{r}x + s$.



Generalized Shuffle Theorem

Theorem (Blasiak, Haiman, Morse, Pun, S., 2023)

$$D^{(r,s)} \cdot 1 = \sum_{\lambda} (q, t \text{ monomial})(\text{symmetric function})$$

Sum over all SE lattice paths lying below the line $y = -\frac{s}{r}x + s$.

Generalized Shuffle Theorem

Theorem (Blasiak, Haiman, Morse, Pun, S., 2023)

$$D^{(r,s)} \cdot 1 = \sum_{\lambda} (q, t \text{ monomial})(\text{symmetric function})$$

Sum over all SE lattice paths lying below the line $y = -\frac{s}{r}x + s$.

Consequences

- Gives second uniform proof of shuffle and rational shuffle theorems.

Generalized Shuffle Theorem

Theorem (Blasiak, Haiman, Morse, Pun, S., 2023)

$$D^{(r,s)} \cdot 1 = \sum_{\lambda} (q, t \text{ monomial})(\text{symmetric function})$$

Sum over all SE lattice paths lying below the line $y = -\frac{s}{r}x + s$.

Consequences

- Gives second uniform proof of shuffle and rational shuffle theorems.
- Entire quantity is q, t symmetric, but not clear combinatorially.

Generalized Shuffle Theorem

Theorem (Blasiak, Haiman, Morse, Pun, S., 2023)

$$D^{(r,s)} \cdot 1 = \sum_{\lambda} (q, t \text{ monomial})(\text{symmetric function})$$

Sum over all SE lattice paths lying below the line $y = -\frac{s}{r}x + s$.

Consequences

- Gives second uniform proof of shuffle and rational shuffle theorems.
- Entire quantity is q, t symmetric, but not clear combinatorially.
- Schur expansion coefficients in $\mathbb{N}[q, t]$

Generalized Shuffle Theorem

Theorem (Blasiak, Haiman, Morse, Pun, S., 2023)

$$D^{(r,s)} \cdot 1 = \sum_{\lambda} (q, t \text{ monomial})(\text{symmetric function})$$

Sum over all SE lattice paths lying below the line $y = -\frac{s}{r}x + s$.

Consequences

- Gives second uniform proof of shuffle and rational shuffle theorems.
- Entire quantity is q, t symmetric, but not clear combinatorially.
- Schur expansion coefficients in $\mathbb{N}[q, t]$
- \rightarrow Some S_n -module M such that $\text{Frob}_{q,t}(M) = D^{(r,s)} \cdot 1$?

Some applications

Some applications

Generalized q, t Catalan numbers

Some applications

Generalized q, t Catalan numbers

- $D^{(r,s)} \cdot 1 = C^{(r,s)}(q, t)s_{(1,\dots,1)} + \dots$

Some applications

Generalized q, t Catalan numbers

- $D^{(r,s)} \cdot 1 = C^{(r,s)}(q, t)s_{(1,\dots,1)} + \dots$
- $C^{(r,s)}(q, t)$ = subclass of “generalized q, t Catalan numbers” of (Gorsky, Hawkes, Schilling, Rainbolt, 2020).

Some applications

Generalized q, t Catalan numbers

- $D^{(r,s)} \cdot 1 = C^{(r,s)}(q, t)s_{(1,\dots,1)} + \dots$
- $C^{(r,s)}(q, t)$ = subclass of “generalized q, t Catalan numbers” of (Gorsky, Hawkes, Schilling, Rainbolt, 2020).
- Generalized Shuffle Theorem proof gives all generalized q, t Catalan numbers as coefficients of a single geometric series or “generating function.”

Some applications

Generalized q, t Catalan numbers

- $D^{(r,s)} \cdot 1 = C^{(r,s)}(q, t)s_{(1,\dots,1)} + \dots$
- $C^{(r,s)}(q, t)$ = subclass of “generalized q, t Catalan numbers” of (Gorsky, Hawkes, Schilling, Rainbolt, 2020).
- Generalized Shuffle Theorem proof gives all generalized q, t Catalan numbers as coefficients of a single geometric series or “generating function.”

KR-homology (generalizes HOMFLY-PT polynomial)

Some applications

Generalized q, t Catalan numbers

- $D^{(r,s)} \cdot 1 = C^{(r,s)}(q, t)s_{(1,\dots,1)} + \dots$
- $C^{(r,s)}(q, t)$ = subclass of “generalized q, t Catalan numbers” of (Gorsky, Hawkes, Schilling, Rainbolt, 2020).
- Generalized Shuffle Theorem proof gives all generalized q, t Catalan numbers as coefficients of a single geometric series or “generating function.”

KR-homology (generalizes HOMFLY-PT polynomial)

- Rational Shuffle Theorem \implies formula for KR homology of (m, n) -torus knots. (Mellit, 2021)

Some applications

Generalized q, t Catalan numbers

- $D^{(r,s)} \cdot 1 = C^{(r,s)}(q, t)s_{(1,\dots,1)} + \dots$
- $C^{(r,s)}(q, t)$ = subclass of “generalized q, t Catalan numbers” of (Gorsky, Hawkes, Schilling, Rainbolt, 2020).
- Generalized Shuffle Theorem proof gives all generalized q, t Catalan numbers as coefficients of a single geometric series or “generating function.”

KR-homology (generalizes HOMFLY-PT polynomial)

- Rational Shuffle Theorem \implies formula for KR homology of (m, n) -torus knots. (Mellit, 2021)
- (Capran, González, Hogancamp, Mazin, 2024+) use Generalized Shuffle Theorem formula to give KR homology of associated “Coxeter knots.”

A Mystery: Convex Curve Conjecture

A Mystery: Convex Curve Conjecture

$D^{(C)} \in \mathcal{E}$ for any convex curve C between $(0, s)$ and $(r, 0)$.

A Mystery: Convex Curve Conjecture

$D^{(C)} \in \mathcal{E}$ for any convex curve C between $(0, s)$ and $(r, 0)$.

Conjecture (Blasiak, Haiman, Morse, Pun, S., 2023)

$D^{(C)} \cdot 1 = \sum_{\mu} a_{C,\mu}(q, t) s_{\mu}$ satisfy $a_{C,\mu}(q, t) \in \mathbb{N}[q, t]$.

A Mystery: Convex Curve Conjecture

$D^{(C)} \in \mathcal{E}$ for any convex curve C between $(0, s)$ and $(r, 0)$.

Conjecture (Blasiak, Haiman, Morse, Pun, S., 2023)

$D^{(C)} \cdot 1 = \sum_{\mu} a_{C,\mu}(q, t) s_{\mu}$ satisfy $a_{C,\mu}(q, t) \in \mathbb{N}[q, t]$.



A Mystery: Convex Curve Conjecture

$D^{(C)} \in \mathcal{E}$ for any convex curve C between $(0, s)$ and $(r, 0)$.

Conjecture (Blasiak, Haiman, Morse, Pun, S., 2023)

$D^{(C)} \cdot 1 = \sum_{\mu} a_{C,\mu}(q, t) s_{\mu}$ satisfy $a_{C,\mu}(q, t) \in \mathbb{N}[q, t]$.



- S_n -module?

A Mystery: Convex Curve Conjecture

$D^{(C)} \in \mathcal{E}$ for any convex curve C between $(0, s)$ and $(r, 0)$.

Conjecture (Blasiak, Haiman, Morse, Pun, S., 2023)

$D^{(C)} \cdot 1 = \sum_{\mu} a_{C,\mu}(q, t) s_{\mu}$ satisfy $a_{C,\mu}(q, t) \in \mathbb{N}[q, t]$.



- S_n -module?
- $s_{(1,\dots,1)}$ coefficient appears in context of “positroid varieties” (Galashin, Lam, 2024).

The Extended Delta Theorem

The Extended Delta Theorem (BHMPS, 2023)

For operators Δ_{h_l} and $\Delta'_{e_{k-1}}$ on symmetric polynomials,
$$\Delta_{h_l} \Delta'_{e_{k-1}} e_n = \sum (q, t \text{ monomial}) (\text{LLT symmetric polynomial})$$

Sum over “decorated Dyck paths.”

The Extended Delta Theorem

The Extended Delta Theorem (BHMPs, 2023)

For operators Δ_{h_l} and $\Delta'_{e_{k-1}}$ on symmetric polynomials,
$$\Delta_{h_l} \Delta'_{e_{k-1}} e_n = \sum (q, t \text{ monomial})(\text{LLT symmetric polynomial})$$

Sum over “decorated Dyck paths.”

- Originally conjectured by (Haglund, Remmel, Wilson, 2018).

The Extended Delta Theorem

The Extended Delta Theorem (BHMPs, 2023)

For operators Δ_{h_l} and $\Delta'_{e_{k-1}}$ on symmetric polynomials,
$$\Delta_{h_l} \Delta'_{e_{k-1}} e_n = \sum (q, t \text{ monomial}) (\text{LLT symmetric polynomial})$$

Sum over “decorated Dyck paths.”

- Originally conjectured by (Haglund, Remmel, Wilson, 2018).
- Shuffle theorem is a special case.

The Extended Delta Theorem

The Extended Delta Theorem (BHMPs, 2023)

For operators Δ_{h_l} and $\Delta'_{e_{k-1}}$ on symmetric polynomials,
$$\Delta_{h_l} \Delta'_{e_{k-1}} e_n = \sum (q, t \text{ monomial})(\text{LLT symmetric polynomial})$$

Sum over “decorated Dyck paths.”

- Originally conjectured by (Haglund, Remmel, Wilson, 2018).
- Shuffle theorem is a special case.
- Consequence: Schur expansion coefficients in $\mathbb{N}[q, t]$.

The Extended Delta Theorem

The Extended Delta Theorem (BHMPS, 2023)

For operators Δ_{h_l} and $\Delta'_{e_{k-1}}$ on symmetric polynomials,
$$\Delta_{h_l} \Delta'_{e_{k-1}} e_n = \sum (q, t \text{ monomial}) (\text{LLT symmetric polynomial})$$

Sum over “decorated Dyck paths.”

- Originally conjectured by (Haglund, Remmel, Wilson, 2018).
- Shuffle theorem is a special case.
- Consequence: Schur expansion coefficients in $\mathbb{N}[q, t]$.

Proof Technique

$$\Delta_{h_l} \Delta'_{e_k} e_n = \sum_{\alpha} D_{\alpha} \cdot 1$$

The Loehr-Warrington Conjecture

Loehr-Warrington Conjecture

$$s_{\lambda}^{(m,1)} \cdot 1 = \pm \sum (q, t \text{ monomial})(\text{symmetric function})$$

Sum over “nested Dyck paths”

The Loehr-Warrington Conjecture

Loehr-Warrington Conjecture

$$s_{\lambda}^{(m,1)} \cdot 1 = \pm \sum (q, t \text{ monomial})(\text{symmetric function})$$

Sum over “nested Dyck paths”

Rational Loehr-Warrington Theorem, (BHMPS, 2021+)

$$s_{\lambda}^{(m,n)} \cdot 1 = \pm \sum (q, t \text{ monomial})(\text{symmetric function})$$

Sum over “dens” in a “nest.”

The Loehr-Warrington Conjecture

Loehr-Warrington Conjecture


$$s_{\lambda}^{(m,1)} \cdot 1 = \pm \sum (q, t \text{ monomial})(\text{symmetric function})$$

Sum over “nested Dyck paths”

Rational Loehr-Warrington Theorem, (BHMPs, 2021+)

$$s_{\lambda}^{(m,n)} \cdot 1 = \pm \sum (q, t \text{ monomial})(\text{symmetric function})$$

Sum over “dens” in a “nest.”



Catalanimals

Common theme: $\mathcal{E} \curvearrowright 1 = \text{Combinatorial sum}$

Catalanimals


Common theme: $\mathcal{E} \curvearrowright 1 = \text{Combinatorial sum}$

Bridge:

Catalanimals

Catalanimals


Common theme: $\mathcal{E} \curvearrowright 1 = \text{Combinatorial sum}$

Bridge:  Catalanimals

- Catalanimal = symmetric multivariate Laurent series.
- Governed by 3 subsets of $R_+ = \{(i, j) \mid i < j\}$ and $\gamma \in \mathbb{Z}^n$.

Catalanimals

Common theme: $\mathcal{E} \curvearrowright 1 = \text{Combinatorial sum}$


Bridge:  Catalanimals

- Catalanimal = symmetric multivariate Laurent series.
- Governed by 3 subsets of $R_+ = \{(i, j) \mid i < j\}$ and $\gamma \in \mathbb{Z}^n$.

$$H(R_+, R_+, \{(1, 3)\}, (1, 1, 1))$$

Catalanimals

Common theme: $\mathcal{E} \curvearrowright 1 = \text{Combinatorial sum}$

Bridge:  Catalanimals


- Catalanimal = symmetric multivariate Laurent series.
- Governed by 3 subsets of $R_+ = \{(i, j) \mid i < j\}$ and $\gamma \in \mathbb{Z}^n$.

$$H(R_+, R_+, \{(1, 3)\}, (1, 1, 1))$$

$$= \text{Symmetrization of } \left(\frac{z_1 z_2 z_3 (1 - q t z_1 / z_3)}{\prod_{1 \leq i < j \leq 3} (1 - q z_i / z_j) (1 - t z_i / z_j)} \right)$$

Catalanimals

Common theme: $\mathcal{E} \curvearrowright 1 = \text{Combinatorial sum}$


Bridge:  Catalanimals

- Catalanimal = symmetric multivariate Laurent series.
- Governed by 3 subsets of $R_+ = \{(i, j) \mid i < j\}$ and $\gamma \in \mathbb{Z}^n$.

$$\begin{aligned}
 & H(R_+, R_+, \{(1, 3)\}, (1, 1, 1)) \\
 &= \text{Symmetrization of } \left(\frac{z_1 z_2 z_3 (1 - q t z_1 / z_3)}{\prod_{1 \leq i < j \leq 3} (1 - q z_i / z_j) (1 - t z_i / z_j)} \right) \\
 &= s_{111} + (q + t + q^2 + q t + t^2) s_{21} + (q t + q^3 + q^2 t + q t^2 + t^3) s_3
 \end{aligned}$$

Catalanimals

Common theme: $\mathcal{E} \curvearrowright 1 = \text{Combinatorial sum}$

Bridge:  Catalanimals

- Catalanimal = symmetric multivariate Laurent series.
- Governed by 3 subsets of $R_+ = \{(i, j) \mid i < j\}$ and $\gamma \in \mathbb{Z}^n$.

$$\begin{aligned}
 & H(R_+, R_+, \{(1, 3)\}, (1, 1, 1)) \\
 &= \text{Symmetrization of } \left(\frac{z_1 z_2 z_3 (1 - q t z_1 / z_3)}{\prod_{1 \leq i < j \leq 3} (1 - q z_i / z_j) (1 - t z_i / z_j)} \right) \\
 &= s_{111} + (q + t + q^2 + q t + t^2) s_{21} + (q t + q^3 + q^2 t + q t^2 + t^3) s_3
 \end{aligned}$$

Thank you!